

©Copyright 1955 by Medical Encyclopedia Inc., New York, N. Y.

**©COPYRIGHT UNDER THE INTERNATIONAL COPYRIGHT UNION. ALL RIGHTS RESERVED.
THIS BOOK IS PROTECTED BY COPYRIGHT. NO PART OF IT MAY BE DUPLICATED OR
REPRODUCED IN ANY MANNER WITHOUT WRITTEN PERMISSION FROM THE PUBLISHER.
PRINTED IN THE UNITED STATES OF AMERICA.**

Distributors outside U.S.A.: Interscience Publishers, New York-London

CONTENTS

Foreword

| | |
|---|---|
| <i>by Henry Welch, Ph.D., and Félix Martí Ibáñez, M.D</i> | 7 |
|---|---|

Chapter I

| | |
|--------------|----|
| Introduction | 13 |
|--------------|----|

Chapter II

| | |
|---|----|
| Mechanism of the Antibiotic Growth Effect | 19 |
| <i>The Influence of Environmental Bacteria on the Growth of Animals</i> | 21 |
| <i>Morphologic Changes Produced by Feeding Antibiotics</i> | 25 |
| <i>The Effects of Antibiotics on the Growth of Germ-free Animals</i> | 26 |
| <i>Effects on Intestinal Flora</i> | 29 |
| <i>Effect on Rumen Microorganisms</i> | 33 |
| <i>Antibiotic-tolerant Intestinal Microorganisms as a Source of Unidentified Growth Factors</i> | 34 |
| <i>Growth of Animals Receiving Hydrolyzed Penicillin</i> | 36 |
| <i>Effects of Prolonged Usage of Dietary Antibiotics</i> | 38 |

Chapter III

| | |
|---|----|
| <i>Effect of Feeding Antibiotics on the Requirement for Nutrients</i> | 43 |
| <i>Effects on Vitamin and Mineral Requirement</i> | 44 |
| <i>Effect on Protein Requirement</i> | 50 |
| <i>Effects on Nitrogen Metabolism</i> | 53 |

Chapter IV

| | |
|--|----|
| <i>Antibiotics in Animal Production</i> | 57 |
| <i>Antibiotics in Poultry Production</i> | 58 |
| <i>Antibiotics in Swine Production</i> | 61 |

| | |
|--|----|
| <i>Effect on Carcass Quality</i> | 64 |
| <i>Antibiotics in the Diets of Ruminants</i> | 65 |
| <i>Antibiotics in Beef Cattle Feeding</i> | 68 |
| <i>Antibiotics in Lamb Feeds</i> | 69 |
| <i>The Effect of Dietary Antibiotics on Reproduction in Animals</i> | 71 |
| <i>Egg Production in Poultry</i> | 77 |
| <i>Antibiotics and Egg Production in Hens with Chronic Respiratory Disease</i> | 78 |

Chapter V

| | |
|--|-----|
| <i>Physiologic and Other Effects of Dietary Antibiotics</i> | 81 |
| <i>Effects of Antibiotics on the Endocrine System</i> | 81 |
| <i>Antibiotics and Liver Physiology</i> | 84 |
| <i>Effect of Antibiotics on Blood Formation</i> | 88 |
| <i>Pharmacology of Antibiotics</i> | 89 |
| <i>Effects on Growth of Human Subjects</i> | 90 |
| <i>Miscellaneous Nutritional Effects</i> | 92 |
| <i>Effects on Guinea Pigs</i> | 93 |
| <i>Public Health Aspects of Feeding Antibiotics to Animals</i> | 94 |
| <i>Conclusions</i> | 95 |
| <i>Bibliography</i> | 97 |
| <i>Index</i> | 121 |

Foreword

One of the great economic developments in the field of antibiotics is the use of these drugs in feeds for the promotion of animal growth. During the past few years there has been a steady increase in the amount of antibiotics — largely penicillin, chlortetracycline oxytetracycline, and bacitracin — used for this purpose. In 1954 alone, some 490 000 pounds of these drugs were incorporated into feeds rendering the farmer tremendous savings. Because antibiotics stimulate the growth of swine, chicks, and poults, they cut down the feed consumption of these animals, since they can be sent to slaughter much sooner and in a healthier condition than they would have been prior to the use of these drugs.

A unique phenomenon — perhaps without precedent in the history of medicine — is indeed the present use of the so-called antibiotics as stimulants of growth and nutrition in both human beings and animals.

The opposite phenomenon has already been recorded in history namely the use of such nutritive elements as vitamins, mineral salts, and amino acids as therapeutic agents. Vitamins were known and developed primarily as a component of the normal diet of human beings and animals only recently were they synthesized and developed as powerful therapeutic drugs. But never before has a medicinal drug been used in animal feeding or in the diet of humans to stimulate nutrition and promote the speed and extent of growth.

This new application for antibiotics will surely not only greatly widen the horizon of their usefulness, but also markedly affect the nutritional and economic problems of the world probably so much so that antibiotics may in the future become a most valuable element in solving the ever-growing disproportion between increasing world population and food supplies.

The use of antibiotics in nutrition has also opened up new avenues for research into such basic philosophical problems of biology as the influence of environmental bacteria on animal growth the correlation between the biochemical action of antibiotics and the anatomical changes in the morphology of animals fed with these substances the philosophical question of "germ free" animal life versus the growth of animals whose bodies, as well as their environment, are literally soaked in bacteria, and finally the correlation between the prolonged administration of antibiotics to animals and their vitamin and mineral requirements.

These and other vital problems will eventually be solved thanks to the searching light projected by antibiotics and antibiotic research upon the field of nutrition.

We were indeed fortunate that Dr. Thomas H. Jules prepared the monograph on "Antibiotics in Nutrition." This is the fourth in the monograph series designed to embrace the principal antibiotics and their application, not only in clinical medicine but in other allied fields of usefulness as well. Doctor Jules is a pioneer in this field. He, together with his colleagues, was the first to demonstrate the effectiveness of chlor tetracycline in stimulating the growth of chickens.

In the preparation of this monograph, Dr. Jules has completely covered the development of the use of antibiotics in nutrition. Following a brief introduction in Chapter I, he discusses in some detail the mechanism of the antibiotic growth effect. He covers the effect of feeding antibiotics on the requirement for other nutrients in Chapter III. Chapters IV and V explain the use of antibiotics in animal production and the physiological and other effects of dietary antibiotics. A bibliography of more than 400 references is testimony to the thoroughness with which Dr. Jules has covered the subject of antibiotics in animal nutrition. All of the studies done to date concerning the antibiotic effect on the growth of human subjects, are

also included, and the public health aspects of feeding antibiotics to animals is explored.

The correlation of this large mass of data has been a monumental task, but it is so well done that this monograph will be of great value to public health and research workers, veterinarians, economic specialists, and those interested in the impact of the ever widening use of antibiotics in our daily life.

Doctor Jukes has written his monograph in a clear, terse, easily readable style. We consider it a very worthwhile and valuable addition to this series.


HENRY WELCH, Ph.D.
*Director, Division of Antibiotics,
Food and Drug Administration;
Editor-in-Chief, Antibiotics &
Chemotherapy and Antibiotic
Medicine*

FELIX MARTI IBANEZ, M.D.
*International Editor, Antibiotics &
Chemotherapy; Associate Editor,
Antibiotic Medicine*

"OFTEN in our laboratory discussions for the last several years I have told the young research workers in my group that there should be much interest in raising young animals such as rabbits, guinea pigs, dogs and chicks from birth on pure food materials. By this latter expression I refer to food products which have been freed completely from common microbes by laboratory methods. Without wishing to speak positively I wish to state that I would undertake this investigation if I had time with the expectation that life under these conditions would become impossible. By such an approach one could perhaps attempt to study the digestive process by the systematic addition of microbe-free nutritive materials or single strains of microbes or mixed cultures of known microbes. The egg of the chicken lends itself without serious difficulty to this type of experiment. The outside of the shell could be kept free from bacteria at the moment the chick hatched and then the chick could be placed in a germ free environment ventilated by sterile air and sterile food (water, milk and grains) could readily be introduced from the outside.

"Whether the result would be positive and would confirm the pre-conceived idea which I have put forward or whether it would be negative or even reversed, that is to say that life under these conditions would be actually stimulated, it would be of great interest to try such an experiment."

(PASTEUR, L., as quoted by Duclaux, E.,
Comptes rendus Volume 100, page 68, 1885)




CHAPTER I Introduction

The use of antibiotics in the control of harmful bacteria is a culmination of years of searching for what has been aptly termed a "magic bullet"¹ a substance that will kill a parasite without harming its host. Antibiotics when added to the food of animals are able to improve their nutrition by producing changes in the intestinal bacteria. This effect was discovered accidentally; the "magic bullet" reached an unseen target. The growth rate of young chickens was found to be increased by adding streptomycin,² chlortetracycline,³ or chlortetracycline fermentation residues^{4, 5} to their diet. The work with chlortetracycline residues led promptly to studies with other species and it was found, perhaps not unexpectedly that the growth promoting effect was even more marked when certain intestinal infections were present. It is now well established that the growth rate of young animals is usually increased by adding relatively small amounts of certain antibiotics to the diet. This effect on growth occurs in apparently healthy animals on diets that are adequate in all known nutritional factors and the effect is also observed on certain deficient diets.


Many antibiotics are toxic to animals but there are certain outstanding and familiar exceptions, which include three antibiotics that are widely used in animal feeds: chlortetracycline, oxytetracycline,[†] and penicillin.⁶ These three substances are well known for their effectiveness in animals against many disease-producing bacteria and for the absence of harmful effects produced by their use in therapeutic doses. They are commonly used in animal feeds, and bacitracin is used to some extent. These four antibiotics together with streptomycin have been used in many experimental nutritional studies since 1950 and this

¹ The trade name of Lederle Laboratories Division, American Cyanamid Co., for chlortetracycline is Aureomycin.


[†] The trade name of Chas. Pfizer & Co., for oxytetracycline is Terramycin.




Benzylpenicillin



Streptomycin




Chlorotetracycline



Oxytetracycline

FIG. 1. Structural formulas of penicillin G (benzylpenicillin) streptomycin, chlorotetracycline, and oxytetracycline.

monograph will be largely concerned with a review of these studies. The structural formulas of chlortetracycline, oxytetracycline, streptomycin, and penicillin are shown in figure 1. Their distinctive structural chemistry excites interest in the biochemical mechanism by which such widely differing substances inhibit bacterial growth while not injuring the host to which they are administered. The other antibiotic that is used as a supplement to feeds, bacitracin is a mixture of complex polypeptides. Bacitracin A has recently been reported by Craig and co-workers to have the structure



It is nephrotoxic when injected⁷ but appears to be absorbed very poorly from the gastrointestinal tract.

The antibiotics are a group of soluble organic substances that are produced by microorganisms and that are characterized by their property of inhibiting at low concentrations the growth, activity or multiplication of other microorganisms. Antibiotic like substances are also produced by the flowering plants and by animals. Such plants as garlic, hops, and tomatoes contain antibacterial compounds termed respectively "alline," "lupulon," and "tomatine," while lysozyme, a protein with antibacterial properties, is present in egg white and in the secretions of the eye. These substances are not included among the antibiotics because the definition of this term is restricted to compounds that are produced by microorganisms. However the extensive occurrence of antibacterial substances in more complex organisms serves to emphasize their widespread nature. This is a useful reminder of the fact that the introduction of small quantities of antibiotics into the diet of animals is not necessarily an entirely novel procedure.

The discovery and development of the useful antibiotics have been based on empirical screening procedures that were designed to find an antibacterial substance that was harmless to animals. It may be presumed that such a substance must block certain essential enzyme systems in the bacterial cell without appreciably affecting any of the essential systems in the cells of the host. The

mechanism of this biochemical selectivity has been studied from various stand points.^{8, 9} There may be differences in the biochemical mechanisms of bacteria as compared with animals so that it is possible to block the metabolism of a pathogenic microorganism without harming the host, or there may be differences in permeability so that an injurious substance can enter the bacterial cell but not the cells of the host.

Penicillin has been reported to exert its toxic effects on microorganisms only under conditions where their growth is possible.¹⁰ Gale⁸ found that penicillin inhibited the formation of combined glutamate in *Staphylococcus aureus* cells while peptide glutamate accumulated in the medium. Bacitracin was found to behave similarly to penicillin.¹¹ In contrast, chlortetracycline and chloramphenicol tended to inhibit the formation of the peptide rather than the uptake of glutamate.¹² Penicillin has also been found to produce changes in the metabolism of nucleic acid.¹³ Streptomycin appears to inhibit the oxaloacetate-pyruvate condensation in *Escherichia coli* and Umbreit et al¹⁴ have suggested that animal cells are not similarly affected because they are less permeable to the antibiotic. Other biochemical reactions in the bacterial cells were unaffected even by high concentrations of streptomycin.

It is evident from these and other examples that differences between the antibiotics exist with respect to their antibacterial mechanisms as well as to their chemical structures. The only known biochemical property common to the antibiotics under consideration is their antibacterial effect. They do not, for example, all possess a single and distinctive chemical group, which might have a growth promoting effect on animals. The strong inference is that any such effect is secondary to the antibacterial action of antibiotics on the microorganisms in the digestive tract.

Before the discovery of the growth-promoting effect of antibiotics on animals, many investigations were made of the results produced by adding sulfonamides to purified diets. It was found that under such conditions deficiencies of biotin, folic acid, and vitamin K were produced.¹⁵⁻¹⁹ However these deficiencies did not appear when these three substances were added to purified diets or when rats received diets of natural foods to which sulfonamides were added. The results were taken to indicate that rats were able to obtain a supply of biotin, folic acid, and vitamin K from a nondietary source when the rats were maintained on a purified diet that was deficient in these three vitamins and that this nondietary source originated in a bacterial fermentation that took place in the intestinal tract. It could be deduced that sulfonamides could depress or change the intestinal bacteria in a manner that interfered with the production of these vitamins, which are needed by the host animal. However

the sulfonamides do not interfere with the assimilation of vitamins that are added to the diet. The reason that deficiencies of biotin, folic acid, and vitamin K developed was because they were not added to purified diets for rats, since deficiencies of these vitamins had not appeared before sulfonamides were added. In contrast, thiamine, riboflavin, pyridoxine and pantothenic acid were added to the diets because if these additions were not made deficiencies developed even in the absence of sulfonamides.

The concept of a symbiotic relationship between animals and the bacteria in their gastrointestinal tract is an old one. The studies with rats on diets with sulfonamides emphasized this relationship and experiments with ruminants showed that these animals were even more versatile, for it was found that they could largely dispense with certain water-soluble vitamins, which were shown to be produced by fermentation in the rumen.²⁰ Furthermore simple nitrogenous compounds such as urea, can be transformed by the rumen microorganisms into amino acids, which could replace part of the protein requirement of ruminants.

In contrast with these beneficial activities of the microorganisms of the digestive tract, there are many familiar examples of pathogenic bacteria and protozoa that can invade the alimentary canal to produce clinical disease. The sulfonamides and antibiotics have made some of their greatest contributions to medicine in controlling enteric diseases produced by organisms of this type.

A few investigators in the 1940's described experiments with rats and chicks in which antibacterial substances produced increases in the growth rate of the animals. Martin²¹ found that the mortality of young rats was prevented and their growth rate was increased by feeding 1 mg. of sulfanilamide daily but it was not clear whether an infection was present. Moore and co-workers²² found that succinylsulfathiazole and streptomycin increased the growth of chicks on a purified diet. These authors made the important suggestion that the dietary supplements inhibited intestinal bacteria that were producing toxic materials or were rendering certain dietary vitamins unavailable to the animal, but that the possibility that the agents were acting systemically could not be overlooked. Morehouse and Mayfield²³ found that 3-nitro-4-hydroxyphenylarsonic acid increased the growth of chicks and turkeys when added to natural diets. This observation was made during studies of the coccidiostatic effect of this substance. A pharmacologic study of chlortetracycline by Harned and co-workers²⁴ led to an observation that the addition of this antibiotic to the diet increased the growth rate of chicks. The effect was attributed to the elimination of some infection. All these studies in which the growth rate of animals was improved by antibacterial substances appeared to indicate that the effect was

due to suppression of harmful microorganisms, and indeed such an effect would not be surprising in view of the association of weight loss with clinical disease. Nevertheless, the possibility had been suggested by Moore and co-workers² that unidentified intestinal bacteria could have an inhibitory effect upon growth without causing clinical disease.

Investigations in our laboratory which were first reported in 1949^{4,5} showed that a growth-promoting effect was produced in chicks when crude preparations of *Streptomyces aureofaciens* cultures were added to the diet. These experiments were carried out with chicks that were being used to assay the vitamin B₁₂ content of *S. aureofaciens* fermentations. The chicks were apparently healthy and were kept in cages that were frequently cleaned. Furthermore, the cages were sterilized with steam at four week intervals. Even with these precautions, the crude fermentation materials containing chlortetracycline produced more rapid growth than could be obtained with crystalline vitamin B₁₂ or liver extract. Samples of the fermentation materials, which had been examined in the chick growth test and had been found to contain the new growth factor were sent early in 1949 to other investigators including Catron, Couch, Cunha, Krider McGinnis, and Stingsen, who confirmed the observations during the summer of 1949 and who also showed that other species, such as turkeys and pigs, showed a marked growth response to the fermentation materials. The response was then found to be due to the presence of chlortetracycline in the fermentation materials.²³ Unexpectedly small amounts of chlortetracycline and other antibiotics were found to produce an effect on growth.²⁴⁻²⁷ These amounts were so small, 5 to 10 parts per million of diet, that it was economically feasible to incorporate them in commercial diets for farm animals.

The nature and mechanism of the growth-promoting effect produced in young animals by antibiotics has excited the interest and attention of many investigators and some of their findings will be discussed in succeeding chapters.

CHAPTER II Mechanism of the Antibiotic Growth Effect

The action of antibiotics in increasing the growth rate of young animals may be termed for convenience the "antibiotic growth effect." The most obvious explanation of the effect is to postulate that it is due to the action of antibiotics on the intestinal bacteria. In support of this are the following observations:

1. A number of different antibiotics will produce the effect on growth but they have no special chemical properties in common. Furthermore, their antibacterial effects are exerted through different mechanisms. It is therefore unlikely that these antibiotics would all have a stimulatory effect on animal cells and it is much more probable that the effect on growth is secondary to changes in the intestinal bacteria, since many of these bacteria are known to be susceptible to the common antibiotics.

2. There have been many observations that the antibiotic growth effect may diminish or disappear entirely when steps are taken to keep bacteria away from the animals. Exceptions to these findings have been noted. However, if the growth effect is due to the direct effect of the antibiotic upon the animals, it should be consistently obtainable in clean surroundings, which is not the case.

3. Very little is known of the normal effects of the intestinal bacteria on the nutrition of the host animal. For example, it has been shown that feeding antibiotics brings about an increase in the storage of vitamin A in the liver^{28, 29} and that a similar increase takes place when the animals are placed in a highly sanitary environment without antibiotics.³⁰ This may imply that the effect on Vitamin A is secondary to changes in the gut wall, which the antibiotic has produced by its action on the intestinal microflora; indeed, such findings may be a clue to hitherto unrecognized biochemical activities of the intestinal bacteria.

4. The growth-promoting effects of certain antibiotics when injected may be due to their excretion into the intestine. This has been shown to occur in

calves under conditions where growth was improved following of chlortetracycline.³¹

5. The antibiotic growth effect can be produced by certain antibiotics which do not readily enter the blood stream when given by mouth. The antibiotics streptomycin³² and bacitracin.³³ It is of interest that bacitracin has been found to produce a growth response in chicks when added to the diet³⁴ but when implanted subcutaneously³⁵ for injected bacitracin has been reported to be excreted in the urine³⁶ rather than the bile so that it is possible that bacitracin may not reach the intestine. In contrast, such antibiotics as tetracycline and penicillin, which have been found to produce growth responses when injected,^{31, 37, 38} are known to be excreted into the bile.^{31, 39}

6. A large number of investigations have shown that antibiotic treatment is a requirement for certain B vitamins in animals on diets deficient in these vitamins. The B vitamins differ from each other with respect to their chemical structure and physiology but they share the common property of being produced by intestinal bacteria, and there are indications that antibiotic-tolerant strains of these bacteria will produce increased amounts of B vitamins.⁴⁰

7. Antibiotics disappear rapidly from the tissues of animals after administration is stopped.⁴¹ However under certain experimental conditions the dietary effect of antibiotics in sparing vitamin B₁₂ may persist after their withdrawal from the diet,⁴² thus suggesting that a firmly established change in the intestinal flora has taken place.

8. Several investigators have reported that the growth of unsupplemented animals gradually tends to improve from experiment to experiment when other animals are receiving antibiotics, indicating the elimination of certain harmful bacteria from the environment.

9. No growth effect was found when antibiotics were injected into growing chick embryos⁴³ or were fed to germ free chicks⁴⁴ at levels that would be expected to affect the growth rate of "contaminated" animals.

10. The beneficial effect of antibiotics on dietary liver necrosis in chicks was found to disappear after several weeks of feeding, suggesting that the effect was due to a change in the intestinal flora, which later became tolerant to the antibiotic so that the change was reversed.⁴⁵

The Influence of Environmental Bacteria on the Growth of Animals

The effect of antibiotics in improving growth of animals led to the

sion that bacteria that inhibited growth might be normally present in the intestinal tract. If this assumption were correct, the inhibitory effect should be absent in animals that are free from bacteria, and such animals should not show a growth response to antibiotics. Evidence to test the assumption was sought from several approaches from the concept of "disease level," from comparing animals in "clean" and heavily contaminated surroundings, from measuring the growth of chick embryos, and from studying "germ free" chicks and turkeys.

The Disease Level Concept Experiments with *S aureofaciens* fermentation products in the diets of swine were carried out in 1949 and 1950 by a number of investigators including Cunha et al,⁴⁶ Catron,⁴⁷ Terrill and Krider⁴⁸ and Carpenter.⁴⁷ Striking effects were noted in the control of diarrhea and in the improvement of growth. In many herds, bloody diarrhea was endemic and yielded to treatment with the antibiotic supplement so that the affliction largely disappeared and growth was markedly increased. These observations led to an emphasis by various investigators on the postulation that the antibiotic growth effect in pigs varied with the disease level,^{47 49, 51} a term that was first introduced by Catron,⁴⁷ which he has defined as the degree or amount of feed lot contamination with bacteria or virus infestation that causes pigs to scour.⁴⁸ The definition could probably be extended to include other species of animals in addition to pigs and other pathologic manifestations in addition to diarrhea. This proposal is now well substantiated and it is supported by observations on other farm animals, such as turkeys, chickens, and calves. It seems evident that certain subacute infections can often occur in young animals and that these infections are accompanied by slow growth and often by diarrhea. Certain antibiotics will relieve these conditions when added to the diet so that a marked increase in growth rate is observed when antibiotics are administered. However it is also evident that a somewhat less marked effect of antibiotics on growth can be produced in animals that appear to be healthy as judged by a complete absence of external clinical signs. It is not clear whether the growth promoting effect of antibiotics on these apparently healthy animals differs only qualitatively from the effect of antibiotics on the growth of animals that are suffering from frank infections. However the preponderance of evidence is that both phenomena are part of the same picture and the effect of antibiotics in both cases is to eliminate harmful bacteria.

Relation of Clean and Dirty Environments to the Antibiotic Growth Effect The finding that a high disease level magnified the antibiotic growth effect in pigs leads to the possibility that the effect may disappear in animals if the bacterial contamination of the environment is kept low. A report by Coates and

to initiate the proliferation of a thriving and abundant microflora. Taken as a whole, the findings demonstrate that under some sanitary conditions, and perhaps by chance, it is possible to stop certain antibiotic-susceptible and growth-depressing microorganisms from becoming established in the intestinal tract of experimental animals. This is shown by the fact that these animals may grow as well without as with antibiotic supplementation until an outside source of infection is introduced. However the random tendency of natural contamination can lead to a different result so that animals that are kept scrupulously clean may nevertheless acquire and harbor a microflora whose growth-depressing action is eliminated by feeding an antibiotic. The experimental findings in which antibiotics were fed to animals in clean and dirty environments have led to the conclusion that the antibiotic growth effect may be due to the suppression of a growth-depressing microflora that is usually present in healthy animals but that may sometimes be absent under special sanitary conditions.

Antibiotics and the Growth of Chick Embryos The developing chick embryo grows rapidly in the absence of an intestinal flora and its weight increases twenty-six fold during the period between the seventh and seventeenth days of incubation.⁶² This is a greater rate of increase than takes place at any time after hatching. We concluded that a living organism whose tissues were prolif

Table III

Weight of Chick Embryos at 17 Days as Affected by
Injections of Antibiotics

| Antibiotic* | Amount Injected, mg | Survival ratio | Average weight | |
|------------------------|---------------------------|-------------------|-------------------|---------------|
| | | | Whole egg, Gm. | Embryo Gm. |
| Chlortetracycline | 1.2 | 19:30 | 54.1 | 12.0 |
| Penicillin | 1.2 | 20:30 | 53.7 | 12.6 |
| | 6.0 | 22:30 | 53.6 | 10.4 |
| Streptomycin | 1.2 | 20:30 | 54.6 | 12.5 |
| | 6.0 | 23:30 | 54.2 | 12.2 |
| None (distilled water) | | 26:36 | 53.7 | 13.3 |

The antibiotic was divided into three equal doses, which were injected on three successive days.

erating so rapidly might be unusually sensitive to the effects of a growth-stimulating substance. Accordingly chicken eggs containing living embryos were injected with sterile solutions of antibiotics and the incubation period was continued. The injections were started on the seventh day and on the seven-teenth day the embryos were separated from the membranes and weighed.⁶⁴ No growth-promoting effect of the antibiotics was found, and no evidence of toxicity as judged by deformities or depression of growth was observed except that the embryos on the higher level of penicillin were somewhat smaller than the others (table III). The results indicated that the antibiotics tested did not increase the growth rate of embryonic tissue and the observation helps to support the idea that the antibiotic growth effect is exerted through the intestinal flora, since these are absent from the developing chick embryo. Lepine and co-workers⁶⁵ found that slight but consistent diminutions in the proliferation of chick embryo heart and lung fibroblasts and epithelial cells in vitro were caused by chloramphenicol or chlortetracycline, 10 μ g/ml. of culture medium. Higher levels of the antibiotics produced marked inhibitory effects on the growth of these tissue cultures. In experiments with brain tissue, 10 μ g./ml. of chloramphenicol and 100 μ g./ml. of chloramphenicol or chlortetracycline inhibited cell migration while 10 μ g./ml. of chlortetracycline had no effect.

Morphologic Changes Produced by Feeding Antibiotics

Gordon⁶⁶ made a number of comparisons of the morphology of germ-free chicks with chicks exposed to ordinary contamination, with and without antibiotic feeding. Penicillin and oxytetracycline were the antibiotics that were used. A marked reduction, 33 to 50 per cent, of the weights of the small intestine was noted in the germ free animals and in the antibiotic fed conventional animals, by comparison with the conventional nonantibiotic-fed animals. Simultaneously there was a less marked increase in the weight of the thymus gland. No difference between the antibiotic-fed and nonantibiotic-fed germ-free animals was observed with respect to the weights of these organs. The author concluded that antibiotics created "germ-free-like conditions in the conventional animals.

It is indeed striking that without reducing the total bacterial count in the intestine of chicks the antibiotics will produce morphologic conditions similar to those in germ-free animals. This observation points strongly to the probability that antibiotic-sensitive bacteria, which produce a tissue reaction, are normally present in the digestive tract.

Table V

Summary of Results with Germ-free Animals In which the Weights at 4 weeks with and without Antibiotic Supplementation Were Compared

| Antibiotic | Level per Kg of diet mg | Animals | No. of experiments showing | |
|-----------------|-------------------------|---------|----------------------------|---------------|
| | | | Increased wt | Decreased wt. |
| Streptomycin | 45 | Chicks | 0 | 1 |
| Streptomycin | 70 | Chicks | 1 | 0 |
| Penicillin | 11 | Chicks | 1 | 0 |
| Penicillin | 11 | Turkeys | 1 | 0 |
| Penicillin | 46 | Chicks | 1 | 1 |
| Penicillin | 46 | Turkeys | 3 | 2 |
| Oxytetracycline | 25 | Chicks | 3 | 0 |
| Oxytetracycline | 25 | Turkeys | 0 | 1 |
| Oxytetracycline | 50 | Chicks | 1 | 3 |

Taken Luckey and co-workers^{44, 71}

somewhat variable but the over-all experience with these two drugs indicated a negative response. The levels of antibiotics in the diet ranged from 23 mg. of chloramphenicol to 50 mg. of oxytetracycline per Kg. of diet and the number of chicks per experimental group was necessarily small, ranging from 2 to 6. Similar results were obtained with turkeys which received 50 mg. of penicillin per Kg. of diet so that the author concluded that "no stimulation is seen when germ-free birds are fed antibiotics."

More recently Luckey⁷¹ found that in three out of four experiments with a lower level of oxytetracycline, 25 mg./Kg. of diet, the birds receiving the antibiotic grew more rapidly than the unsupplemented birds and slight increases in the growth rate, 3 to 7 per cent greater than the controls, were noted in two experiments in which a lower level of penicillin, 11 mg./Kg. of diet, was fed while in three experiments with turkeys receiving 46 mg. of penicillin per Kg. of diet, the weights of the supplemented birds were respectively 1, 7 and 13 per cent higher than those of the controls. He inferred that the difference between these and the earlier results was that the levels of oxytetracycline

and penicillin used in 1952 were higher and therefore growth-depressing. However the small number of birds involved and the great variability in the individual weights make it impossible to arrive at a definite conclusion that a growth effect was observed at these lower levels of antibiotics. It seems unlikely that levels of 50 mg of oxytetracycline or penicillin would have a specific depressing effect on the growth of germ free birds when much higher levels of these antibiotics do not slow the growth of birds under ordinary conditions. The variability encountered in these experiments may be seen in table IV in which the weights of all the germ-free birds receiving oxytetracycline as described by Luckey are summarized.

A condensed summary of the results described in Luckey's two communications on antibiotics and the growth of germ free birds is given in table V. Table V shows that the antibiotic-fed birds were heavier than the controls at 4 weeks in 11 of the 19 experiments. This would seem to lead to the conclusion that there was about an equal chance of the antibiotics increasing or decreasing growth, and the over-all result appears to confirm the conclusion reached by Luckey in 1952.

Effects on Intestinal Flora

Many studies have been made of the bacteriology of the intestinal contents of animals fed antibiotics in efforts to throw some light upon the antibiotic growth effect. The results have varied greatly and Sieburth and co-workers⁷³ pointed out that variations may be due to different basal diets, different bacteriologic procedures, the age of the birds studied and the types and levels of antibiotics used. Furthermore, their investigations showed that extreme variations in the cecal and fecal microflora were found even from day to day.

Examples of the differences noted by various investigators may be seen in table VI. These differences are so numerous that we hesitate to review the literature in detail or to attempt to draw generalizations with respect to the various bacterial types. Thus Kratzer and co-workers⁷² found a five- to ten fold increase in the number of yeasts in the excreta of chicks and turkey poultz fed streptomycin, while Williams and co-workers⁷⁴ found little or no change in the yeast count of chicks fed chlortetracycline. Sieburth and co-workers⁷⁵ drew attention to the marked reduction in intestinal clostridia produced by feeding antibiotics and this was confirmed by Elam and co-workers⁷⁶ and by Williams and co-workers.⁷⁴ However Williams and co-workers found that feeding live cultures of hemolytic toxin producing clostridia did not reduce the growth of chicks that were not receiving antibiotics, and no growth-depressing effect was

found when cell-free clostridia toxins were fed to unsupplemented chickens. Elam and co-workers⁷⁸ found that feeding fecal clostridia depressed the growth of chicks in clean quarters but not in old quarters. In later communication, Sieburth and co-workers⁷⁹ concluded that little significance could be attached to the observation that the toxin-producing clostridia are greatly reduced by antibiotic supplementation, because the population of these organisms is small in comparison with other types. These authors noted that a consistent reduction in the numbers of bacteria was found only in the material obtained in the small intestine and not in other regions of the digestive tract. They studied the oxygen uptake and carbon dioxide evolution of the intestinal contents in vitro, and they found that these values were lowered in the case of chicks fed a supplement of chlortetracycline, perhaps indicating a decreased use of nutrients by the microflora, which might permit greater utilization of nutrients from the diet by the host animal. The mesenteric blood vessels along the small intestine were noted to be more dilated in the birds receiving the antibiotic supplemented diet.

Smyser and associates¹⁰³ found that dietary penicillin produced during 4 weeks a steady increase in the numbers of *Clostridium perfringens* in the feces of chicks. This observation seemed to rule out the possibility that the growth effect was related to a suppression of *Cl. perfringens*. Chlortetracycline produced only slight changes in the count.

Another source of difficulty in carrying out investigations of the effects of dietary antibiotics on intestinal bacteria is because the presence of the antibiotic in intestinal contents may prevent the growth of the bacteria which are plated out.¹⁰⁴

Effect on Rumen Microorganisms

A number of investigators have made observations on the effect of antibiotics on the rumen microflora. Loosli and co-workers¹⁰⁰ did not observe any striking difference in the total bacterial count or in the morphologic type of bacteria present based on stained slide examination in studies with Holstein calves fed chlortetracycline. In experiments with beef heifers, Neumann and co-workers¹⁰⁶ stated that the types of organisms found in the animals receiving chlortetracycline were much less diverse than in the controls, which suggested that the normal bacterial flora had been disturbed. Keeler and Knodt¹⁰⁷ found that rumen samples taken from calves fed oxytetracycline showed consistently lowered ability to digest cellulose in vitro as compared with the samples from control calves. These differences disappeared when the antibiotic supplement-

of indicator microorganisms in the rumen when chlortetracycline was fed to dairy calves. The supplemented animals were superior to the controls with respect to gains in body weight, feed intake, and efficiency of feed utilization. These investigators and Kesler and Knodt reported that the antibiotic supplementation had no significant effect on the thiamine and riboflavin levels of the rumen contents although Chance and co-workers¹⁰⁹ found that the amount of riboflavin in the rumen contents was lower when 0.5 Gm. of chlortetracycline was fed daily than when the antibiotic-free ration was fed. In another investigation Chance and co-workers¹¹⁰ reported that a definite increase occurred in the total bacterial count of the rumen contents and feces when chlortetracycline was included in the ration. The number of rumen streptococci decreased in the two fistulated animals that were used in the study. Horn and co-workers,¹¹¹ in studies with yearling steers, reported that the rumen contents of animals receiving chlortetracycline showed a decreased ability to digest fiber in vitro and similar observations were made by Radisson and co-workers¹¹² however these investigators noted that, when alfalfa hay was added to the culture medium, no depressing effect of chlortetracycline on cellulose digestion occurred, indicating that alfalfa hay contained a factor that prevented chlortetracycline from depressing the digestion of cellulose in vitro by cultures of rumen bacteria. It is evident that studies of cellulose digestion in vitro do not necessarily reflect the behavior of the rumen bacteria in the living animal. Rusoff et al¹¹³ did not find differences between control calves, chlortetracycline injected calves, and those fed chlortetracycline orally with respect to the rumen contents. Measurements were made of pH and the content of eight of the B vitamins in the rumen fluid together with an examination of the rumen flora and the degree of fiber digestion in vitro. Lodge and co-workers¹¹⁴ found that the digestion of cellulose in vitro by the rumen microorganisms was lower in the case of cows that had received a daily supplement of 240 mg of chlortetracycline from an early age. The presence of chlortetracycline was demonstrated in the rumen liquid. These authors also noted that the addition of 1.6 μ g of chlortetracycline per ml. of fermentation mixture severely inhibited cellulose digestion in vitro by the microorganisms from the control cows but had very little effect on the digestion of cellulose by the microorganisms obtained from the antibiotic-fed cows perhaps indicating the establishment of chlortetracycline-tolerant cellulolytic microorganisms in the antibiotic-fed animals.

Huhtanen and co-workers¹¹⁵ have described techniques for studying the digestion of fiber in vitro by rumen microorganisms. Depressions of the digestion was discontinued. Hibbs and co-workers¹⁰⁸ found changes in the ratings

tion of fiber were observed when antibiotics were added. However no depression of the digestion of cellulose as present in alfalfa was observed¹¹⁶ to result from feeding chlortetracycline to sheep thus recalling the results reported by Radiason.¹¹⁵

Studies of the effect of chlortetracycline on the rumen microorganisms were reported by Mann and co-workers.¹¹⁷ The antibiotic was fed in doses of 40 to 60 mg to young calves for several weeks following which the animals were slaughtered and the contents of their paunches and abomasums were examined. The chlortetracycline-fed calves showed no marked bacteriologic differences from the controls and the authors commented that there was little difference between the supplemented and control calves in the final development of a typical viable streptococcal population of the rumen. The supplemented calves had larger and less acid rumen contents than the controls and reached a rumen pH that was considered to be suitable for intensive rumen bacterial and protozoan action at a much earlier age.

Antibiotic tolerant Intestinal Microorganisms as a Source of Unidentified Growth Factors

Following the addition of an antibiotic to the diet, microorganisms that are tolerant to the presence of the antibiotic in culture media may be recovered from the intestinal contents.¹¹⁸ One possible explanation of the antibiotic growth effect in animals could be that these tolerant microorganisms produce vitamin like growth factors that are not present in sufficient quantities in the diet to enable maximum growth of the host animal to take place and that are not produced by the intestinal flora at a sufficient rate in the absence of antibiotics in the diet. There have been several investigations of this possibility. Kratzer and co-workers¹¹⁹ found that the number of yeasts increased five- to tenfold in the intestinal tract of young turkeys that received streptomycin. These yeasts were plated out and grown in large flasks following which they were harvested and fed to chicks and turkeys. There was no consistent growth response and, although slight increases were found in some experiments, they were less than those obtained with antibiotic supplements.

A fresh attack on this problem was made by Romoser and co-workers¹²⁰ who noted that *E. coli* and *Aerobacter aerogenes* increased in numbers in the ceca of penicillin fed chicks. These penicillin resistant organisms were isolated, grown in mass cultures under various conditions, and fed to chicks. In some experiments the cultures were lyophilized. The chicks received basal diets that were considered suitable for obtaining responses to unknown growth

factors." Little or no chick growth response was obtained when either organism was added to the diet in the absence of an antibiotic but a significant growth response was obtained to penicillin alone and a greater response to the combination of penicillin and cultures of the organisms as dietary supplements. It was considered in view of the small amount (0.22 per cent) of the supplement that the results were due to effects of the viable organisms rather than to unidentified growth factors in the cultures, and the possibility was raised that the organisms "aided" the antibiotic in reducing the number of inhibitory intestinal bacteria. However no experiments with heat inactivated cells were included.

Along similar lines Anderson and co-workers^{11 12 13} isolated various strains of *E. coli* from penicillin fed chicks and fed the cultures to other chicks. One strain of *E. coli* produced a growth response in the presence of penicillin but not in its absence while a second strain somewhat improved growth under both conditions. In another experiment only one out of five supplements, the culture filtrate, increased the growth rate of the chicks and then only in the presence of penicillin. Similar preparations were tested with turkey poult¹⁴ but no significant weight increases were found except that responses were obtained to penicillin itself. In a further report¹⁵ the conclusion was drawn that the growth response of chicks to antibiotics was associated with an increase in intestinal coliforms because the growth response and the increase took place simultaneously. Roberts¹⁶ has reported that the growth of chicks was increased by either *E. coli* or fish solubles and Blaylock and co-workers¹⁷ found that either of these supplements produced a growth increase of 11 per cent in chicks during the first three weeks. The control diet for such experiments ought to be supplemented with the unfermented culture medium, for the medium may contain supplements that improve the diet. The comparatively small growth responses that are obtained with experimental animals when sources of "unknown factors" such as bacterial and yeast cultures, are added to the diet present certain difficulties of interpretation. Much depends on the basal diet that is used to evaluate "unknown growth factors" in materials such as fish solubles, liver yeast, and distillers solubles. The diet should be complete and balanced with respect to the known growth requirements of the animal, including amino acids, vitamins, and minerals. Recently Scott et al¹⁸ and Couch and co-workers¹⁹ have found that chicks will show a growth response to the ash of certain sources of unknown growth factors and similar results have been obtained in our laboratory. There is no evidence that antibiotic-adapted forms of *E. coli* or other intestinal bacteria are better sources of nutritional factors than those bacterial forms that normally inhabit the intestinal tract.

The results with *E. coli* and other intestinal organisms when fed to either antibiotic-supplemented or unsupplemented animals indicate that cultures of these organisms will not consistently produce a growth response when added to the diet. Furthermore, when feeding the culture produces a growth response in the absence of antibiotics in the diet, the response may be less than the response to adding an antibiotic. It is possible that antibiotic-tolerant strains of intestinal organisms may displace harmful bacteria from the digestive tract. The investigations just cited^{125, 126} probably indicate that beneficial nutritional effects may be produced by antibiotic-tolerant intestinal organisms.

Growth of Animals Receiving Hydrolyzed Penicillin

The effect of penicillamine (β β -dimethylcysteine) on rats fed a low-choline diet was described by Wilson and duVigneaud.¹²⁷ They found that the L isomer was toxic and its injurious effects were counteracted by choline and ethanolamine. The D-isomer which is a hydrolytic product of penicillin, was not toxic. Experiments in our laboratories with chicks showed only a slight growth response to an enzymatic hydrolyzate of penicillin (table VII). The response appeared to be not more than 10 per cent of that given by the unhydrolyzed antibiotic. L-penicillamine was found to depress growth markedly when added at a level of 3 Gm./Kg. to a purified diet containing choline.

Elam and co-workers¹²⁸ reported that penicillin, inactivated by autoclaving, produced a growth response when injected into chicks but not when added to

Table VII

Effect of Procaine Penicillin and Penicillinase-Hydrolyzed Penicillin, on the Growth of Chicks

| Supplement per Kg of diet Gm | Weights at 25 days, Gm. | |
|---------------------------------|----------------------------|-----|
| None | 226, | 229 |
| 2 mg. procaine penicillin | 329 | 324 |
| 5 mg procaine penicillin | 306, | 325 |
| 5 mg hydrolyzed penicillin | 277 | 282 |
| 10 mg hydrolyzed penicillin | 278, | 294 |

There were 10 chicks in each of duplicate groups.

Table VIII

Microbiologic and Chick Assays of Penicillin in Feed

| Treatment of feed | Per cent of penicillin remaining | |
|--------------------------|----------------------------------|--------------|
| | Microbiologic assays | Chick assays |
| Steamed 5 minutes* | 43 | 64 46 |
| Steamed 15 minutes | 21 | 28 56, 50 14 |
| Steamed 30 minutes | 18 | 20 20 32 30 |
| Stored 2 weeks at 34 C. | 23 | 29 |
| Stored 4 weeks at 34 C. | 12 | 19 |
| Stored 4 weeks at 34 C.† | 48 | 67 |
| Stored 4 weeks at 34 C.† | 17 | 40 |

Dibenzylethylenediamine penicillin.

†Procaine penicillin two different samples.

the diet. An examination of their results shows that the response with the injected hydrolyzate was obtained in one experiment but not in a second experiment. Fell and Stephenson¹²⁹ reported that DL-penicillamine produced a growth response when injected but not when added to the diet however in their experiments penicillin did not produce a growth response when added to the diet. Taylor and Gordon¹³⁰ carried out 3 experiments with pigs in which penicillin inactivated by three different methods was compared with active penicillin for its growth-promoting effect on pigs. The pigs receiving penicillin grew at a rate of 14.3 per cent greater than the controls and with inactive penicillin the growth rate was 9.4 per cent more than the controls. No penicillin was detected in the gut, serum, or urine of the experimental animals. Williams and co-workers¹³¹ found that inactivated penicillin retained not more than a small part of the ability of the original antibiotic to promote the growth of chicks. It was found possible to plot a dosage response curve to levels of penicillin on a purified sucrose-casein diet so that feeds containing penicillin could be assayed with chicks. Chick diets containing 25 or 50 mg./Kg of penicillin were stored under various conditions or steamed and were assayed microbiologically and with chicks. The rates of inactivation of penicillin were

very similar when measured by either procedure as shown in table VIII. The chick assays gave slightly higher values but there were rather wide variations in the chick assays. The conclusion from the results in table VIII is that the hydrolyzate did not have more than a small portion of the activity of the original penicillin when tested with chicks. It is possible that hydrolyzed penicillin, even though inactive for the test organism *Micrococcus pyogenes* var *aureus*, might retain some slight activity against some of the intestinal bacteria. Recently we have compared the growth response to penicillin and D-penicillamine with chicks on the sucrose-casein diet and D-penicillamine was found to produce a growth response corresponding to about 3 per cent of that produced by procaine penicillin.

Effects of Prolonged Usage of Dietary Antibiotics

Reduction of the Antibiotic Growth Effect Following the Prolonged Use of Antibiotics in Diets When antibiotics are routinely used in the diets of animals, one can speculate that prolonged use, year after year might produce selective changes in the environmental bacteria so that antibiotic-resistant forms would gradually become more common. If some of these resistant bacteria were pathogenic or harmful, then one would expect that the antibiotic growth effect would disappear and that animals would grow at a rate corresponding to the rate seen in similar animals before the introduction of antibiotic feeding.

Actually the recorded observations have been in the opposite direction and the prolonged use of antibiotics has often resulted in a gradual improvement in the growth rate of unsupplemented animals in quarters where antibiotics have been routinely used for several years. Perhaps certain growth-depressing bacterial forms tend to disappear from the environment so that the increased growth rate of animals is eventually seen in both the supplemented and unsupplemented groups. It is also possible that antibiotic-resistant bacteria may produce increased quantities of certain unidentified growth factors. The possibility of an environmental clean-up due to antibiotic feeding came to our attention in 1949 and 1950 when the response to antibiotics in chicks in our laboratory appeared to diminish over a long series of experiments, although the chicks receiving antibiotic supplements continued to reach a weight of 300 Gm. or more at 25 days.

Waibel and co-workers¹²³ compared the weights of chicks at 8 weeks of age with the percentage growth response of chlortetracycline or penicillin in a series of experiments lasting from August, 1950 to July 1953. The quantitative response to antibiotics tended to vary inversely with the growth of the control




FIGURE 2

groups and the average weight of the control groups in the third year of the study was greater than during the first two years. A wide variety of diets was used. A scatter diagram of these results is in figure 2.

Lisby and Schaible¹²² summarized their findings with 3900 chicks in 146 comparisons between groups of birds that were fed antibiotics or arsenic acids and groups that received the unsupplemented control diets. The results are shown in table IX. A steady increase in growth rate and a decrease in mortality of the unsupplemented birds were noted. The authors pointed out that the results do not support a concept that the continued use of antibiotics in feeds can lead to the proliferation of resistant strains of organisms that can cause antibiotics to become ineffective. Instead, the authors state, the continuous use of antibiotics over a long period "produces an environment with a lowered germ load or disease potential. After long time usage these growth promotants

Table IX

Growth Responses to Dietary Antibiotics and Arsonic Acids in Successive Years Compared with Control Group in Experiments with Chickens

| Period | Increase in weight of supplemented birds by comparison with control birds, % | | Increase in weight of control birds with respect to 1950-1951 controls, % | Mortality to 6 wks. of age % |
|-----------|--|---------------|---|------------------------------|
| | Antibiotics | Arsonic acids | | |
| 1950-1951 | 19.0 (75)† | 16.0 (2) | 0 | 8.5 |
| 1951-1952 | 12.8 (32) | | 7.4 | 8.2 |
| 1952-1953 | 10.0 (4) | 3.8 (5) | 15.3 | 4.6 |
| 1953-1954 | 3.3 (19) | 3.2 (9) | 19.1 | 2.8 |

Taken from Libby and Schaible ¹⁰

†Number of comparisons in parentheses.

are still capable of suppressing the germ load, but less of a challenge is presented. Thus, birds that are grown on the premises where birds have or have had antibiotics or arsonic acids in their feeds benefit to some extent from the improvement in environment.

"The reduction in the mortality of chicks that we have observed during the last 4 yr adds support to the postulation that a reduced germ load develops. As can be seen in column 5, prior to the inclusion of antibiotics and arsenicals in chick rations, the mortality of battery-raised chicks up to 6 wk of age was about 8.5 percent, whereas the mortality of birds in these same batteries in recent years has declined to less than 3 percent. In addition, fewer cull chicks and less variability in growth rate within pens are now being noted.

"From the practical point of view this study indicates that antibiotics and arsonic acids would be of greatest value in promoting growth (i) when first used in a poultry-feeding enterprise and (ii) under the environmental conditions usually surrounding farm feeding. If once used and then discontinued, feeding performance may still be somewhat improved, because the contamination level has been lowered."

Experiments with pigs have also shown that the growth of unsupplemented controls may tend to improve from year to year in surroundings where antibiotics are fed, thus Catron has found in an unpublished summary of a five

year study (1949 to 1953) involving 690 pigs that the average daily gains of antibiotic-fed pigs in successive years were 1.64, 1.60, 1.60, 1.67 and 1.57 lb., while the corresponding figures for unsupplemented pigs were 1.33, 1.44, 1.58, 1.57 and 1.57 lb. Hanson and Ferrin¹³⁴ obtained the following results in successive years with preweaning pigs kept from weaning to 200 lb. in the dry lot.

The basal diet was changed somewhat between 1951 and 1953 so that the results from one year to the next were not directly comparable however the results indicated that the general use of antibiotics for three years had controlled certain intestinal infections so well that these infections were not a problem in 1953. A similar trend is indicated in a report by Richardson¹³⁵ who used individually fed littermate pigs in four successive experiments each of which lasted between 14 and 21 weeks. The average daily gain in pounds of the unsupplemented controls rose from 1.38 in 1952 and 1953 to 1.61 in 1954 while the corresponding groups that received chlortetracycline gained respectively 16, 20.9 and 4 per cent more than the controls in successive experiments thus showing a decrease in the antibiotic growth response due to improvement by the unsupplemented controls. No response was obtained to oxytetracycline.

These experiments with chickens and pigs in which the antibiotic growth effect was shown to diminish in a series of experiments in the same rearing quarters while the growth of the unsupplemented controls improved are of great interest. The results indicate that the antibiotic growth effect is not due to a direct action on the animal but is due to a suppression of harmful bacteria and that the antibiotics or their effects may become "spread around the animal quarters so that the suppression may gradually extend throughout the

Table X

Results in Successive Years with Preweaning Pigs Kept from Weaning to 200 lb. in the dry lot¹³⁴

| Year | Average daily gain lb. | | | |
|------|------------------------|-----------------|--------------------|-----------------|
| | 3 to 8 weeks | | Weaning to 200 lb. | |
| | Without antibiotic | With antibiotic | Without antibiotic | With antibiotic |
| 1951 | 0.37 | 0.64 | 1.40 | 1.63 |
| 1952 | 0.43 | 0.71 | 1.45 | 1.63 |
| 1953 | 0.63 | 0.73 | 1.74 | 1.79 |

premises. An additional possibility is that certain antibiotic-tolerant bacteria, which become established through the use of diets containing antibiotics, may have a beneficial effect on nutrition. Clearly it is time to question the soundness of the dire prediction that the prolonged feeding of antibiotics would encourage the widespread development of resistant strains of dangerous bacteria¹³⁶ in the poultry rearing industry. From the practical standpoint, the effects of feeding antibiotics to growing farm animals for prolonged periods have led to unexpected benefits in the elimination of certain endemic types of undiagnosed morbidity. Slinger and Pepper¹³⁷ noted that syndrome involving poor growth, high mortality dermatitis, and diarrhea in young turkeys was overcome by the use of antibiotics and, furthermore, that this syndrome, which was one of the major problems involved in rearing poultz in Ontario, has largely disappeared following the inclusion of antibiotics in commercial diets.

The continuous administration of streptomycin and oxytetracycline at high levels (0.02 and 0.04 per cent) in the diet of aged rats, beginning at the age of 11 months, was not found to be beneficial as regards longevity in studies by French et al.¹³⁸ The average life span of the supplemented animals of both sexes was decreased by about 10 per cent. It was thought that, by analogy with their results on a high fat diet, an increased efficiency of food utilization might have been responsible for the deleterious effect. However an examination of the results indicates that the difference in the life span between treated and control animals did not reach significance at the 5 per cent level. No beneficial or deleterious effects of the antibiotic supplementation were shown by gross and histologic examination.

Effect of Feeding Antibiotics

CHAPTER III On the Requirement for Nutrients

Improvements in the utilization of protein and of certain vitamins and minerals have been noted to be associated with the feeding of antibiotics in some experiments. These improvements are separate from the antibiotic growth effect, which has been obtained on diets that were adequately furnished with the known requirements for growth and in some cases also with additional amounts of certain vitamins and sources of "growth factors" such as liver.⁵

Antibiotics have been shown in many investigations to enhance the utilization of certain nutrients, shown by a lower requirement for the nutrient for the production of maximum growth in the presence of the antibiotic, by a greater growth response to the antibiotic at suboptimal levels of the nutrient than when an optimal level of the nutrient is supplied, by a greater growth response to a small increment of the nutrient in the presence of the antibiotic than in its absence, by increases in the tissue levels of the nutrient, or by its increased urinary excretion or deposition in the egg when the antibiotic was fed. It appears in the case of some of the vitamins that antibiotics improve their uptake from the gut or their synthesis by the intestinal bacteria, or both. It also appears that the uptake of calcium and manganese may be improved by antibiotics.

In the case of the protein requirement, the situation is more obscure. In some investigations it has appeared that the protein requirement or the requirement for a specific amino acid for maximum growth was lowered by adding antibiotics and in others no protein sparing effect was noted. These differences may reflect variations in the intestinal bacteria of the animals in the various experiments.

There is a large amount of experimental evidence that establishes an effect of antibiotics in improving the efficiency with which food is converted to an

increase in body weight in growing animals. This effect may be an indication of an improvement of the uptake of various nutrients from the intestinal tract and this in turn can lead to measurable effects of dietary antibiotics on the requirement for protein, vitamins, and energy depending upon the manner in which the postabsorptive utilization of the nutrients takes place.

Effects on Vitamin and Mineral Requirements

B Complex Vitamins Emerson and Smith¹³⁹ produced a biotin deficiency in rats by adding streptomycin to a purified diet. This deficiency can also be produced by adding sulfonamides to purified diets for rats. Streptomycin appears to differ from some of the other antibiotics in this effect. In 1948 we made an unsuccessful attempt to produce folic acid deficiency in rats on a purified diet by adding a high level of chlortetracycline.

The first preparations of antibiotics that were widely used in animal feeds contained chlortetracycline and vitamin B₁₂ and the response that this combination produced in animals on "all vegetable" diets led to the wide impression that there was some interrelation between the antibiotic and vitamin B₁₂ in the growth response. In some investigations it was shown that an antibiotic supplement had a sparing effect upon the vitamin B₁₂ requirements of chicks or pigs but in other experiments the growth response to the antibiotic and the vitamin appeared to be independent of each other. Sparing effects on other vitamins were reported by many investigators.

Common and co-workers¹⁴⁰ found that blood levels of riboflavin and calcium in chicks were increased when chlortetracycline was fed, perhaps indicating that the absorption of these substances from the intestine was improved by the antibiotic. Biely and March¹⁴¹ reported that chlortetracycline had a sparing effect on riboflavin, niacin, and folic acid in the growth of chicks. Some evidence for a sparing effect of chlortetracycline on the riboflavin requirements of chicks on purified diets was noted in our laboratory.⁴³ Oleson and co-workers¹⁴² found a sparing effect of chlortetracycline on the vitamin B₁₂ requirements of chicks while Stokstad and Jukes¹⁴³ found that chlortetracycline reduced the vitamin B₁₂ requirement of chicks in some experiments but not in others. The high mortality of young chicks associated with a deficiency of vitamin B₁₂ was reduced by feeding chlortetracycline. Walbel and co-workers¹⁴⁴ found that penicillin or chlortetracycline markedly increased the growth of chicks fed diets containing suboptimal amounts of thiamine, but, when the level of thiamine in the diet was adequate, the growth response was only slight, thus indicating a sparing effect on the thiamine requirement. The

authors suggested that synthesis of thiamine by the intestinal flora might have been improved by the antibiotic. In another publication Walbel and co-workers¹⁴⁶ observed that the addition of penicillin to the diet of laying hens resulted in increased deposition of biotin and folic acid in the egg yolk.

Linkswiler et al.¹⁴⁵ noted that rats on inadequate levels of pyridoxine showed a somewhat greater growth response to chlortetracycline than rats on adequate levels of the vitamin, thus indicating a sparing effect of the antibiotic on the requirement of vitamin B₆. These investigations were extended considerably by Lish and Baumann¹⁴⁷ who compared the growth-promoting effects of various antibiotics for rats on diets deficient in B vitamins. The experimental data were not conclusive because the growth-promoting effects of the antibiotics were in most cases measured only at suboptimal levels of the vitamins; however, in the case of pantothenic acid there was some evidence of a sparing action. Chlortetracycline was found by McDaniel and Daft¹⁴⁸ to accentuate the signs of vitamin B₆ deficiency in rats while penicillin had a beneficial effect on growth and survival and in reducing the severity of dermatitis. This is in contrast with the findings for chlortetracycline reported by Linkswiler et al.¹⁴⁵

Coates and co-workers¹⁴⁹ studied the vitamin requirements of chicks on purified diets deficient in the various members of the vitamin B complex as influenced by the addition of penicillin to the diets. No sparing effect was found for thiamine, riboflavin, pyridoxine, or pantothenic acid but the requirements for biotin and folic acid were reduced. In the case of niacin, the requirement appeared to be increased. Nelson and Scott¹⁵⁰ found no effect of chlor tetracycline or penicillin on the niacin requirement of chicks on a purified diet. They noted that the antibiotic growth effect disappeared when the diet was severely deficient in niacin, which was in contrast with our findings,⁴³ although we noted that the antibiotic growth effect disappeared or was greatly diminished when chicks were severely deficient in riboflavin, pantothenic acid, or folic acid. The food intake is extremely low under highly deficient conditions so that very little growth takes place. Sarett¹⁵¹ found that the oral administration of 4 to 6 Gm. of streptomycin daily was followed by a marked decrease in the urinary excretion of biotin in human subjects. The effects on the excretion of other B vitamins were variable and not marked. The quantitative urinary excretion of certain products of tryptophane metabolism was not changed by the administration of streptomycin. In studies with rats, Halevy and co-workers¹⁵² found that streptomycin reduced the urinary excretion of folic acid and biotin. The folic acid content of the liver and the amount of biotin in the contents of the cecum were reduced. Varying effects were produced by the other antibiotics: chlortetracycline had a slight sparing effect on niacin, folic

acid, and biotin, oxytetracycline, on niacin and folic acid, but no effects on the requirement for these vitamins were found for penicillin. In an investigation by Sauberlich¹³³ on the requirement of weanling rats for certain B vitamins, sparing effects on the requirement for thiamine and pantothenic acid were obtained with chlortetracycline and penicillin and the requirement for pyridoxine was spared by penicillin as judged by the fact that marked growth responses were obtained to the antibiotics on the deficient diets while their inclusion in the completely vitamin supplemented diet had no effect upon growth of the animals. In contrast, sulfamerazine depressed growth on both the supplemented and deficient purified diets. Schendel and Johnson¹³⁴ studied the effects of penicillin and chlortetracycline on rats receiving suboptimum levels of thiamine and pantothenic acid administered both orally and subcutaneously. Marked increases in growth were produced by the antibiotics and the growth responses were absent when the diets were adequately supplemented with thiamine and pantothenic acid. The results were not affected by the mode of administration of the vitamins. The amount of pantothenic acid in the intestinal contents of rats on the deficient diet was increased when chlortetracycline was given. Along similar lines, Chow and co-workers¹³⁵ found that the oral administration of any of several antibiotics increased the apparent formation of vitamin B₁₂ by the intestinal bacteria as indicated by a rise in the butanol-soluble radioactive fraction following the administration of cobalt⁶⁰.

The effects of antibiotics on the requirement of animals for vitamin B₁₂ have been studied with varying results. Several reports have shown definite improvements produced by feeding antibiotics on the hatchability of hens eggs or on growth of rats, chicks, or pigs when the animals were fed B₁₂ deficient diets while in other investigations no changes were produced.

Johansson and co-workers^{87, 136} found that growth responses were produced in rats by adding chlortetracycline and vitamin B₁₂ singly or in combination to a diet deficient in vitamin B₁₂. The responses appeared to be independent of each other and no increase in the apparent vitamin B₁₂ content of the kidney and liver tissue was produced by feeding chlortetracycline, but increases were found, not unexpectedly when 25 µg/Kg. of diet of B₁₂ were added. Of interest was the observation that chlortetracycline produced increases in the vitamin B₁₂ activity as determined with *E. coli* 113-3 in the supernatant liquid obtained by suspending the contents of the colon and cecum in water and centrifuging off the solids. However *E. coli* 113-3, responds to cobalt-containing B₁₂-like substances that will not alleviate vitamin B₁₂ deficiency in animals.

Cravioto-Munoz et al¹³⁷ found that chlortetracycline produced a marked growth response in rats on a B₁₂-deficient, purified diet to an extent that led

these authors to state that chlortetracycline could replace vitamin B₁₂ in the diet of the rat. Autoclaved chlortetracycline was ineffective. No unsupplemented group was included.

The effect of antibiotics on the vitamin B₁₂ requirements of pigs has varied in different experiments. For example Catron and co-workers⁵⁸ found that chlortetracycline and vitamin B₁₂ both increased the growth rate of pigs on a corn-soybean diet and that the growth response to the vitamin was the same in the presence or absence of the antibiotic, while in another investigation¹⁰⁹ vitamin B₁₂ produced a 14 per cent growth increase in the absence of chlor tetracycline and no increase in its presence. Sheffy and co-workers,¹¹⁰ in a study with baby pigs on a semisynthetic milk, recorded the following daily gains: no supplement, 0.26 lb; vitamin B₁₂, 0.38 lb; chlortetracycline, 0.40 lb; B₁₂ plus chlortetracycline, 0.47 lb. These results indicated a sparing effect of chlortetracycline on the vitamin B₁₂ requirement. On the other hand, baby pigs that were highly deficient in vitamin B₁₂ did not show a growth response to chlortetracycline in a study by Wahlstrom and Johnson.¹⁰⁸ Monson et al.⁶⁰ made an interesting study with chicks that received diets containing insufficient amounts of folic acid, which were supplemented with antibiotics. They found that coliform organisms, which produced increased amounts of cellular folic acid *in vitro*, appeared in the duodenum and ileum following the administration of antibiotics. Simultaneously the folic acid content of the liver of the chicks was increased. This may be a further indication that antibiotic-resistant intestinal bacteria are in part responsible for the beneficial nutritional effects of dietary antibiotics.

It is evident from various investigations already described, which may be taken as representative of the large volume of work in this field, that dietary antibiotics in many cases improve the nutrition of animals with respect to the B vitamins, but in other cases the antibiotics are without any effect on the B vitamin requirement. A few exceptions have been noted, especially the case of streptomycin which appears to reduce the synthesis of biotin in the intestine. Antibiotics improve the production of B vitamins by the intestinal bacteria or the absorption of B vitamins into the blood stream.

Some effects of dietary antibiotics in modifying the requirements for animals for B complex vitamins were expected in view of the well known relationship of the intestinal and rumen microflora to synthesis of these vitamins. There was less basis for expecting an effect on the utilization of minerals and fat soluble vitamins. However experimental work soon uncovered such relationships.

Vitamin A and Carotene Burgess and co-workers²⁹ found that dietary

penicillin improved the utilization of vitamin A and carotene in chicks. Significant increases were found in the vitamin A content per Gm. of liver and in the blood level of carotenoids when penicillin was fed and the chicks were compared with control birds at 28 days of age. Along similar lines, Coates and co-workers²⁰ noted that the liver storage of vitamin A was improved by feeding penicillin to chicks and that the conversion of β -carotene to vitamin A in the intestinal wall was increased. The same effects were produced by placing the chicks in isolation units, which prevented contact with "infections." Further investigations in this field were made by Almqvist and Maurer²¹ who employed a diet containing only vegetable sources of vitamin A activity in contrast to the diet used by Burgess and co-workers, which contained both carotenoids and fish oil sources of vitamin A activity. The findings are shown in table XI, liver vitamin A was increased slightly by the lowest level of antibiotic fed and

Table XI

Vitamin A and Carotene in Chick Livers as Related to Feeding of antibiotics²⁰

| Supplement per Kg. of diet | Liver vitamin A (I U/Gm.) | | Liver carotene (μ g./Gm.) | | Average body wt. Gm. | |
|-------------------------------|------------------------------|--------|-----------------------------------|--------|-------------------------|--------|
| | 8 wk. | 10 wk. | 8 wk. | 10 wk. | 8 wk. | 10 wk. |
| None | 45 | 83 | 2.6 | 5.6 | 980 | 1363 |
| 4.5 mg. antibiotic mixture† | 55 | 94 | 3.1 | 7.6 | 989 | 1389 |
| 55 mg. chlortetracycline | — | 115 | — | 8.8 | 1070 | 1387 |
| 110 mg. chlortetracycline | 113 | 115 | 4.8 | 7.2 | 1076 | 1446 |

*Practical diet containing only plant sources of vitamin A activity approximately 11 I U/Gm.

†Chlortetracycline plus penicillin.

higher levels led to further elevations in both liver vitamin A and carotene. It was concluded that maintenance of a healthier intestinal condition in the presence of an antibiotic was responsible for improved conversion or absorption of carotene or both. It is of interest to compare the effect of antibiotics in preventing liver necrosis,⁴² an effect that is also produced by vitamin E, with their effect on vitamin A metabolism. Vitamin E has been reported in some

investigations to improve the utilization of vitamin A, possibly by protecting it against oxidation.

No sparing effect of chlortetracycline on vitamin A was found in experiments with rats by Hartsook et al.¹²¹ and chlortetracycline appeared to enhance the vitamin A deficiency syndrome when added to a diet free from the vitamin, as judged by earlier appearance of the signs of the deficiency and a shorter time of survival.

In contrast Murray and Campbell¹²² found that the addition of chlortetracycline to a vitamin A free diet for rats increased their response to graded doses of vitamin A acetate. The mean increase of many tests was 24 per cent. The effect was also observed when chlortetracycline supplementation was begun 24 hours after the last dose of vitamin A, which made it appear that the improved utilization was not due to an increase in absorption of vitamin A from the intestine. The time required for depletion of the liver stores of vitamin A was lengthened by dietary chlortetracycline.

The storage of vitamin A in the livers of pigs fed various supplements was measured by Barber and co-workers.¹²³ They found that on the basal diet the livers contained 12.8 units/Gm. of vitamin A, with chlortetracycline added, 19.3 and 19.6 units/Gm. with penicillin, 13.1 units/Gm. and with a mixture of penicillin and streptomycin residue 13.2 units/Gm. Chlortetracycline but not penicillin increased the liver storage of vitamin A.

Penicillin was found to produce increases in the vitamin A content of the liver of chicks on a normal diet and to increase the levels of carotenoids, ascorbic acid, and phospholipid phosphorus in their blood plasma.¹²⁴

High¹²⁵ found that chlortetracycline, 50 mg./Kg. of diet, increased the deposition of vitamin A from carotene in the liver and kidneys. Rats receiving 80 μ g. carotene as the sole source of vitamin A deposited 50 μ g. of vitamin A in the liver and kidneys while chlortetracycline-fed rats deposited 63 μ g. The storage of preformed vitamin A was not affected by the antibiotic. Penicillin had no effect on storage of vitamin A from carotene-fed rats. These results are similar to those reported by Barber and co-workers with pigs¹²³ in that chlortetracycline but not penicillin increased liver vitamin A storage on a diet containing carotene. The results with chicks differ for both Coates and Burgess reported increases in the liver vitamin A of chicks following the addition of penicillin to a diet containing carotene.

Vitamin D Calcium and Phosphorus Investigations by Ross and Yacowitz¹²⁶ indicated that penicillin appeared to decrease the vitamin D requirement for normal bone calcification. An increase in bone ash of chicks at three weeks was observed to be produced by adding penicillin to a diet containing

penicillin improved the utilization of vitamin A and carotene in chicks. Significant increases were found in the vitamin A content per Gm. of liver and in the blood level of carotenoids when penicillin was fed and the chicks were compared with control birds at 28 days of age. Along similar lines, Coates and co-workers²⁰ noted that the liver storage of vitamin A was improved by feeding penicillin to chicks and that the conversion of β -carotene to vitamin A in the intestinal wall was increased. The same effects were produced by placing the chicks in isolation units, which prevented contact with infections.* Further investigations in this field were made by Almquist and Maurer²¹ who employed a diet containing only vegetable sources of vitamin A activity in contrast to the diet used by Burgess and co-workers, which contained both carotenoids and fish oil sources of vitamin A activity. The findings are shown in table XI. Liver vitamin A was increased slightly by the lowest level of antibiotic fed and

Table XI

Vitamin A and Carotene in Chick Livers as Related to Feeding of antibiotics²²

| Supplement per Kg. of diet | Liver vitamin A (I. U./Gm.) | | Liver carotene (μ g./Gm.) | | Average body wt., Gm. | |
|-------------------------------|--------------------------------|--------|-----------------------------------|--------|--------------------------|--------|
| | 8 wk. | 10 wk. | 8 wk. | 10 wk. | 8 wk. | 10 wk. |
| None | 45 | 83 | 2.6 | 5.6 | 980 | 1363 |
| 4.5 mg. antibiotic mixture† | 55 | 94 | 3.1 | 7.6 | 989 | 1389 |
| 55 mg. chlortetracycline | — | 115 | — | 8.8 | 1070 | 1387 |
| 110 mg. chlortetracycline | 113 | 115 | 4.8 | 7.2 | 1076 | 1446 |

*Practical diet containing only plant sources of vitamin A activity approximately 11 I. U./Gm.

†Chlortetracycline plus penicillin.

higher levels led to further elevations in both liver vitamin A and carotene. It was concluded that maintenance of a healthier intestinal condition in the presence of an antibiotic was responsible for improved conversion or absorption of carotene or both. It is of interest to compare the effect of antibiotics in preventing liver necrosis,²³ an effect that is also produced by vitamin E, with their effect on vitamin A metabolism. Vitamin E has been reported in some

investigations to improve the utilization of vitamin A, possibly by protecting it against oxidation

No sparing effect of chlortetracycline on vitamin A was found in experiments with rats by Hartsook et al¹⁸¹ and chlortetracycline appeared to enhance the vitamin A deficiency syndrome when added to a diet free from the vitamin, as judged by earlier appearance of the signs of the deficiency and a shorter time of survival.

In contrast Murray and Campbell¹⁸² found that the addition of chlortetracycline to a vitamin A free diet for rats increased their response to graded doses of vitamin A acetate. The mean increase of many tests was 24 per cent. The effect was also observed when chlortetracycline supplementation was begun 24 hours after the last dose of vitamin A, which made it appear that the improved utilization was not due to an increase in absorption of vitamin A from the intestine. The time required for depletion of the liver stores of vitamin A was lengthened by dietary chlortetracycline.

The storage of vitamin A in the livers of pigs fed various supplements was measured by Barber and co-workers¹⁸³. They found that on the basal diet the livers contained 12.8 units/Gm. of vitamin A with chlortetracycline added, 19.3 and 19.6 units/Gm. with penicillin, 13.1 units/Gm., and with a mixture of penicillin and streptomycin residue, 13.2 units/Gm. Chlortetracycline but not penicillin increased the liver storage of vitamin A.

Penicillin was found to produce increases in the vitamin A content of the liver of chicks on a normal diet and to increase the levels of carotenoids, ascorbic acid, and phospholipid phosphorus in their blood plasma.¹⁸⁴

High¹⁸⁵ found that chlortetracycline 50 mg/Kg. of diet, increased the deposition of vitamin A from carotene in the liver and kidneys. Rats receiving 80 µg. carotene as the sole source of vitamin A deposited 50 µg. of vitamin A in the liver and kidneys while chlortetracycline-fed rats deposited 63 µg. The storage of preformed vitamin A was not affected by the antibiotic. Penicillin had no effect on storage of vitamin A from carotene-fed rats. These results are similar to those reported by Barber and co-workers with pigs¹⁸³ in that chlortetracycline but not penicillin increased liver vitamin A storage on a diet containing carotene. The results with chicks differ for both Coates and Burgess reported increases in the liver vitamin A of chicks following the addition of penicillin to a diet containing carotene.

Vitamin D Calcium and Phosphorus Investigations by Ross and Yacowitz¹⁸⁶ indicated that penicillin appeared to decrease the vitamin D requirement for normal bone calcification. An increase in bone ash of chicks at three weeks was observed to be produced by adding penicillin to a diet containing

suboptimal levels of vitamin D. These results may be compared with earlier observations by Migicovsky and co-workers¹⁶⁷ who found that penicillin increased the uptake of calcium¹⁶⁸ when fed to chicks on a low calcium diet. Gabuten and Shaffner¹⁶⁹ found that dietary penicillin increased the breaking strength of eggs and the blood calcium level in hens and Bogdonoff and Shaffner¹⁶⁰ reported that chlortetracycline, oxytetracycline, penicillin, and bacitracin increased the plasma calcium level of male chicks when added to a diet containing 2.6 per cent calcium. Penicillin was also fed with lower levels of calcium and was found to increase the plasma calcium level.

Lindblad and co-workers¹⁷⁰ found that the growth response of chicks and turkey poults to chlortetracycline was greater on higher than on lower levels of calcium and phosphorus.

Manganese Metabolism. Manganese is needed for growth and the prevention of a bone deformity termed "perosis" in chicks and poults. An effect of chlortetracycline on the utilization of manganese was described by Pepper and co-workers.¹⁷¹ They found that the amount of manganese requirement to prevent perosis in chicks was lowered by adding chlortetracycline at a level of 10 mg/Kg of diet.

Effect on Protein Requirement

A large number of experiments have been carried out to investigate the effect of dietary antibiotics on the level of protein required by poultry and swine for maximum growth. The economic importance of this question is obvious, for the supply of proteins of high biologic value may limit the numbers of poultry and pigs that can be produced on a nationwide basis. In some experiments it was found possible to produce maximum growth with slightly lower levels of protein in the presence of dietary antibiotics than in their absence. Other investigations have indicated that no difference in the protein requirement was produced by feeding antibiotics.

The protein requirement for maximum growth of chicks was found by Machlin and co-workers¹⁷² to be 19 per cent with a diet containing chlortetracycline and 21 per cent if the antibiotic was omitted. Weakley and co-workers¹⁷³ concluded that bacitracin, chlortetracycline, and penicillin all had a sparing effect on the protein requirement of chicks. Chlortetracycline and penicillin were found to improve the growth of chickens at four different levels of protein, and there was evidence that the u^{174} of both protein and energy was improved by the antibiotics, but $\left\{ \begin{array}{l} \text{of both protein and} \\ \text{indices} \end{array} \right.$ the

protein level for maximum growth was lowered.¹⁷⁴ Scott and co-workers¹⁷⁵ concluded that in their experiments the protein requirement of chicks was not changed by adding 15 mg. of chlortetracycline per Kg of diet. Growth and nitrogen retention were measured in chicks on various levels of protein with and without penicillin by Saxena and co-workers.¹⁷⁶ The increase in weight that accompanied penicillin feeding was the same at all levels small but not significant increases in nitrogen retention were found in the birds receiving the penicillin supplement at the lowest level of protein (15 per cent). Turkeys showed an improved response to penicillin as the level of fishmeal in the diet was increased,¹⁷⁷ and the greatest feed efficiency was obtained with penicillin and the highest level of fishmeal. Slinger and co-workers¹⁷⁸ deduced that the requirement for lysine was increased, simultaneously with the growth rate, by adding antibiotics to a diet containing 20 per cent protein. The lysine deficiency which manifests itself in young turkeys by the appearance of white feathers was not seen when the antibiotic-supplemented diet contained 28 per cent protein. Biely et al¹⁷⁹ found that chlortetracycline and penicillin did not affect the protein requirement of chicks or their requirement for lysine or tryptophane but did reduce the mortality on the tryptophane-deficient diet. Matterson and co-workers¹⁸⁰ found that chicks fed a ration containing chlor tetracycline required at least 21 per cent of protein for best results as measured by growth and feed efficiency. There was no indication that the antibiotic reduced the protein requirement. White-Stevens and Zisbel¹⁸¹ approached the problem from the standpoint of the levels of both protein and antibiotic. Their results with chicks showed that 10 parts per million of chlortetracycline was more effective as judged by growth on the low than on the high protein diet, while higher levels of chlortetracycline were most effective at the higher protein levels, the optimum being 22 per cent protein with 100 parts per million of chlortetracycline.

Studies with pigs have led to results very similar to those reported for chicks. There is some evidence for a slight lowering of the protein requirements for maximum growth, but in many experiments the antibiotic growth effect has been observed at varying protein levels without evidence of interaction. It has also been noted that feeding an antibiotic may improve nitrogen retention.

It is customary in feeding pigs to reduce the protein content of their diets stepwise during the growing period and this practice complicates an evaluation of the effect of antibiotic supplementation on protein requirement. Catron and co-workers¹⁸² investigated the effects of chlortetracycline on the growth of pigs receiving various levels of protein from weaning to 200 lb. The results are summarized in table XII. Each protein level was reduced by a step of 3

Table XII

Relation of Chlortetracycline 10 mg/lb. of Diet on Average Daily Gain and Food Utilization in Growing Pigs*

| Protein levels | Chlortetracycline in diet | Average daily wt gain, lb. | | | Initial-200 | |
|----------------|------------------------------|----------------------------|--------|---------|---------------|-------------------------|
| | | Initial† | 75-150 | 150-200 | Av daily gain | Feed per 100 lb of gain |
| 20 to 17 to 14 | — | 1.21 | 1.53 | 1.81 | 1.51 | 390 |
| | + | 1.31 | 1.79 | 1.94 | 1.68 | 354 |
| 18 to 15 to 12 | — | 1.29 | 1.58 | 1.81 | 1.56 | 383 |
| | + | 1.28 | 1.73 | 2.03 | 1.66 | 360 |
| 16 to 13 to 10 | — | 1.26 | 1.67 | 2.02 | 1.63 | 372 |
| | + | 1.34 | 1.83 | 1.80 | 1.67 | 355 |
| 14 to 11 to 8 | — | 1.23 | 1.64 | 1.54 | 1.50 | 374 |
| | + | 1.26 | 1.84 | 1.88 | 1.66 | 359 |

*The protein levels of the diets were lowered when the pigs reached a weight of 75 lb and again when the weights reached 150 lb.¹⁴²

†Sixteen pigs per group. Initial weight 29.9 to 37.5 lb.

per cent when the pigs reached 75 lb and again at 150 lb. In the absence of chlortetracycline, the rate of gain for the entire feeding period varied significantly among the several groups of protein levels, and the results indicated that in the presence of the antibiotic the protein requirement was lowered so that the pigs receiving successive levels of 14, 11, and 8 per cent of protein in the presence of the antibiotic gained as well as those receiving higher levels of protein in the absence of the antibiotic. The experiments were carried out with Duroc pigs on a diet consisting predominantly of corn and soybean meal. Jensen and associates¹⁴³ fed protein levels of 10, 12, 14, 16, 18, and 20 per cent, with and without 5 mg of chlortetracycline or oxytetracycline per lb. of diet, to pigs weighing 31 to 36 lb. until the pigs reached a weight of 200 lb. The

Table XIII

Increased Growth Rate in Rats Fed Chlortetracycline with Soybean Diets

| Supplement/100 Gm. diet | Average daily gain (Gm.) | | | |
|-----------------------------------|--------------------------|---------|---------------------|---------|
| | Raw soybean meal | | Heated soybean meal | |
| | Males | Females | Males | Females |
| None | 0.42 | 0.78 | 2.36 | 1.85 |
| Chlortetracycline 20 mg | 1.50 | 1.29 | 2.60 | 2.33 |
| DL-methionine 0.5 Gm. | 2.33 | 1.97 | 3.36 | 2.58 |
| Chlortetracycline plus methionine | 3.02 | 2.01 | 4.02 | 2.40 |

average daily gains of the pigs appeared to reach a maximum at between 16 and 18 per cent protein without antibiotics and at 14 per cent protein with antibiotics, but the effect was not statistically significant.

Effects on Nitrogen Metabolism

Streptomycin was found to have no effect on the utilization of food protein in rats by Black and Bratzler¹⁸⁴ An all-vegetable diet supplemented with vitamin B₁₂ was used. In another study¹⁸⁵ a supplement of antibiotics was used at a lower level, 10 per cent, of dietary protein, and again no effect was found on the nitrogen retention of paired fed rats due to the supplement. The utilization of energy was improved by the antibiotics.

Carroll and co-workers^{186, 187} investigated the effects of chlortetracycline and methionine on the utilization of nitrogen in diets containing either raw or heated soybeans when fed to rats. The amount of nitrogen absorbed from the diet during passage through the small intestine was increased by chlortetracycline but not by methionine. The antibiotic also improved the absorption of lysine, leucine, methionine and cystine. A marked increase in growth rate accompanied the feeding of chlortetracycline as shown in table XIII.¹⁸⁷ The lower availability of methionine in raw as compared with heated soybeans is well established. The authors concluded that a sparing action of chlortetracycline for dietary methionine was suggested by these results, either due to an

CHAPTER IV Antibiotics in Animal Production

The theoretical and experimental examinations of the mode of action of antibiotics give way to practical considerations when the effect of these substances in the diet is measured in terms of animal production, and it is in this field that antibiotics have made one of their greatest contributions. The steady increase of the human population makes it incumbent on chemical technology to furnish means of increasing the supply of food. The use of antibiotics has increased the supply of meat by lowering the death losses of young farm animals and by improving the efficiency of conversion of feed into meat. Laster¹⁸⁶ stated that the feeding of antibiotics to farm animals has brought about a new era in livestock production. The following appraisal of the effects of antibiotics on swine production is summarized from an article by Catron.¹⁸⁷ The practical effective feeding level of suitable antibiotics for swine is approximately 5 mg./lb. of feed and therapeutic levels commonly used in treating disease are 30 to 100 times as high. Antibiotics are effective in either dry lot or pasture feeding. Their use increases growth rate by 10 to 20 per cent from weaning to 200 lb. under average feed lot conditions with an increase in feed consumption of 5 to 10 per cent and a saving of about 5 per cent of feed required per unit of gain in growing and fattening pigs. Antibiotics produce the maximum growth response when fed to young pigs and the increase in gains in pigs weighing from 100 to 200 lb. is less. When antibiotic feeding is discontinued after pigs reach 75 to 125 lb. cessation of accelerated growth rate occurs and the growth rate tends to follow that of the control animals; however, the pigs previously fed antibiotics maintain their early growth advantage. The number of runts is reduced and the uniformity of pigs receiving antibiotics is several times greater than that of unsupplemented controls. The response to antibiotic feeding appears to be in proportion to the disease level; the practice controls a high percentage of nonspecific enteritis (scours) in swine but com-

Table XV

Effect of Chlortetracycline In the Presence of Chronic Respiratory Disease in Chicks

| Pleurapneumonia-like inoculum applied | Av. wt., Gm., at 4 wk. when chlortetracycline added, Gm./ton | | |
|---------------------------------------|--|-----|-----|
| | 0 | 50 | 100 |
| None (control) | 327 | 349 | 338 |
| Inoculum applied | 283 | 324 | 348 |

The reduction was evident in the weights at 4 weeks except on the higher level of chlortetracycline (table XV)

In an experiment under commercial conditions where chronic respiratory disease was endemic in which 5000 birds were used in each to two groups, the results at 10 weeks are shown in table XVI.

Various treatments as shown below were used in a flock of chickens in which signs of chronic respiratory disease appeared at the age of 6 weeks while the birds were on a diet containing 4 Gm./ton of procaine penicillin. The flock was divided into five groups each with about 4000 birds (table XVII).

The results are of great interest in that the efficiency of feed conversion was much lower in group 1 (unsupplemented) than in group 4 which grew at the

Table XVI

Effect of Feeding Antibiotics on Chicks with Chronic Respiratory Disease

| Supplement/ton of diet | Per cent mortality | Av. wt., Kg. | Gm. feed/Gm. gain |
|--|--------------------|--------------|-------------------|
| 4 Gm. procaine penicillin | 10.68 | 1.30 | 3.11 |
| 4 Gm. procaine penicillin plus 100 Gm. chlortetracycline | 2.82 | 1.40 | 2.81 |

There were 5000 chicks used in each group.

Table XVII

Results with Various Levels of Chlortetracycline on Chicks with Chronic Respiratory Disease

| Gm. of chlortetracycline added/ ton of diet | | Av final wt., Kg | Mortality % | Gm feed/ Gm gain |
|--|-------------|---------------------|----------------|---------------------|
| 6 to 8 wk. | 8 to 12 wk. | | | |
| None | None | 1.49 | 10.9 | 4.75 |
| 400 | 100 | 1.63 | 6.1 | 2.90 |
| 200 | 200 | 1.57 | 5.1 | 3.57 |
| 100 | 100 | 1.47 | 5.7 | 3.69 |
| 70 mg/liter in drinking water | None | 1.56 | 7.8 | 3.86 |

same rate, indicating that the effect of the antibiotic was to improve the utilization of nutrients, rather than to increase the appetite. In other experiments, the food intake per unit of body weight per day was about 20 per cent lower in groups receiving high levels of chlortetracycline than in controls. The findings by White-Stevens and his collaborators indicate the effect of a subacute disease in depressing the rate of growth and the utilization of nutrients by chicks and the action of chlortetracycline in reversing these nutritional sequelae of the disease.

Antibiotics in Swine Production

Pigs under farm conditions have been shown to be outstanding among domestic animals in growing more rapidly and efficiently when small amounts of antibiotics, especially chlortetracycline, are added to the diet. A comparison of various antibiotics, summarizing work published in 1950 to 1953 was made by Braude and co-workers²²⁵ and table XVIII, which has been published elsewhere,⁴¹ was condensed from their review. It may be seen from table XVIII that chlortetracycline, oxytetracycline, penicillin, and streptomycin have given the best results in promoting the growth of swine. Erythromycin was reported to produce no response.²²⁶

Table XVIII

Growth and Efficiency of Feed Utilization of Pigs Receiving Various Antibiotics²²⁵

| Antibiotic | Growth index (unsupplemented = 100) | | Feed required per unit of gain (unsupplemented = 100) | |
|-------------------|--|--------|---|-------|
| Chlortetracycline | 135.9 | (187)* | 90.2 | (146) |
| Penicillin | 110.6 | (53) | 94.3 | (44) |
| Streptomycin | 115.2 | (50) | 94.4 | (41) |
| Oxytetracycline | 123.7 | (23) | 93.9 | (17) |
| Butyracillin | 109.0 | (12) | 103.0 | (10) |
| Chloramphenicol | 105.5 | (6) | 98.2 | (6) |
| Neomycin | 93.3 | (4) | 87.6 | (3) |
| Polymyxin | 96 | (1) | 100 | (1) |
| Subtilin | 89 | (1) | 130 | (1) |

*Figures in parentheses indicate number of comparisons.

Dried whole cultures and fermentation residues containing chlortetracycline and vitamin B₁₂ were supplied by our laboratory early in 1949 to investigators who were engaged in studying the nutrition of swine. Marked growth responses, associated with the prevention of diarrhea, were soon observed and were communicated to us in the summer of 1949 by Catron, Cunha, and Krider. The effectiveness of the chlortetracycline fermentation residue was contrasted with the lack of activity of a vitamin B₁₂ concentrate from the streptomycin fermentation in a publication by Cunha and co-workers.⁴⁶ Further publications appeared in 1950.^{26, 27, 42, 43, 227} The effects of the supplement in abolishing scouring, and in producing its most marked effects where bloody diarrhea was rampant, promptly led to the establishment of the disease level concept as an interpretation of the antibiotic growth response in pigs. This concept has been reinforced by the observation that the response in pigs has progressively diminished in many laboratories where antibiotics are fed, while the growth of unsupplemented pigs has improved and the incidence of diarrhea has lessened.

Many of the diets used in the experiments with pigs were deficient in vitamin B₁₂ and consisted predominantly of corn and soybean meal so that the growth responses obtained to the crude chlortetracycline fermentation materials was due to both vitamin B₁₂ and the antibiotic. Experiments with crystalline antibiotics and vitamin B₁₂ soon established that both supplements were needed for optimum growth of pigs.²⁰

The use of antibiotics in diets for baby pigs, such as creep feeds and artificial milks, was found to be especially beneficial in preventing mortality and producing rapid early growth. Carpenter²² found that baby pigs receiving a creep feed containing chlortetracycline grew 43 per cent more rapidly than controls and Wahlstrom and co-workers²³ found that chlortetracycline increased the growth of baby pigs on a purified diet by 31 per cent. No effect was obtained with penicillin. Similar results were found by Noland and co-workers²⁴ and by Williams and co-workers.²⁵

Runtiness or unthriftiness in young pigs may often be successfully treated or prevented by adding antibiotics to the diet.^{22, 23} In one investigation²⁴ unthrifty pigs that were fed a diet containing 11 mg/Kg. of chlortetracycline gained three times as much weight in a five week period as did the unsupplemented controls.

The effect of the type of protein concentrate on modifying the growth response to antibiotics in pigs has been studied by several investigators. In the first published investigations with pigs, Burnside and co-workers²⁶ found that the crude fermentation containing chlortetracycline and vitamin B₁₂ produced a markedly greater response in pigs when peanut meal was used as compared with either soybean meal or fishmeal. Barber and co-workers¹⁴ found that the response to antibiotics was much less on a diet containing fishmeal than on an all vegetable diet and similar findings were described by Robinson et al.²⁴ These findings may be related to sparing effects of the antibiotics on the requirement for amino acids. Becker and co-workers²⁷ found in contrast to the work by Barber and Robinson that the growth response of pigs to chlortetracycline or to a mixture of antibiotics was greater with menhaden fishmeal than with soybean meal as the supplementary protein. In the absence of antibiotic, the pigs showed a growth response to tryptophane when added to the corn-fishmeal diet thus raising the question of a possible sparing effect of the antibiotic supplement. In other investigations, the response to antibiotics varied with different sources of protein but no consistent trend can be perceived and no doubt the results depend to some extent on variations in the nutritional quality of the various samples of meal that were used.^{23, 24}

The effect of implanting pellets of antibiotics on the growth of pigs was

Table XVIII

Growth and Efficiency of Feed Utilization of Pigs Receiving Various Antibiotics²²³

| Antibiotic | Growth index (unsupplemented = 100) | | Feed required per unit of gain (unsupplemented = 1) | |
|-------------------|--|-------|---|-------|
| Chlortetracycline | 135.9 | (187) | 90.2 | (146) |
| Penicillin | 110.6 | (53) | 94.3 | (44) |
| Streptomycin | 115.2 | (50) | 94.4 | (41) |
| Oxytetracycline | 123.7 | (23) | 93.9 | (17) |
| Bactracin | 109.0 | (12) | 103.0 | (10) |
| Chloramphenicol | 105.5 | (6) | 98.2 | (6) |
| Neomycin | 93.3 | (4) | 87.6 | (3) |
| Polymyxin | 96 | (1) | 100 | (1) |
| Subtilin | 89 | (1) | 130 | (1) |

*Figures in parentheses indicate number of comparisons.

Dried whole cultures and fermentation residues containing chlortetracycline and vitamin B₁₂ were supplied by our laboratory early in 1949 to investigators who were engaged in studying the nutrition of swine. Marked growth responses associated with the prevention of diarrhea, were soon observed and were communicated to us in the summer of 1949 by Catron, Cunha, and Kruse. The effectiveness of the chlortetracycline fermentation residue was contrasted with the lack of activity of a vitamin B₁₂ concentrate from the streptomycin fermentation in a publication by Cunha and co-workers.⁴⁶ Further publications appeared in 1950.^{24, 27, 48, 49, 227} The effects of the supplement in abolishing scouring, and in producing its most marked effects where bloody diarrhea was rampant, promptly led to the establishment of the disease level concept as an interpretation of the antibiotic growth response in pigs. This concept has been reinforced by the observation that the response in pigs has progressively diminished in many laboratories where antibiotics are fed, while the growth of unsupplemented pigs has improved and the incidence of diarrhea has lessened.

Many of the diets used in the experiments with pigs were deficient in vitamin B₁₂ and consisted predominantly of corn and soybean meal so that the growth responses obtained to the crude chlortetracycline fermentation materials was due to both vitamin B₁₂ and the antibiotic. Experiments with crystalline antibiotics and vitamin B₁₂ soon established that both supplements were needed for optimum growth of pigs.⁸⁰

The use of antibiotics in diets for baby pigs such as creep feeds and artificial milks, was found to be especially beneficial in preventing mortality and producing rapid early growth. Carpenter²²⁸ found that baby pigs receiving a creep feed containing chlortetracycline grew 43 per cent more rapidly than controls and Wahlstrom and co-workers²²⁹ found that chlortetracycline increased the growth of baby pigs on a purified diet by 31 per cent. No effect was obtained with penicillin. Similar results were found by Noland and co-workers⁹⁴ and by Williams and co-workers.²³⁰

Runtiness or unthriftiness in young pigs may often be successfully treated or prevented by adding antibiotics to the diet.²³¹⁻²³⁴ In one investigation²³⁴ unthrifty pigs that were fed a diet containing 11 mg/Kg of chlortetracycline gained three times as much weight in a five week period as did the unsupplemented controls.

The effect of the type of protein concentrate on modifying the growth response to antibiotics in pigs has been studied by several investigators. In the first published investigations with pigs, Burnside and co-workers²³⁵ found that the crude fermentation containing chlortetracycline and vitamin B₁₂ produced a markedly greater response in pigs when peanut meal was used as compared with either soybean meal or fishmeal. Barber and co-workers¹⁶² found that the response to antibiotics was much less on a diet containing fishmeal than on an all-vegetable diet and similar findings were described by Robinson et al.²³⁶ These findings may be related to sparing effects of the antibiotics on the requirement for amino acids. Becker and co-workers²³⁷ found in contrast to the work by Barber and Robinson that the growth response of pigs to chlortetracycline or to a mixture of antibiotics was greater with menhaden fishmeal than with soybean meal as the supplementary protein. In the absence of antibiotic, the pigs showed a growth response to tryptophane when added to the corn fishmeal diet thus raising the question of a possible sparing effect of the antibiotic supplement. In other investigations, the response to antibiotics varied with different sources of protein but no consistent trend can be perceived and no doubt the results depend to some extent on variations in the nutritional quality of the various samples of meal that were used.²³⁸⁻²⁴³

The effect of implanting pellets of antibiotics on the growth of pigs was

lambs appeared in 1951 when Jordan and Bell²⁷⁶ found that chlortetracycline promoted the growth of lambs and decreased enterotoxemia.

Many investigators of antibiotics in the feeding of ruminants, conducted for the most part with chlortetracycline, were published in the years between 1951 and 1955. Reviews have been written by Knodt²⁷⁸ and Lassiter.¹⁹⁴ Chlortetracycline has been reported to increase the growth rate, feed consumption, and efficiency of feed utilization of young calves and lambs and to reduce the incidence of certain diseases including scours, pneumonia, and liver abscesses in calves, and enterotoxemia in lambs. With older ruminants, digestive upsets have sometimes been encountered, but it is often possible to forestall these by regulating the composition of the ration and the dosage of the antibiotic. The most marked effects have been found with unthrifty animals. Chlortetracycline and other antibiotics have been found to depress the digestion of cellulose *in vitro* by mixed cultures of rumen microorganisms, but such effects do not seem to be evident in the intact animal.

Effects on Dairy Calves Marked increases in the rate of growth of dairy calves during the first 5 or 6 months of life have been observed by various investigators when chlortetracycline was added to the ration. The most marked effects were observed in calves which were housed in contaminated quarters where the incidence of scours was high²⁷⁷ but responses were also obtained by Bartley and co-workers²⁷⁸ when the unsupplemented animals were gaining at a normal rate. Two levels of chlortetracycline were compared, 15 mg. and 45 mg./day and the higher level was found more effective in promoting growth and in reducing the incidence of various types of infections. In further studies,²⁷⁹ levels of 45 and 90 mg. of chlortetracycline were compared and were found to be equally effective in reducing the incidence of infections. Much larger doses, 200 to 800 mg. of chlortetracycline per 100 lb. of calf weight, were fed to calves without producing anorexia or weight loss or any other visible detrimental effects and one calf not previously supplemented, was given 2.5 Gm. of chlortetracycline daily between 16 and 20 weeks of age without deleterious effects. Many investigations with chlortetracycline in the diets of dairy calves have been carried out by Rusoff and co-workers.^{280, 281}

^{280, 281, 284} The antibiotic was found to increase the rate of gain of both Jersey and Holstein calves by 20 per cent over that of the control animals through 16 weeks of age. Comparisons were made between oral supplementation and intramuscular injection and the injected animals were found to grow more rapidly than those receiving the antibiotic orally. These observations led the authors to conclude that the mode of action of chlortetracycline in producing the antibiotic growth response was not due to an effect in the rumen because

the injected antibiotic was shown to be present in the bile and not in the rumen contents. At autopsy the animals receiving chlortetracycline showed larger skeletal size and more meat than did the controls. This was thought to indicate that the antibiotic was producing an effect on the pituitary gland. The authors were unable to detect any chlortetracycline in the pituitary, spleen, thymus, and muscular tissues so that it is difficult to implicate the pituitary gland on the basis of these studies. No effects on the yield and composition of the milk were produced by feeding supplements of 130 mg of chlortetracycline daily to dairy cows²⁸⁴ nor were physiologic effects produced when chlortetracycline was fed to bulls at a rate of 300 mg daily.²⁸⁵ A decrease in thickness of duodenal and jejunal sections of the intestinal wall and an increase in the ileal section were found in calves receiving chlortetracycline.²⁸⁶

Richardson and co-workers²⁸⁷ found no growth-promoting effect of chlortetracycline when given by intramuscular injection to dairy calves. The gains in 4 weeks were 8.2 lb for unsupplemented calves, 12.0 lb for calves receiving two levels of oral chlortetracycline, 9.0 lb for calves implanted subcutaneously with chlortetracycline, and 8.2 lb for calves injected intramuscularly with chlortetracycline.

It was noted by Roy and co-workers²⁸⁸ that the beneficial effect of colostrum on calves appeared to be related to the prevention of scouring. Accordingly the effects of chlortetracycline and colostrum were compared in experiments in which 40 calves were used, and it was found that the calves receiving chlortetracycline without colostrum were protected against scours in a manner that resembled the effect of colostrum when fed without chlortetracycline. The authors made the interesting suggestion that colostrum might be important for its antibacterial effect. The feeding of chlortetracycline to colostrum-deprived calves increased live-weight gain significantly and reduced the incidence of scouring and the occurrence of abnormally high rectal temperatures. Colostrum-fed calves behaved similarly to the chlortetracycline-fed calves when compared with unsupplemented controls.

No effect on milk production and feed consumption was found when chlortetracycline was fed to mature lactating dairy cows which received 130 mg daily for 30 days,²⁸⁹ 300 to 500 mg daily²⁹⁰ or 700 mg daily for 10 days.²⁹⁰ The antibiotic was not detected in the milk in the latter investigation.

Antibiotics in Beef Cattle Feeding

The addition of chlortetracycline or penicillin at high levels ranging up to 600 mg daily to the rations of beef cattle was found in earlier investigations

to be followed by various effects including a loss of appetite, a depression of the apparent digestibility of crude fiber and a decrease in nitrogen retention.^{111, 291, 292} Such apparent digestive disturbances were noted to disappear when the feeding period was prolonged.¹⁰⁴ In later investigations it was found that chlortetracycline was tolerated better when fed at lower levels and when the diet was high in roughage, thus Tovrea²⁹³ found the following average daily gains in an experiment lasting 90 days with 20 beef cattle per group in which the initial weight of the animals was about 660 lbs. when chlortetracycline was given in doses of 0, 72, 90 and 108 mg per animal per day the average daily gain was 2.76, 3.12, 3.11, 3.00 lb., respectively. In a second experiment, an unsupplemented group gained 2.61 lb. daily per head, a group receiving 80 mg of oxytetracycline daily 2.65 lb. and a group receiving 80 mg. of chlortetracycline daily 2.81 lb.

Matrushima and co-workers²⁹⁴ found that calves which were fed 15 mg chlortetracycline daily grew more rapidly than control calves and were free from scouring which occurred frequently in the controls. The calves were slaughtered at 7.5 to 10 months and 90 per cent of the supplemented calves had normal livers as compared with 10 per cent of the controls, 70 per cent of which had abscessed livers, presumably from infections which entered through the portal circulation. Perry and co-workers²⁹⁵ observed a marked decrease in the incidence of scours when chlortetracycline was fed at 10 mg /100 lb of body weight daily to beef calves. The daily gain was 1.78 lb. in the supplemented group as compared with 1.59 lb. for the controls during the 80 day experimental period. In the same investigation, the effect of chlortetracycline on older beef animals was examined. The addition of chlortetracycline to a concentrate-type of ration for yearling heifers and steers, starting at 9.1 mg /lb of grain and increasing stepwise to 27.2 mg resulted in daily gains of 1.64 lb. as compared to 1.52 for controls. In other experiments, marked improvements in rate of growth and cellulose digestion were observed when chlortetracycline was added to high roughage rations (table XX) containing supplement A, which consists of 65.1 per cent soybean meal, 14 per cent molasses, 14 per cent alfalfa meal, 5.2 per cent bonemeal, 1.7 per cent cobalt, and a concentrate supplying 2270 units of vitamin A and 200 units of vitamin D per pound of supplement.

Duitsman and Kessler^{296, 297} compared bacitracin and chlortetracycline for 122 day periods in two experiments, each of which involved 10 animals per group fed 122 days on a ration of silage, milo and cottonseed meal with the result that in the first experiment those animals fed no supplement had an average daily gain of 1.54 those with 45 mg daily of bacitracin added, 1.6

those with 45 mg daily of chlortetracycline added, 1.72. In the second experiment the average daily gains were 1.45 for those fed no antibiotic supplement, 1.84 for those fed 45 mg daily of bacitracin, and 1.52 when 45 mg daily of chlortetracycline was added.

The experimental work with beef cattle in which favorable results were reported was carried out principally with chlortetracycline. A level of about 10 mg/100 lb of body weight was suitable and the best responses were on rations high in roughage. Setbacks may occur when antibiotics are given at high levels, especially to animals that are receiving high-concentrate diets.

Antibiotics in Lamb Feeds

The oral administration of fairly high doses of antibiotics to lambs was found to produce digestive disturbances by Colby and co-workers,²⁹⁹ who noted that lambs receiving daily 100 mg of chlortetracycline or penicillin lost weight and developed diarrhea while Turner and Hodgetts²⁹⁸ noted that a single dose of 750 mg. of chlortetracycline to Merino sheep was followed by loss of appetite and weight. Lower supplementary levels are better tolerated. Jordan²⁹⁶ and Jordan and Bell²⁹⁷ found that lambs on fattening rations tolerated low dosage with chlortetracycline and there was some reduction of enterotoxemia. In further experiments,³⁰⁰ chlortetracycline was fed to lambs at levels of 4.3 to 10.8 mg. daily and slight improvements in growth were seen in three of four trials. In another report, lambs receiving chlortetracycline in a pelleted ration required 10 per cent less feed per unit of gain than controls.³⁰¹ Hatfield and Garrigus³⁰²⁻³⁰³ studied the effects of chlortetracycline when 10 mg /lb was added to a concentrated feed which was offered free choice with alfalfa hay. The supplemented lambs grew about 10 per cent faster than the controls and produced slightly better carcasses.

These investigators³⁰³ summarized the results of three experiments involving 190 lambs. Rations supplemented with chlortetracycline at levels of 5.5 to 7.6 mg /lb gave small but consistently higher average daily gain and produced daily gains which were $.055 \pm .014$ lb higher than those of the controls. The feed efficiencies and carcass grades of the supplemented lambs were higher than those of the controls. The average performance of lambs receiving diets containing oxytetracycline and penicillin was approximately the same or slightly better than that of controls but the differences were not significant. Chlortetracycline did not eliminate enterotoxemia but the authors concluded that "the improvement in gain and efficiency was sufficient to show that chlortetracycline has a practical place in lamb rations."

Bridges et al³⁰⁶ fed chlortetracycline to lambs at levels of 1.1 to 15 mg./lb. of diet. Increases in the rate of gain resulting from feeding the antibiotic were small and were not statistically significant. Slight improvements in efficiency of feed utilization and in carcass grade and dressing percentage were noted in the antibiotic-fed lambs. Effects of the composition of the ration on the response to chlortetracycline were recorded by Elliott and Ellsworth³⁰⁷ who found an effect of the antibiotic on growth with 80 per cent of roughage and 20 per cent of grain but not with higher levels of grain while no relation between roughage level and response was found by Botkins and Paules³⁰⁸ who reported that the average daily gain was 0.287 lb. for the controls and 0.304 for the lambs receiving 10 mg. of chlortetracycline per lb. of ration. In an earlier investigation, Botkins and Paules³⁰⁹ found no response to chlortetra-

Table XX

Effect of Chlortetracycline on Growth of Beef Cattle Receiving various rations²⁹³

| Days of experiment | No. animals per group | Additions to supplement A | Chlortetracycline addition mg./day | Average | |
|--------------------|-----------------------|-------------------------------------|------------------------------------|------------------|-----------------|
| | | | | Initial wt., lb. | Daily gain, lb. |
| 56 | 5 | Ground corn cobs | 0 | 682 | 1.69 |
| 56 | 5 | Ground corn cobs | 75 | 680 | 2.18 |
| 56 | 5 | Ground corn cobs | | 678 | 1.83 |
| 161 | 9 | Ground corn cobs | 0 | 713 | 1.26 |
| 161 | 9 | Ground corn cobs | 75 | 713 | 1.53 |
| 168 | 9 | Ground cobs plus ground yellow corn | 0 | 711 | 2.18 |
| 168 | 9 | Ground cobs plus ground yellow corn | 75 | 715 | 2.16 |
| 49 | 12 | Green chopped corn | 0 | 1021 | 2.28 |
| 49 | 12 | Green chopped corn | 75 | 1032 | 2.71 |

First week 25 mg., second week 50 mg., then 75 mg. daily

cycline 30 mg per lamb per day during a 117 day feeding period. Reductions in scouring and enterotoxemia were found to be associated with feeding chlortetracycline by Kunkel and co-workers^{210, 211} as follows in 118 without antibiotic 8.5 per cent died of enterotoxemia and 9.3 per cent died of all causes in 232 lambs fed 5 to 15 mg of chlortetracycline per lb of diet, 17 died of all causes with no deaths from enterotoxemia. Enterotoxemia in lambs is commonly accompanied by clostridial infection of the digestive tract. No effects of antibiotics, chlortetracycline, oxytetracycline and penicillin, were obtained in an experiment with lambs by Kinsman and Riddell.²¹² Bohman and co-workers²¹³ fed graded levels of chlortetracycline, 5 to 20 mg./lb of grain mix, to 140 lambs from 2 to 4 weeks of age to market weight, with or without added B vitamin. No detrimental or beneficial effects were noted on rate of gain dressing percentage or carcass grade.

The experimental work with lambs indicates that a variable but predominantly favorable effect is obtained when lambs are fed a low level of chlortetracycline, about 10 mg /lb of diet. Many of the observed effects appear to be related to the suppression of harmful bacteria. High dosage with antibiotics may cause digestive disturbances. Good results have been found when antibiotics were fed to cull or scrub lambs.

The Effect of Dietary Antibiotics on Reproduction in Animals

A number of investigators have studied the results of adding antibiotics to the diet on reproduction in chickens, pigs, and, in a few cases, laboratory animals. Most of the measurable effects have been beneficial. In a number of cases, no effects were found. The interpretation of such results may be complicated if the maternal diet is inadequate with respect to certain vitamins which are "spared" by an antibiotic.

Antibiotics do not readily pass through the placental barrier into the fetus as judged from studies with pigs nor do antibiotics tend to become concentrated in the egg yolk. Carpenter and Larson²¹⁴ fed 1 Gm. of chlortetracycline three times daily to sows, starting two days before the expected farrowing date. The baby pigs were bled within the first few hours after birth and before they had nursed, but no antibiotic was found in their blood serum, although it was present at levels of 0.03 to 1.3 μ g /ml. in the serum of all 5 of the sows.

Various investigators have studied the birth weight and survival rate of young pigs as affected by adding antibiotics, in most cases chlortetracycline, to the maternal diet. In some of the earlier experiments, crude supplements

Table XXI
Effect of Chlortetracycline on Reproduction in Pigs

| Investigators | Level of chlortet- racycline Gm /ton | No of sows and gilts | | Av no. pigs/ litter | Av birth wt., lb. | Pigs born dead, % |
|---|---|-------------------------|----------|---------------------------|----------------------|----------------------|
| | | Started | Farrowed | | | |
| Carpenter and | 0 (a) | 16 | 15 | 7.0 | 2.95 | 7.1 |
| Lanson ³¹⁴ | 20 (a) | 23 | 17 | 8.3 | 2.98 | 4.8 |
| | 0 (b) | 14 | 14 | 10.6 | 2.9 | 18.9 |
| | 20 (b) | 14 | 14 | 8.6 | 3.1 | 5.8 |
| Catron ¹⁹⁷ | 0 | 20 | 18 | 8.9 | 2.89 | 9 |
| | 40 | 20 | 19 | 9.7 | 2.84 | 16 |
| | 0 | 10 | 7 | 6.9 | 2.90 | 10 |
| | 20 | 11 | 10 | 6.1 | 2.88 | 5 |
| | 0 | 25 | 20 | 10.0 | 2.90 | 10 |
| | 40 | 25 | 19 | 10.3 | 2.86 | 8 |
| | 0 | 24 | 20 | 10.3 | 2.9 | 4 |
| | 10 | 24 | 22 | 10.0 | 3.0 | 11 |
| | 0 | 30 | 25 | 8.8 | 3.1 | 5 |
| | 10 | 30 | 26 | 10.6 | 3.0 | 6 |
| Davey Green and Stevenson ³¹⁸ | 10 | 16 | 4 | 10.0 | 2.68 | 5.0 |
| | 50 | | 5 | 7.2 | 3.06 | 28 |
| | 100 | | 3 | 6.7 | 3.08 | 4.5 |
| | 10 | 8 | 3 | 7.0 | 2.66 | 38 |
| | 50 | 8 | 4 | 9.5 | 2.13 | 8.4 |
| | 100 | 8 | 7 | 8.7 | 2.51 | 3.4 |
| Vesta ³¹⁵ | 0 | 13 | 11 | 8.8 | 2.45 | 6.2 |
| | 20 | 26 | 25 | 9.4 | 2.58 | 4.7 |
| Beeson ³¹⁷ | 0 | 13 | 12 | 9.8 | 2.51 | 0.9 |
| | 10 | 13 | 12 | 9.1 | 2.36 | 2.8 |
| | 20 | 13 | 10 | 9.4 | 2.60 | 7.4 |
| | 30 | 13 | 12 | 9.4 | 2.82 | 0.9 |

were used which contained vitamin B₁₂ as well as an antibiotic so that the effects could not be definitely attributed to the latter. Carpenter²¹⁰ fed a supplement containing chlortetracycline and vitamin B₁₂ for 49 to 90 days prior to parturition without observing any effect on the size of the litter, the average birth weight, or the number of pigs born dead. Further studies²¹⁴ in which the supplement was fed through two gestation periods led to similar results, although slight differences were observed in favor of the supplement (table XXI). Vestal and co-workers²¹³ used a similar supplement and found that the control pigs had an average birth weight of 2.45 lb. as compared with 2.58 lb. for the pigs from the sows that received the supplement at a level supplying 9 Gm. of chlortetracycline and 9 mg./ton of vitamin B₁₂. No effects of such a supplement on the birth weights were found in studies by Dyer.²¹⁶

Beeson and co-workers²¹⁷ fed various levels of crystalline chlortetracycline to sows during gestation (table XXI). They found that heavier and stronger pigs were produced at birth when the level of chlortetracycline in the maternal diet was 20 or 30 Gm./ton than when the level was 0 or 10 Gm. The antibiotic had no effect on the number of pigs per litter. The authors summarized their work as follows: "Gilts fed Aureomycin at levels of 20 to 30 Gm. per ton of feed farrowed heavier and stronger pigs at birth than the gilts receiving no antibiotics or a level of 10 Gm. of Aureomycin. Aureomycin had no significant effect on the number of pigs farrowed or the weaning weight of the pigs. Pigs from the sows receiving the high level of antibiotic (30 Gm. per ton) during pregnancy were stronger and more uniform in appearance at weaning. The percentage of the pigs weaned was not significantly influenced by antibiotic feeding but was slightly higher for the high Aureomycin group."

The effect of chlortetracycline on reproduction in swine through successive reproductive cycles and generations was exhaustively studied by Davey and co-workers.²¹⁸ Diets containing relatively high levels of the antibiotic, 50 and 100 mg./lb. of feed, were included together with a lower level in order to accelerate or intensify the effects of continuous antibiotic feeding. The high levels produced no adverse effects in number of pigs born per litter, birth weights, or weaning weights as compared with a low level. The results by Davey and co-workers in table XXI represent experiments in consecutive years. The results in the second year were obtained with the daughters of the females used in the first year so that the effects of chlortetracycline were measured through two generations. The lower average weights of the second year pigs was stated to be probably due to the use of younger sows in the second year.

The effects of antibiotics in the diet of the sow on the growth of baby pigs

are usually complicated by the fact that the young animals consume part of the mother's food, and the experimental results are difficult to evaluate.

Maddock and associates²¹⁹ found that chlortetracycline was not detectable in the blood and milk of sows receiving the antibiotic at levels up to 20 mg./lb. of diet. However when single oral doses of chlortetracycline up to 4 Gm. were given by capsule, a maximum level of 0.74 μ g./ml. was found in the blood serum and 2.21 μ g./ml. in the milk whey.

The effect of dietary antibiotics on the composition of hen's egg has an important bearing on any effects on hatchability or embryonic development. When any of several antibiotics were injected into eggs, the injections were well tolerated by the developing embryo (table III) at levels that far exceeded those that have been found in eggs following the administration of high levels of antibiotics to laying hens. Broquist and Kohler⁴¹ found that no antibiotic potency was detectable in eggs laid by hens receiving a diet containing 20 or 200 Gm./ton of chlortetracycline, and only small amounts, ranging from 0.15 to 3.1 μ g. per egg, were found when the antibiotic level was 2000 Gm./ton of diet. Wong and co-workers²²⁰ were unable to detect the presence of oxytetracycline and penicillin in eggs even when the hens were fed the antibiotics at levels of 1000 Gm./ton of diet. Clearly any effects of the antibiotics on hatchability when fed at the usual levels are probably exerted indirectly rather than through their presence in the egg. This view is supported by publications which indicate that the content of certain vitamins in the egg may be increased following the feeding of antibiotics to hens. Walbel and co-workers¹¹⁵ found increases in the biotin and folic acid content of egg yolks when penicillin was fed to laying hens at levels of either 5 or 200 mg./Kg. of diet. These authors suggested that the increases were due to changes in the intestinal flora brought about by the antibiotic. The basal diet was not deficient in biotin or folic acid.

The effect of certain antibiotics in increasing the hatchability of eggs laid by hens on diets deficient in vitamin B₁₂ presents some unsolved problems. Several investigators have reported that this improvement in hatchability was not accompanied by an increase in the vitamin B₁₂ content of the egg or that the increase in vitamin B₁₂ was insufficient to account for the improvement in hatchability.^{22, 221, 222} Sizemore and co-workers²²² reported in addition that the feeding of a supplement containing chlortetracycline to pullets during the growing period made them resistant to vitamin B₁₂ deficiency of the breeder diet, measured by a depression in hatchability which could be corrected by vitamin B₁₂, and that this resistance persisted after the cessation of antibiotic feeding for a period of several months. This was further emphasized by their

later observations¹² in which the resistance continued into the second year after withdrawal of the supplement. Laying house mortality egg production, and fertility were not influenced by the growing or the breeder diets. However the presence of a vitamin B₁₂-antibiotic supplement in the growing diet resulted in increased hatchability during the two year period in all four comparisons (table XXII) and the addition of chlortetracycline to the breeder diet also resulted in increased hatchability. Embryonic mortality showed similar trends. The effect of the antibiotic during the laying period was beneficial and the addition of a supplement containing vitamin B₁₂ and chlortetracycline to the rearing diet was followed by improvements in the subsequent reproductive performance. The authors suggested that the prolonged feeding of the antibiotic may influence the intestinal flora or may make available a nutrient to

Table XXII

Relation of Diets during Rearing Period and Laying Period to Mortality of Hens, Egg Production and Hatchability of Fertile Eggs in Rhode Island Red Hens

| Series | Diet during rearing period | Mortality during 2 yr | Egg production during 2 yr | Hatchability | |
|--------|----------------------------|-----------------------|----------------------------|--------------|-------------|
| | | | | 2d yr | 1st & 2d yr |
| A | 1 | 55.0 | 47.0 | 46.4 | 53.9 |
| A | 1 + AFS† | 55.0 | 47.9 | 33.2 | 59.4 |
| A | 2 | 40.0 | 42.6 | 38.3 | 43.8 |
| A | 2 + AFS | 60.0 | 45.6 | 73.6 | 79.2 |
| | Average | 52.5 | 45.8 | 47.9 | 59.1 |
| B | 1 | 30.8 | 42.1 | 61.3 | 66.8 |
| B | 1 + AFS | 43.9 | 41.9 | 74.3 | 74.4 |
| B | 2 | 62.5 | 43.8 | 67.5 | 63.8 |
| B | 2 + AFS | 47.5 | 45.0 | 71.6 | 73.3 |
| | Average | 46.2 | 43.2 | 68.6 | 69.6 |

Series A received no antibiotic during the two year laying period and series B received a supplement of crystalline chlortetracycline.

† AFS = Antibiotic vitamin B₁₂ supplement in rearing diet.

increase hatchability. It was not possible to explain the of the apparent vitamin B₁₂ content of the egg yolks for when the birds received chlortetracycline even though the eggs was higher. The presence of an unidentified "hatchability" of hens receiving an antibiotic as a supplement to a diet B₁₂ was also noted by Elam and co-workers.²² Further, known vitamin like factor in the eggs of hens which had supplements was provided by Slinger and co-workers^{22,23} as Hershberger^{22,24} Slinger and co-workers found that the progeny had received penicillin were 3 per cent heavier at 10 weeks from hens without the antibiotic supplement. The chicks on a commercial diet that was supplemented with penicillin and Bentley and Hershberger fed an all vegetable diet to laying hens the effect of vitamin B₁₂ and antibiotics in the diet of hens on the progeny. The addition of chlortetracycline or bacitracin to hens increased the growth of the progeny on vitamin B₁₂ deficient though the egg yolks, livers of the hens and the livers of the chicks showed no difference in vitamin B₁₂ content as determined with Leichmann's. Furthermore, feeding procaine penicillin to the hens increase the growth of their progeny when both the hens and received diets containing additional amounts of vitamin B₁₂. The chicks again raise the possibility that a growth promoting factor vitamin B₁₂ may be stored in the egg as a result of feeding antibiotics. They fed egg yolks to rats which received a basal corn soybean diet or a diet vitamin B₁₂ and iodinated casein. Egg yolks from hens receiving chlortetracycline, bacitracin, or penicillin produced more rapid growth in rats from unsupplemented hens.

Egg Production in Poultry

The nutritional requirements of chickens and turkeys for egg production in many respects less critical than the requirements for hatchability. In my terms this may be expressed in the statement that on certain deficiencies hens will continue to lay eggs at a fairly good rate though the hatchability of the eggs falls to extremely low levels, or in extreme cases, the eggs will hatch at all. The essential nature of certain vitamins, like acid, and vitamin B₁₂ or hatchability under sanitary conditions are hens are usually have been found to have no effect on the hatchability of eggs in the absence of riboflavin. High rate of egg production is not possible without adequate riboflavin.

duction. Increases have been observed to accompany the feeding of antibiotics when the rate is depressed by certain unfavorable environmental conditions. Several investigators have noted improvements in hatchability when antibiotics were fed, and in some cases a marked increase was noted when the diet was deficient in vitamin B₁.

No increase in hatchability and no consistent increase in egg production were noted when antibiotics were fed by Sunde et al,²²⁷ although the highest egg production was in a group receiving a supplement containing chlortetracycline. Berg and co-workers²²⁸ found no effects on egg production or hatchability when oxytetracycline or chlortetracycline were fed at levels ranging from 4 to 15 mg/Kg of diet. The reproductive performance of all the groups was excellent. Walbel and co-workers¹⁴⁴ stated without presenting data that the addition of penicillin to a diet of natural foodstuffs resulted in no measurable effect on egg production, egg weight, or hatchability over a 10 month period. The biotin and folic acid contents of the egg yolk were greater in the supplemented groups. Lillie and Bird²²⁹ found no significant effects of a supplement containing chlortetracycline on egg production or hatchability and Brown and co-workers²²⁹ reported no effects with penicillin in a similar experiment with laying hens. Other investigators who have reported that antibiotics were without effect on egg production include Petersen and Lampman²³⁰ who used penicillin streptomycin and oxytetracycline, Halbrook and co-workers²³¹ who used chlortetracycline with both chickens and turkeys, Sherwood and Milby²³² who used chlortetracycline, oxytetracycline, and penicillin, Carpenter and Duckworth²³³ who used chlortetracycline, Mariakulandia and co-workers²³⁴ who used oxytetracycline and Johnson²³⁴ who used penicillin.

Elam and co-workers²³⁵ reported increases in egg production when antibiotics were added to the diet of pullets. 10 birds were used in each group. A decrease in egg production was noted in hot weather by Heywang²³⁶ who found that the addition of 50 or 100 Gm of chlortetracycline per ton of diet helped to prevent the decrease. Balloun²³⁷ noted that the addition of chlortetracycline at a high level (100 Gm./ton of diet) improved egg production in pullets which had gone into a laying slump. Egg production increased from the January average of 29.4 per cent to 40.7 per cent in February in two groups where chlortetracycline was added to the diet at the rate of 100 Gm./ton. The improved rate of production was continued in March. No increase in production occurred in unsupplemented controls during the same periods.

Heywang²³⁸ found that the addition of chlortetracycline at levels of 50 or 100 Gm./ton to the diet of laying pullets resulted in greater egg production during periods of hot weather than was observed in unsupplemented controls.

This may indicate a diminished resistance to chlortetracycline-sensitive injurious microorganisms during hot weather (average mean temperature 88 F)

White-Stevens and Zeibel²³⁹ studied the effects of chlortetracycline during various periods of the growing period on egg production by pullets. A significant and consistent month to-month difference in egg production was found between hens receiving 50 Gm. of chlortetracycline per ton of diet throughout the growing and laying period as compared with controls. Three groups of 65 hens were used for each treatment. The per cent production (eggs per 100 hens per day) for 12 month periods was as follows: there was 65.9 per cent egg production without antibiotic, 69.1 per cent when chlortetracycline was fed before egg-laying started, 67.3 per cent when chlortetracycline was fed only after egg-laying started, and 69.8 per cent when chlortetracycline was fed throughout the 12 month period. Carlson and Kohlmeier²⁴⁰ selected physically superior pullets and fed them a diet containing 50 to 100 Gm./ton of chlortetracycline, following which these birds produced more eggs than controls which had appeared to be physically superior. Carlson and co-workers²⁴¹ also noted that egg production in turkeys was improved slightly when chlortetracycline was added to a diet which contained supplementary vitamin B₁₂.

Antibiotics and Egg Production in Hens with Chronic Respiratory Disease

The appearance of signs of chronic respiratory disease in laying hens is characteristically accompanied by a drop in egg production. The effect of high levels of chlortetracycline in the diet in the control of chronic respiratory disease was studied by White-Stevens and co-workers.²⁴² In one experiment chronic respiratory disease appeared in a breeder flock containing 296 hens that had been in production for 10 months. The flock was divided into two equal groups and one group received an addition of chlortetracycline, 100 Gm./ton of feed. At the start of the experiment the hens in the supplemented group were laying 7.4 per cent less than the controls but during the final week of the 45 day experimental period the birds receiving the antibiotic produced 7.9 per cent more eggs per hen per day than the controls. A second experiment was carried out using 630 hens per group. Levels of 0, 50, 100 and 200 Gm. of chlortetracycline per ton of feed were used. The highest level was decreased to 100 Gm./ton after five months. The cumulative egg production of the first two groups was equal during a nine month period and egg production in groups 3 and 4 were respectively 8.3 and 3.3 per cent more than the control group. The hatchability for group 1 was 79.0 per cent, for group 2, 83.4 per

cent for group 3, 63.3 per cent and for group 4 83.1 per cent, so that the combined effects of increased egg production and hatchability were in favor of the supplemented groups. Carson and co-workers²⁴² found that egg production in pullets with CRD was improved by injecting a suspension of 50 mg of oxytetracycline in mineral oil. The mean egg production of 143 treated birds for 81 days of the postinjection period was 44.7 eggs as compared with 38.6 eggs for the control birds.

The effect of chlortetracycline on the egg production by turkeys in flocks where sinusitis and bluecomb were occasionally observed was improved by chlortetracycline. The average egg production in two unsupplemented groups totaling 593 birds was 45 and 55 per cent. The production of two corresponding groups receiving 115 Gm./ton of chlortetracycline was 57 and 58 per cent and for two groups with 230 Gm./ton of chlortetracycline, 59 and 61 per cent.²⁴²



CHAPTER V Physiologic and Other Effects Of Dietary Antibiotics

Effects of Antibiotics on the Endocrine System

Chlortetracycline and the Thyroid Gland. Ershoff has reported^{244, 245} that crude supplements derived from the fermentation of penicillin or chlortetracycline prolonged the survival of immature rats which received toxic doses of desiccated thyroid gland. He noted that crystalline chlortetracycline hydrochloride was ineffective thus perhaps indicating the existence of an unidentified nonantibiotic factor in the residues. However Meltes and Ogle²⁴⁶ found that penicillin and neomycin were effective in overcoming the growth-depressing effect of iodinated casein for rats. No goitrogenic effects were observed for the antibiotics. Calesnick and co-workers²⁴⁷ reported that either penicillin or chlortetracycline at the low level of 1 mg/Kg of diet produced a severalfold increase in the weight of thyroid glands of immature white rats with a three-to-fourfold decrease in I^{131} uptake by the gland after 42 days of feeding. They suggested that chlortetracycline and penicillin had a goitrogenic action similar to the thiouracils. Inability to repeat these observations was reported by Libby and Meltes²⁴⁸ and by Grant.²⁴⁹ In studies with rats, even though 20 to 50 times higher levels of the antibiotic were fed than were used by Calesnick, Grant noted in contrast that there was a small but definite increase in the uptake of I^{131} by the thyroid when chlortetracycline was fed to rats. It was also pointed out by Grant that rats receiving 2 per cent propylthiouracil in the diet as used by Calesnick and co-workers lived for less than two weeks in his experiments as opposed to the results reported by Calesnick et al who indicated that their rats lived through the six week period on the propylthiouracil supplement.

No effect on the iodine metabolism of thyroid slices in vitro was found in experiments with chlortetracycline by Taurog and Chaikoff.²⁵⁰ These investi-

gators incubated sheep thyroid tissue slices in Ringer solution containing with or without 10 μ g./ml. of chlortetracycline. After two hours of incubation at 37 C the tissues were homogenized and hydrolyzed with pancreatin and the hydrolysate was chromatographed for separation of the iodine-containing fractions. The chromatograph indicated that both preparations contained approximately the same amounts of radioactive diiodotyrosine, monoiodotyrosine, roxine, and free iodide so that chlortetracycline had no effect upon the uptake of I^{131} or on organic I^{131} formation by sheep thyroid tissue slices. Their findings indicate that any effect of dietary antibiotics on the thyroid is indirect, probably mediated through the intestinal bacteria.

Similarly Telkkä and Kuusisto²³¹ found no differences in the histological picture of the thyroid gland of antibiotic-supplemented rats as compared with the controls. Both chlortetracycline and oxytetracycline were used at levels of 10 mg./Kg. of diet. The relative weight of the gland was practically the same in all three groups. Maghrabi and Turner²³² noted a slight depression of thyroid size in young chicks that received 50 mg. of chlortetracycline per Kg. of feed. A small increase in the weights of the thyroid glands of chicks receiving chlortetracycline was noted at eight weeks of age by Mellen and Waller. The thyroid weight was 67 mg./Kg. of body weight for the unsupplemented birds, 83 mg. for the chlortetracycline-fed birds, and 76 mg. for the birds receiving bacitracin. No such differences were found by Grant²³³ as shown in table XXIII.

Both chlortetracycline and vitamin B₁₂ were reported by Meltes^{234, 235} to counteract the deleterious effects of cortisone on vitamin B₁₂-deficient rats. The effect of the antibiotic would seem to be related to improving the utilization of B₁₂.

Table XXIII

Average Body Weights, Thyroid Weights, and I^{131} Uptake of 10 Week Old Male Chickens 24 Hours after Intravenous Injection of I^{131}

| Chlortetracycline added per ton of diet | Body wt., Gm. | Thyroid wt. mg./Kg. body wt. | I^{131} uptake C.P.M. per mg thyroid |
|--|------------------|---------------------------------|--|
| None | 1259 | 90.1 | 158 |
| 100 Gm | 1359 | 89.4 | 156 |

*There were 10 birds per group.²³⁴

Table XXIV

Partial Protection of Rats Against Alloxan Diabetes by Chlortetracycline

| Treatment | Chlortetra- cyclin per Kg of diet Gm | Av gain in body wt in 28 days Gm. | Average daily | |
|-----------|--|---|------------------|----------------------|
| | | | Diuresis, ml. | Urinary sugar Gm. |
| None | 0 | 28 | | |
| None | 0.2 | 28 | | |
| Alloxan | 0 | 9.6 | 73 | 3.85 |
| Alloxan | 0.2 | 19.0 | 88 | 5.0 |

tion of vitamin B₁₂ or increasing its intestinal synthesis Wahlstrom and Johnson¹⁰⁰ found that cortisone aggravated vitamin B₁₂ deficiency in baby pigs but no alleviation was produced by chlortetracycline.

An effect of chlortetracycline on partially protecting rats against alloxan diabetes was reported¹⁰⁷ and is summarized in table XXIV

The experiment was carried out because it had been observed that patients with severe diabetes had shown improvement during treatment with chlor tetracycline.

The effects of chlortetracycline singly and in combination with diethylstil bestrol and L-thyroxine on the growth of pigs were studied by Barber and co-workers¹⁰⁸ who found evidence for an additive effect on growth and feed efficiency when the combination was fed.

Effects Following Various "Stress" Conditions Baez et al¹⁰⁹ found an 18 per cent greater survival after rotation in a drum in rats receiving oral chlortetra cycline than in untreated controls. These results prompted an examination of the liver slices of the experimental rats, and it was found that the liver preparations from rats receiving chlortetracycline orally or via portal vein in jection did not release vasoactive ferritin (VDM) into the incubation medium after 90 and 120 minutes of anoxia, in contrast to control liver slices. This suggested a mechanism by which chlortetracycline may have partially protected the drummed rats against shock. Fine¹⁰⁰ reported that penicillin and chlortetracycline were ineffective when administered parenterally in prevent

ing irreversible shock induced by hemorrhage in dogs, but these antibiotics together with neomycin produced a resistance to shock when given by mouth.

Chlortetracycline and streptomycin at levels of 50 mg/Kg. of diet were found to increase the tolerance to emetine in rats on a low protein diet.³⁴¹


Radiation Injury Following exposure of the total body to roentgen rays acute radiation sickness, accompanied by hemorrhage and infection, may develop. A number of investigations have shown that protective effects may sometimes result if antibiotics are administered to experimental animals prior to irradiation. Such effects may be important in the event of accidental exposure to radioactive materials. Penicillin, streptomycin, chlortetracycline, and oxytetracycline have all been used in these experiments.

Howland and co-workers³⁴² found that chlortetracycline alleviated diarrhea and prolonged the survival time for five to seven days beyond the control in rats following an LD₅₀ dose of roentgen ray radiation. Chloramphenicol, streptomycin, and penicillin were less effective. Good results were also obtained with chlortetracycline in dogs. Miller and associates³⁴³ found that the mortality of irradiated mice was lowered by streptomycin and to a lesser degree by chlortetracycline but that the protection was less if the mice developed certain infections. Gustafson and Koletsky³⁴⁴ reported that oxytetracycline probably protected against radiation injury by reducing bacteremia, in an experiment in which oxytetracycline was given 72 hours prior to radiation, the mean survival time was raised to 150 days from 106 days.

In further studies by Furth and his associates^{345, 347} the results with dogs and rats were indefinite. In two experiments, each involving 96 dogs, no protective effects were obtained from either chlortetracycline or whole blood transfusions. The authors nevertheless concluded³⁴⁷ that "the prophylactic administration of a broad spectrum antibiotic is still advisable despite the results of this study" in the event of an atomic catastrophe and the consequent exposure to infection.

Antibiotics and Liver Physiology

Antibiotics and Deficiencies of Choline and Methionine. Deficiencies of folic acid and vitamin B₁₂ are well known to lead to increase the disturbances arising from dietary deficiencies of choline and methionine in the diets of rats and chicks. Folic acid has a coenzymatic role via tetrahydrofolic acid (THF) and its derivatives in the synthesis of methionine from homocysteine and "single-carbon unit," which may be derived from formaldehyde, serine, or formate as follows ^{348, 349}



The role of vitamin B₁₂ is not clear but it seems to be connected with the formation or utilization of CH₂OH THFA, the hydroxymethylated coenzyme that is involved in one route for the formation of methionine from homocysteine. The second route for this synthesis is through the combination of betaine or dimethylthetin with homocysteine in the presence of transmethyrase. If the utilization or intestinal synthesis of folic acid and vitamin B₁₂ is improved by adding antibiotics to deficient diets, the effects of deficiencies of choline or methionine such as fatty livers, hemorrhagic kidneys, or necrotic livers, may be alleviated²⁷⁰ and anti-anemia effects of the antibiotic may also be shown.

Hepatic necrosis, a dietary liver disease of rats, has been extensively studied by Gyorgy and co-workers who showed that it was possible to produce the disease on a diet high in yeast and to prevent it by adding vitamin E, cystine or methionine. Chlortetracycline was also found to prevent hepatic necrosis when added to the diet of rats²⁷¹ but in contrast to the other preventive agents the effect of chlortetracycline was found to be only temporary. The mechanism for this effect of the antibiotic was thought to be an inhibition of toxin-producing bacteria in the lower intestine because the necrosis developed predominantly in the left lobe of the liver which derives its portal blood from the large intestine and stomach in contrast to the relatively normal right lobe which receives portal blood from the small intestine. Abell and Beveridge²⁷¹ confirmed the results with chlortetracycline and found that sulfaguanidine had some protective action. A number of antimicrobial agents were tested²⁷² and chlortetracycline was found to give the longest protection, followed by oxytetracycline and streptomycin, then by sulfaguanidine while chloramphenicol, polymyxin, and penicillin were ineffective. Vitamin B₁₂ had no effect. The temporary nature of the protection given by chlortetracycline led to the conclusion that its effect was not direct but was mediated through a transient change in the intestinal flora and that eventually the bacterial

types that were postulated as producing hepatotoxicosis reappeared in an antibiotic-tolerant form. In later studies penicillin was found to be effective.³⁷³ The intestinal flora of the experimental rats on penicillin and chlortetracycline supplements was studied and the type of changes produced by the two antibiotics differed, furthermore, resistant bacteria appeared in the first week while the antinecrogenic effect lasted for five or six months.³⁷⁴

Against the bacterial theory of hepatic necrogenesis were observations with germ-free rats³⁷⁵ which developed the disease when on a restricted food intake but not when on full feed. This makes it appear that some ingredient of the deficient diet could prevent necrosis and that the ingredient, perhaps methionine or cystine, was absorbed in sufficient amounts only when the animals were on full feed under the germ-free or antibiotic-supplemented conditions both of which have been reported to be accompanied by certain changes in the intestinal wall.³⁶ The addition of antibiotics produced a change in the pattern of the urinary excretion of ether-soluble acids as compared with unsupplemented or vitamin E-treated animals.³⁷⁶ Two unidentified acids with R values of 0.45 and 0.80 in butanol-acetic acid water mixtures appeared in the urines of the antibiotic-supplemented rats, these substances had no apparent relation to necrogenesis but their presence may be of much biochemical interest. The urinary excretion of volatile phenolic compounds by human subjects was found to decrease markedly following the administration of antibiotics by mouth. This was thought to be due to a depression of the bacterial formation of phenols from tyrosine in the intestinal tract.³⁷⁷

In other experiments by Gyorgy and his collaborators,³⁷⁸ rats were fed an entirely different diet, deficient in labile methyl and containing sucrose, 33 per cent lard, and 8 per cent casein, which produced cirrhotic changes and kidney injury. The changes were prevented by chlortetracycline, 25 mg daily.³⁷² These findings with rats may be compared with clinical investigations in which chlortetracycline was used in the treatment of hepatitis,^{378, 379} hepatic coma,³⁸⁰ and cirrhosis.^{381, 382} A patient with chronic posthepatic cirrhosis and continuous jaundice of four years duration showed striking improvements, including histopathologic changes, following five months intensive treatment with chlortetracycline.³⁸¹ Three patients with acute fulminating hepatitis recovered following treatment with chlortetracycline.³⁷⁹ However in a controlled study³⁷⁸ no difference in the course of acute viral hepatitis was found between chlortetracycline-treated and control patients when 37 cases were studied in each group. De la Huerga and co-workers³⁸³ reported that dietary chlortetracycline, oxytetracycline, and penicillin suppressed the urinary excretion of trimethylamine in dogs following the ingestion of a test dose of choline. This

effect was thought to be due to the lowering of trimethylamine formation by the intestinal bacteria. The effect disappeared after one or two weeks as if the bacteria had become adapted to the presence of the antibiotic. However Kaplan and co-workers³⁴⁴ were unable to repeat these observations. They found that chlortetracycline had a lipotropic effect on dogs with ligated pancreatic ducts when given in large doses. Similar changes occurred when choline was administered. The effects produced by chlortetracycline reached a maximum in three to four weeks and then receded and leveled off at an intermediate level, in one animal a second response was produced by adding folic acid and vitamin B₁₂ to the treatment with chlortetracycline. Either choline or chlortetracycline were found by Baxter and Campbell³⁴⁵ to prevent renal lesions in rats on diets high in fat and low in choline. Vitamin B₁₂ had some activity but oxytetracycline, chloramphenicol, and penicillin were ineffective. No change in the liver lipids of rats was produced by adding chlortetracycline to a low protein, low fat diet in studies by Cornatzer and Gallo³⁴⁶ and after four weeks there was a decrease in phospholipid synthesis in the group fed chlortetracycline. On the clinical side, 12 cases of chronic liver disease were treated with chlortetracycline, 7 of whom showed an improvement while in the remaining 5 cases there was little or no evidence of benefit.³⁴⁷

These effects of antibiotics may be contrasted with the findings of Seto and Lepper³⁴⁸ who found that an increase in the liver fat of young female rats was produced by a single intravenous injection of chlortetracycline or oxytetracycline, 100 or 150 mg /Kg of body weight as follows: the per cent liver fat, 18 to 24 hours after injection of the chlortetracycline was 15.5 ± 5.4 and the control was 7.1 ± 1.6 the per cent liver fat, 18 to 24 hours after injection of oxytetracycline, was 13.5 ± 6.2 , and the control was 6.0 ± 1.3 .

The liver fat values returned to normal within three days following withdrawal of the drugs. No change in the effect was noted when methionine or various B vitamins were given simultaneously. The experimental intravenous route of administration may have been the reason for the difference between these results and those previously described,^{372, 349} and it is interesting to note that Zimmerman and Humoller³⁵⁰ found that oral chlortetracycline did not effect the endogenous respiration and choline oxidase activity of rat liver tissues while chlortetracycline given parenterally or added in vitro depressed these systems. However Sborov and Sutherland³⁵⁰ found that treatment with chlortetracycline, 2 Gm. daily by mouth in a group of 16 patients with chronic viral hepatitis or hepatic cirrhosis, was followed by increased liver fat as indicated by biopsy. These patients received a diet which averaged 126 Gm. of fat daily so that the increases in liver fat were perhaps to be expected, moreover

toxic properties. Bacitracin may produce renal tubular necrosis, especially in mice. Neither streptomycin nor bacitracin are readily absorbed from the digestive tract when given orally.

Massive doses of chlortetracycline were given to pigs by Cameron⁴⁰⁹ who found no toxic effects when 250 mg./Kg. of body weight was fed daily for 2 days. The level of the antibiotic normally consumed by pigs receiving typical chlortetracycline-supplemented feeds would not exceed 1 mg./Kg. of body weight daily. The pigs were in excellent condition at the end of the experiment and the bone marrow was found to be yellow in color at autopsy but showed no histologic abnormality.

Effects on Growth of Human Subjects

An approach to the study of the effects of antibiotics on the growth of children presents certain difficulties as compared with experiments with animals, first, because every effort must be made with children to exclude the bacterial contaminations that contribute so much to the antibiotic growth effect in animals and, second, because of the slow rate of growth of human beings. The most marked effects of antibiotics on the growth of children may be expected where they are poorly developed due to subnormal diets or to subacute intestinal diseases.

Several encouraging reports have appeared regarding premature infants. Perrini⁴¹⁰ fed chlortetracycline, 25 mg./Kg. of body weight daily to 10 premature infants and found that they gained 8 per cent in weight during a 10 day period as compared with no weight increase for a group of 23 controls. Robinson⁴¹¹ also reported increased growth with premature infants receiving chlortetracycline who gained 29.5 Gm. daily as compared with controls who gained 18 Gm. daily. Five of the 15 controls died from intercurrent infections while all of the supplemented group lived. Lower mortality in a group of chlortetracycline-treated premature infants together with increased growth was reported by Snelling and Johnson⁴¹² there was one death in a group of 47 who received 50 mg. of chlortetracycline daily as compared with 8 deaths in 48 controls. In a study of 57 supplemented premature infants and 56 controls, Coodin⁴¹³ found that oxytetracycline, 25 mg. daily failed to alter the rate of weight gain. The mortality rates were not reported. No effects in increasing weight gains were found in 2 infants studied by Kaye et al.⁴¹⁴

Growth increasing effects for antibiotics on infants were reported by Déchéne⁴¹⁵ and Wetzel and Hopwood,⁴¹⁶ who found that oxytetracycline, streptomycin, and penicillin, 10 to 50 mg. daily by mouth, improved the

growth of school children Scrimshaw et al⁴¹⁷ studied Mayan children 7 to 12 years of age living in the Guatemalan highlands and subsisting on diets low in animal protein. In one experiment, 14 children who received 50 mg. of chlor tetracycline daily for 15 months gained an average of 0.48 cm. of height monthly as compared with 0.33 cm. for 43 control children. In two successive 12 month periods following withdrawal of the antibiotic, they had the same rate of height increase as control groups. In other experiments the increases in height are shown in table XXV. The average monthly weight gains varied widely irrespective of the type of treatment. It was concluded that the oral administration of chlortetracycline daily stimulated growth in stature in primary school children under the experimental conditions encountered but the response to penicillin was variable.

Carter⁴¹⁸ carried out a prolonged investigation of a study of the effects of administering 75 mg. of chlortetracycline twice daily to mentally defective children for periods up to three years at the Florida Farm Colony. The children were mentally deficient spastic cases and were almost entirely helpless. The average yearly gain in weight for the supplemented group was 6.5 lb. while the control group averaged 1.9 lb. in yearly weight gain. The supplemented group had a lower incidence of gastrointestinal disorders than the controls.

The effect of chlortetracycline on weight gains of undernourished children in Kenya was studied by Macdougall.⁴¹⁹ Thirteen children in the age range of 1 to 3 years received chlortetracycline, 50 mg. daily for one to seven weeks and

Table XXV

Increases in Height of Children with and without Antibiotics

| Experiment | Length of time | Monthly gains in height cm.* | | |
|------------|----------------|------------------------------|------------------|--------------------------|
| | | Controls | 50 mg penicillin | 50 mg. chlortetracycline |
| 1 | 15 months | 0.33 (43) | | 0.48 (14) |
| 2 | 18 months | 0.43 (23) | 0.41 (32) | |
| 3 | 5 months | 0.35 (38) | 0.36 (38) | 0.44 (35) |
| 4 | 5 months | 0.43 (35) | 0.51 (39) | 0.50 (37) |

The number of children per group is given in parentheses.

Public Health Aspects of Feeding Antibiotics to Animals

The prolonged feeding of comparatively high levels of the common antibiotics to animals has not raised problems in public health as regards the consumption of animal products. Broquist and Kohler⁴¹ found no detectable amounts of chlortetracycline in the liver and muscle of chicks receiving chlortetracycline, 200 mg./Kg. of diet. Barely detectable amounts were present in the blood serum. At unusually high dietary levels, traces of the antibiotic were found in the liver and muscle. The values ranged from less than 0.05 to 0.165 $\mu\text{g./Gm.}$ of muscle when the diet contained 2 Gm./Kg. of chlortetracycline up to 0.18 to 0.39 $\mu\text{g./Kg.}$ of liver with 8 Gm. of chlortetracycline per Kg. of diet. Similarly with pigs the level in the muscle tissue was 0.80 $\mu\text{g./Gm.}$ with 2 Gm. of chlortetracycline per Kg. of diet and undetectable in the muscle with 200 mg./Kg. of diet. In further studies, Kohler and co-workers⁴² found that chlortetracycline in poultry meat was readily destroyed by cooking and the destruction of chlortetracycline proceeded more rapidly than that of oxytetracycline. Chicken breast muscle containing 3 $\mu\text{g./Gm.}$ of chlortetracycline was found to be free from the antibiotic after boiling for 15 minutes or after roasting for 30 minutes at 230 C.

The prolonged administration of chlortetracycline to human subjects has not produced untoward effects. A group of 184 elderly patients were dosed orally with 500 mg. of chlortetracycline daily for prolonged periods in a study by McVay and Sprunt and an equal number in a control group received placebos.⁴³ In the groups with congestive heart failure, there were 15 deaths in the chlortetracycline group who were medicated for 20 months and a decrease in the incidence of respiratory infections in 30 patients as compared with 19 deaths in the control group and a decrease in respiratory infections in 4 control patients. Hematocrit values were higher in the chlortetracycline group. Along similar lines 189 diabetics were observed for an average of 19.5 months, again with a decrease in respiratory infections in the chlortetracycline group. Comprehensive liver function studies showed some evidence of improvement in this group. These observations emphasize the lack of toxicity of chlortetracycline when administered for prolonged periods at high dosage rates. No toxic reactions are imaginable from the traces of chlortetracycline which might be present in the meat of animals fed high levels of the antibiotic. Indeed, these traces of antibiotic were only of the order of 50 $\mu\text{g./lb.}$ or less of muscle tissue even when the chickens received 2000 Gm. of chlortetracycline per ton of diet⁴¹ and the antibiotic was destroyed by ordinary cooking procedures. It

is also worth noting that the antibiotic-tolerant strains of bacteria, which appear in the intestinal tracts of animals receiving dietary antibiotics appear not to be pathogenic: the report of Libby and Schaible¹²³ indicates a downward trend in the mortality of chickens during four years of feeding antibiotics. Markham and co-workers¹²⁴ found that the feeding of high levels of chlortetracycline did not inhibit the development of a normal immune response to vaccination for Newcastle disease and infectious bronchitis in chicks.

Conclusions

An apparent improvement in the nutritional status of animals may be often produced by adding small quantities of certain antibiotics to the diet. The utilization of food is often increased in growing animals by this addition as measured by a consequent decrease in the amount of food required to produce a unit of gain in body weight. The daily rate of gain in weight is increased and the percentage of unthrifty animals is reduced. These effects are of great economic and practical importance in agriculture. Many experimental investigations of the nutritional changes underlying these effects have been made and a sparing action of antibiotics on the requirement for certain nutrients has often been noted.

The mechanism by which the antibiotics produce their effects on nutrition appears to be secondary to their antibacterial action. Variations in the extent of these effects are inevitable in view of the inherent variations in the bacterial population. Changes presumably take place in the intestinal bacteria and these changes are responsible for the nutritional effects. Examinations of the types of bacteria in the intestinal contents of antibiotic-fed animals have led to variable and inconclusive findings as regards any correlation between feeding antibiotics and the establishment of specific bacterial types. When antibiotics are administered, the total numbers of bacteria in samples of the intestinal contents characteristically decrease for a few days and then increase above the original levels. Perhaps antibiotics suppress certain bacterial forms which are inhibitory not only to the growth of the host but also to the growth of some of the other intestinal bacteria.

The prolonged and widespread feeding of antibiotics to farm animals has not led to deleterious results or to the establishment of resistant strains of pathogenic bacteria. There is much evidence which indicates that one of the results is that the premises may become cleaned up so that the growth of the resident animals is improved even in the case of those animals on the same premises which are given unsupplemented diets.

21. MARTIN G. J.: Aminobenzoic acid and sulfonamides in rat nutrition, *Proc. Soc. Exper. Biol. & Med.* 51:56, 1942.
22. MOREHOUSE, N. F., AND MAYFIELD, O. J.: The effect of some aryl anionic acids on experimental coccidiosis infection in chickens, *J. Parasitol.* 33:20, 1946.
23. STOKESAD E. L. R., AND JONES, T. H.: Further observations on the animal protein factor *Proc. Soc. Exper. Biol. & Med.* 73:523, 1950.
24. WHITEHILL, A. R., OLSEN J. J. AND HUTCHINGS B. L.: Stimulatory effect of aureomycin on the growth of chicks, *Proc. Soc. Exper. Biol. & Med.* 74:11 1950.
25. GROSCHKE, A. C., AND EVANS, R. J.: Effect of antibiotics, synthetic vitamins, vitamin B₁₂, and an APF supplement on chick growth, *Poultry Sc.* 29:616, 1950.
26. LUECKE, R. W.; McMILLEN W. N., AND THORP F., JR. The effect of vitamin B₁₂, animal protein factor and streptomycin on the growth of young pigs, *Arch. Biochem.* 26:469 1950.
27. CARPENTER, L. E.: Effect of aureomycin on the growth of weaned pigs, *Arch. Biochem.* 27:469 1950.
28. BURGESS, R. C.; GLUCK, M. BRUNSON G. AND LAUGHLAND, D. H.: Effect of dietary penicillin on liver vitamin A and serum carotenoids in the chick, *Arch. Biochem.* 33:339 1951.
29. ALMQUIST H. J. AND MAURER, S. The effect of antibiotic, antioxidant, and fat on conversion of carotene to vitamin A in the chicken, *Arch. Biochem.* 55:297 1953.
30. COATES, M. E.; HARRISON G. F. KON, S. K. PORTER, J. W. G., AND TROMBSON, S. Y.: Antibiotics in chick nutrition and vitamin A metabolism, *Chem. & Indust.* 7:149, 1952.
31. HESTER, H. H., JR., LANDAGORA, F. T. AND RUBOFF L. L.: The distribution of aureomycin in the body of dairy calves showing a growth response when the antibiotic is administered orally or intramuscularly *J. Animal Sc.* 13:988, 1954.
32. STEEDING, R. B.; GRAEBLE, O. E., AND ROBINSON H. J. Studies on the absorption and excretion of streptomycin in animals, *Proc. Soc. Exper. Biol. & Med.* 60:68, 1945.
33. WELCH, H. Principles and Practice of Antibiotic Therapy New York, Medical Encyclopedia, Inc., 1954, p. 144.
34. MATTHEWSON, L. D., AND SIMON E. P.: A comparison of several antibiotics as growth stimulants in practical chick-starting rations, *Storrs Agric. Exper. Sta. Bull.* No. 275, p. 20, 1951.
35. BRANNON H. D.; HILL, D. C., AND MOTZOK, L.: Effect of subcutaneous implantation of bacitracin on the growth of chicks, *Poultry Sc.* 31:1090, 1952.
36. EAGLE, H. NEWMAN E. V., GRIFT R.; BURKHOLDER, T. M., AND GOODMAN, S. C.: Blood levels and renal clearance in rabbits and man of an antibiotic derived from *B. subtilis* (bacitracin) *J. Clin. Investigation* 26:919 1947.
37. ELAM, J. F. GEE, L. L., AND COUCH, J. R. The effect of feeding and injecting penicillin and aureomycin on the growth and intestinal microflora of the growing chick, *Poultry Sc.* 30:935 1951.
38. DIXON J. AND TRAYER, R. H. Observations on the growth-promoting action of antibiotics in the chick, *Poultry Sc.* 30:910, 1951.
39. COOK, D. L.; LAWLER, C. A., AND GREEN D. M.: Excretion and concentration of antibiotics in bile, *Federation Proc.* 11:833, 1952.
40. MONROE, W. F.; HANPER, A. E.; WENY, M. E.; ELVEHJED, C. A., RYONES, R. A., AND SANLEY, W. B.: Mechanism of the vitamin-sparing effect of antibiotics, *J. Nutrition* 52:628, 1952.

41. BROQUST H P AND KOELLER, A R.: Studies of the antibiotic potency in the meat of animals fed chlortetracycline. In: Antibiotics Annual 1953-1954 New York, Medical Encyclopedia, Inc., 1953, p. 409
42. SIZEMORE, J R.; LILLIE, R J.; BIRD, H R., AND DENTON, C. A. Further studies on the influence of aureomycin in the chick diet upon subsequent reproductive performance of laying hens, Poultry Sc. 34 432, 1955
43. JUKER, T H AND WILLIAMS, W L.: Nutritional effects of antibiotics, Pharm. Rev 5 381 1953.
44. LUCKEY T D Effect of feeding antibiotics upon the growth rate of germ-free birds. In a colloquium: Studies on the growth effect of antibiotics in germ-free animals held June 4, 1953, at Lobund Institute, University of Notre Dame, South Bend, Ind.
45. GYONGY P.; STOKES, J., JR.; SMITH, W H., AND GOLDBLATT H Studies on the use of aureomycin in hepatic disease. II. The effect of aureomycin on experimental dietary hepatic necrosis, Am. J Med. 220 8-11 1950.
46. CUNHA, T J BURNSIDE, J E. BUSCHMAN D M.; GLASSCOCK, R S., PEARSON A M., AND SHEALY A. L.: Effect of vitamin B₁₂, animal protein factor and soil for pig growth, Arch. Biochem. 23 3-4 1949
47. CATRON D V Recent developments in swine nutrition, Vet. Med. 44:215, 1949
48. SPEER, V C. VOHS, R L., CATRON D V., MADDOCK, H. M. AND CUTBERTSON, C. C. Effect of aureomycin and animal protein factor on healthy pigs, Arch. Biochem. 29 45., 1950.
49. TIERHILL, S W AND KRIDER, J L. A comparison of solvent soybean oil meals and the effect of adding aureomycin, dried whey product with whey fermentation solubles and APF in drylot rations of weanling pigs, J Animal Sc. 9 670, 1950
50. CATRON D V MADDOCK, H. M. SPEER, V C., AND VOHS, R L.: Effect of different levels of aureomycin with and without vitamin B₁₂ on growing fattening swine Antib. and Chemo. 1 31 1951
51. CARPENTER, L E. Antibiotics in swine nutrition, Iowa Vet. 22 15 1951.
52. COATES, M E., DICKINSON C. D HARRISON G F; KON S K. CUMMINS, S. H., AND CUTBERTSON W F J Mode of action of antibiotics in stimulating growth of chicks, Nature 168 332, 1951
53. COATES, M E. DICKINSON C D HARRISON G F KON S K. PORTER, J W G.; CUMMINS, S H. AND CUTBERTSON W F J A mode of action of antibiotics in chick nutrition, J Sc. Food Agric. 3 43, 1952.
54. BIRD, H R., LILLIE, R. J AND SIZEMORE, J R. Environment and stimulation of chick growth by antibiotics, Poultry Sc. 31 907 1952.
55. LILLIE, R. J SIZEMORE, J R., AND BIRD, H R. Environment and stimulation of growth of chicks by antibiotics Poultry Sc. 32 406, 1953.
56. HILL, D C. BRANTON H D AND SLINGER, S J: Influence of environment on the growth response of chicks to penicillin, Poultry Sc. 31 9-10 1952.
57. HILL, D C. BRANTON H D SLINGER, S J AND ANDERSON G. W: Influence of environment on the growth response of chicks to penicillin, Poultry Sc. 32 462, 1953.
58. KING, J O L., AND WHITE, E. G. The effects of penicillin supplements on the growth of chicks, Vet. Rec. 65 149 1953.
59. MORRISON A. B HUNSAKER, W G., AND ATTEREN J R. Influence of environment on the response of chicks to growth stimulants, Poultry Sc. 33 491 1954.

60. COATES, M. E.: The mode of action of antibiotics in animal nutrition, *Chem. & Indust.* 50 1333, 1953.
61. HILL, E. G., AND LARSON N. L. Effect of chlortetracycline supplementation on growth and feed utilization of unsuckled baby pigs obtained by hysterectomy *J. Animal Sc.* In press, 1953.
62. YOUNG, G. A., UNDERDAHL, N. R., AND HENZ, R. W.: Procurement of baby pigs by hysterectomy *Am. J. Vet. Res.* 16:123, 1953.
63. NEEDHAM, J.: *Chemical Embryology* London, Cambridge Univ. Press, 1931, Vol. I, p. 384.
64. JUKES, T. H., WILLIAMS, W. L., STORSTAD, E. L. R.; WONG, S. C., AND COX, H. R.: Lack of growth response to antibiotics by sterile chick embryos. Paper presented at Informal Poultry Nutrition Conference, New York, 1952.
65. LEFONK, P.; BARSKI, G., AND MAUREL, J.: Action of chloromycetin and of aureomycin on normal tissue cultures, *Proc. Soc. Exper. Biol. & Med.* 73 253, 1950.
66. GORDON H. A.: A morphological and biochemical approach. In a colloquium: Studies on the growth effect of antibiotics in germ-free animals, held June 4, 1952, at Lobund Institute, University of Notre Dame, South Bend, Ind.
67. COATES, M. E., DAVIES, M. K., AND KOK, S. K.: The effects of antibiotics on the intestine of the chick, *Brit. J. Nutrition* 9:110, 1953.
68. PEPPER, W. F.; SLENGER, S. J., AND MOTREK, I. Effect of aureomycin on the niacin and manganese requirement of chicks, *Poultry Sc.* 32 636, 1953.
69. ELLIOTT R. F., AND OWEN B.: Unpublished observations.
70. REYNOLDS, J. A.; TACKLER, P. C.; ERVIN, R. F.; WAGNER, M.; LUCKY T. K., AND GORDON H. A. Rearing germ-free chickens, *Lobund Rep.* 2:1 116, 1949.
71. LUCKY T. D.: Paper presented at the Informal Poultry Nutrition Conference, Federation Meetings, San Francisco, 1953.
72. SHERBURN, J. McN.; JEFFERY, J. J. HILL, E. G., AND CARPENTER, L. F. Some microbiological observations on the antibiotic-fed chick, *Poultry Sc.* 33:753, 1954.
73. KRATZER, F. H.; GRAU C. R.; STARR, M. P., AND REYNOLDS, D. M.: Growth-promoting activities of antibiotics and yeast cultures for chicks and turkey poult, *Federation Proc.* 10:386, 1951.
74. WILLIAMS, W. L.; TAYLOR, R. R. STORSTAD, E. L. R., AND JUKES, T. H.: Mechanism of the growth-promoting effect of aureomycin in chicks, *Federation Proc.* 10:270, 1951.
75. SHERBURN, J. M.; GUTERBACH, J.; MCGOWAN, J.; STEEN, J. R., AND SCHNEIDER, B. H.: Effect of antibiotics on intestinal microflora and on growth of turkeys and pigs, *Proc. Soc. Exper. Biol. & Med.* 76 15, 1951.
76. ELAM, J. F.; JACOBS, R. L.; FOWLER, J. AND COUCH, J. R.: Effect of dietary clostridia upon growth-promoting responses of penicillin, *Proc. Soc. Exper. Biol. & Med.* 85 643, 1954.
77. SCHENDEL, H. E.; BORG, A. F. AND JOHNSON, B. C.: The effect of antibiotics on the intestinal and cecal microflora of baby pigs fed a synthetic milk, *J. Animal Sc.* 13 904-911, 1954.
78. LINDGREN, N. O. Studies in the growth promoting action of antibiotics in poultry nutrition, *Nord. Vet. Med.* 6 701 706, 1954.
79. MARCH, B., AND BELY J.: The effect of feeding aureomycin on the bacterial content of chick feces, *Poultry Sc.* 31 177 178, 1952.

80. ROSENBERG, M. M.; MORIKAWA, W., AND BURNELL, O. A.: The effect of increasing concentrations of tetracycline on the growth and intestinal microflora of chicks, *Poultry Sc.* 31:708, 1952.
81. GUZMAN-GARCIA, J., SALLER, W. B., AND BAUMANN, C. A.: Microorganisms in the intestines of rats fed penicillin, *J. Nutrition* 49:647, 1953.
82. SKILVINT, A. V., AND LETHBRIDGE, H.: Influence of antibiotic feeding on intestinal flora in swine, *Mitteilungen Nr 3/4*, 1954.
83. ANDERSON, G. W., CUNNINGHAM, J. D., AND SLINGER, S. J.: Effect of tetracycline and certain phenylalanine acid derivatives on the growth and intestinal flora of turkey poult, *J. Nutrition* 48:539, 1952.
84. ELLSWORTH, S. A., HUFFMAN, C. F.; SMITH, C. K., AND RALSTON, N. P.: Effect of feeding antibiotics to dairy calves. I. Aureomycin and bacitracin feed supplements, *J. Sc. Food & Agric.* 5:183, 1954 abstr. in: *Mich. Agric. Exper. Sta. Quart. Bull.* 36:160, 1953.
85. SCALET, J. V.; KULP, W. L., AND CLEVELAND, R. C.: Effect of tetracycline on fecal microflora of rats. I. Interrelation of diet and tetracycline, *Proc. Soc. Exper. Biol. & Med.* 81:532, 1952.
86. KRATZER, F. H.; STARR, M. P.; REYNOLDS, D. M., AND GRAU, C. R.: Effects on growth of chicks and poult of feeding yeast isolated from streptomycin-fed poult, *Proc. Soc. Exper. Biol. & Med.* 81:812, 1952.
87. JOHANSSON, K. R.; PETERSON, G. E., AND DECK, E. C.: Effects of dietary aureomycin upon the intestinal microflora and the intestinal synthesis of vitamin B₁₂ in the rat, *J. Nutrition* 49:135, 1953.
88. ELAM, J. F.; GEE, L. L., AND COUCH, J. R.: Effect of feeding penicillin on the life cycle of the chick, *Proc. Soc. Exper. Biol. & Med.* 77:200, 1951.
89. ERECHTAKI, A., AND SANFORD, P.: The action of antibiotics on intestinal flora of poultry. I. Bacterial counts in chickens fed varying levels of aureofac., *Poultry Sc.* 32:837, 1953.
90. SLINGER, S. J., HAUSER, M. M., AND PEPPER, W. F.: The correlation between feed consumption and fecal flora in chicks, *Poultry Sc.* 32:923, 1953.
91. ANDERSON, G. W., SLINGER, S. J., AND PEPPER, W. F.: Effect of dietary microorganisms on the growth and cecal flora of chicks, *Poultry Sc.* 31:905, 1952.
92. BRANSON, H. D., AND ANDERSON, G. W.: Antibiotics and the growth of ducks, *Poultry Sc.* 31:908, 1952.
93. COOK, F. D., BLACKLEY, R. M., MACGREGOR, H. L., AND ANDERSON, G. W.: The influence of procaine penicillin on microflora in various intestinal loci of the turkey poult, *Poultry Sc.* 31:912, 1952.
94. BRIDGES, J. H., DYER, L. A., AND POWERS, J. J.: Penicillin and streptomycin affect the microflora of the intestinal tract of pigs, *J. Animal Sc.* 13:90, 1953.
95. SLINGER, S. J., HAUSER, M. M., AND PEPPER, W. F.: The correlation between feed consumption and fecal flora in chicks, *J. Nutrition* 52:73, 1954.
96. NOLAN, P. R.; STEPHENSON, E. L.; NELSON, T. S., AND TUCKER, D. L.: Response of baby pigs to antibiotics, *J. Animal Sc.* 10:1059, 1951.
97. LARSON, N. L., AND CARPENTER, L. E.: Effect of antibacterial substances on fecal clostridia populations, *Proc. Soc. Exper. Biol. & Med.* 79:227, 1953.
98. RUSOFF, L. L., ALFORD, J. A., AND HYDE, C. E.: Effect of type of protein on the

response of young dairy calves to aureomycin with data on the intestinal microflora, *J Dairy Sc.* 36:45 1953.

99. ANDERSON G. W., CUNNINGHAM, J. D. AND SLENGER, S. J.: Effect of aureomycin on the intestinal flora of chickens fed diets varying in protein content, *Canad. J. Comp. Med.* 17:83, 1953.
100. ROMONEN, G. L.; SROOG, M. S., COMBS, G. F., AND PELICAN, M. J.: Effect of antibiotics and diet composition on cecal bacteria and growth of chicks, *Antib. & Chem.* 2:42, 1952.
101. SCALETTI, J. V., KULP W. L., AND CLEVELAND, R. C.: Effect of tetracycline on fecal microflora of rats. II. Therapeutic level of tetracycline, *Proc. Soc. Exper. Biol. & Med.* 83:599, 1953.
102. RHODES, R. A., SABLE, W. B., MORGAN W. J., HANFORD, A. E., AND ELVEHJEM, C. A.: Stimulation and inhibition by antibiotics of intestinal bacteria in chicks, *J. Nutrition* 53:289 1954.
103. SMITH, C. F., JR.; CLEVELAND, R. C., KULP W. L., AND MATTHEWS, L. D.: Effect of dietary antibiotics on number of *Clostridium perfringens* in feces of chickens, *Antib. & Chem.* 2:363, 1952.
104. RIVERA, J. A., AND SROOG V. M.: The effect of tetracycline on the intestinal flora, *Gastroenterology* 17:546, 1951.
105. LOOMIS, J. K.; WAREHAMAN R. H., AND GALL, L. S.: Antibiotic studies with dairy calves, *J Dairy Sc.* 34:500 1951.
106. NEUMANN, A. L., SWAPP R. R., AND GALL, L. S.: The long-time effect of feeding aureomycin to fattening beef cattle, with bacteriological data, *J. Animal Sc.* 10:1058, 1951.
107. KESLER, E. M., AND KNOTT C. B.: Effect of tetracycline on growth and rumen function in Holstein calves, *J. Animal Sc.* 11:768, 1952.
108. HIBBS, J. W.; CONRAD, H. R., AND POUNDEN W. D.: A high roughage system for raising calves based on the early development of rumen function. V. Some effects of feeding aureomycin with different ratios of hay to grain, *J Dairy Sc.* 37:724, 1954.
109. CRANCK, C. M., DUNCAN, C. W., HUFFMAN C. F., AND LUECKE, R. W.: Antibiotics in rumen digestion and synthesis. II. The effect of aureomycin on the concentration of some amino acids and B vitamins in the rumen, *J Dairy Sc.* 36:495 1953.
110. CRANCK, C. M., SMITH, C. K., HUFFMAN C. F. AND DUNCAN C. W.: Antibiotics in rumen digestion and synthesis. III. The effect of aureomycin on rumen microorganisms with special reference to the streptococci and coliform groups, *J Dairy Sc.* 36:743, 1953.
111. HOOD L. H., JR., SWAPP R. R., AND GALL, L. S.: A study of the effects of antibiotics upon the digestion of feed nutrients by yearling steers with bacteriological data, *J. Animal Sc.* 12:921, 1953.
112. RADISSON J. J., BARTLEY E. E., FOUNTAINE, F. C., AND ATKINSON F. W.: Effect of aureomycin and/or grass juice concentrate on cellulose digestion in an artificial rumen, as influenced by different sources of nutrients, *J. Animal Sc.* 12:929, 1953.
113. RUNOFF L. L., FUSSELL, J. M., HYDE, C. E., AND CROWN R. M.: Parenteral administration of aureomycin to young calves with a note on mode of action, *J Dairy Sc.* 37:488, 1954.
114. LODGE, J. R., MILLS, F. T. AND JACOBSON N. L.: The effect of aureomycin feeding to dairy cattle on the *in vitro* digestion of cellulose by the rumen microorganisms, *J Dairy Sc.* 37:663, 1954.

115. HUITANEN C. V.; SAUNDERS, R. K., AND GALL, L. S. Fiber digestion using the miniature artificial rumen, *J Dairy Sc.* 37:328, 1954.
116. HUITANEN C. N.: Unpublished observations 1955
117. MANN S O, MASSON F M., AND OXFORD A. E.: Effect of feeding aureomycin to calves upon the establishment of their normal rumen microflora and microfauna, *Brit J Nutrition* 8:46, 1954
118. JOHANSSON, K. R. PETERSON G E., AND DICK, E. C.: Effects of parenteral and oral administration of antibiotics to rats fed a vitamin B₁₂ deficient diet, *Federation Proc.* 12 418, 1953.
119. ROMOSER, G L., SHORS M S AND COOPER, G. F. Effect of orally administered penicillin-resistant microorganisms on growth of chicks, *Proc. Soc. Exper Biol & Med.* 83:17 1953.
120. ANDERSON, G W.; SLINGER, S J AND PEPPER, W F: Bacterial cultures in the nutrition of poultry I. Effect of dietary bacterial cultures on the growth and cecal flora of chicks, *J Nutrition* 50:35 1953.
121. ANDERSON G W SLINGER, S. J.; PEPPER, W F AND HAUSER, M. M. Bacterial cultures in the nutrition of poultry II. Effect of dietary coliform cultures on the growth and cecal flora of poult, *J Nutrition* 50:47 1953.
122. HAUSER, M M., ANDERSON G W PEPPER, W F., AND SLINGER, S J: Further evidence on the relation of coliforms to the growth response of chicks to antibiotics, *Poultry Sc.* 33 21, 1954.
123. ROBERTS, R. S. The nutritional properties of killed *B coli*, *Brit. J Nutrition* 8 353 1954.
124. BLAYLOCK, L. G. VOIRA, P.; PATTERSON E. B AND MCGINNIS, J: Evidence for the production of chick growth factors by *Escherichia coli*, *Poultry Sc.* 33 6, 1954.
125. SCOTT M. L., MORRISON A. B AND NORMAN, L. C.: An unknown mineral required by the chick, *Federation Proc.* 14 278, 1955.
126. COUCH, J R.; REED, B L., CAMP A. A.; DANNENBURG, W N AND ROZACEK E. E.: Distillers dried solubles in the diets of chicks *Proc. 10th Distillers Feed Conference*, 47 1955.
127. WILSON J E., AND DUVERNEAUD, V L-penicillamine as a metabolic antagonist, *Science* 107 653, 1948.
128. ELAM, J F GEE, L L., AND COUCH J R. Function and metabolic significance of penicillin and bacitracin in the chick, *Proc. Soc. Exper Biol & Med.* 78 832, 1951
129. FELL, R. V AND STEPHENSON E. L. The effect of penicillin and penicillamine on chick growth when injected and when fed orally *Poultry Sc.* 32 1092, 1953.
130. TAYLOR, J H AND GORDON W S. *Nature*. In Press, 1955.
131. WILLIAMS, W L.; ESPARTE, R. G AND STOKSTAD, E. L. R.. Comparison of chick and microbiological methods for assay of penicillin in feeds, *Federation Proc.* 12 290, 1953.
132. WAINEL, P E. ABBOTT O J BAUMANN C. A., AND BIRD, H. R. Disappearance of the growth response of chicks to antibiotics in an "old" environment, *Poultry Sc.* 33 1141, 1954
133. LEBY D A., AND SCHAEFER, P J. Observations on growth responses to antibiotics and amino acids in poultry feeds, *Science* 111 33, 1955.

134. HANSON L. E., AND FERRIN E. F.: Comparative values of antibiotics and minerals for growing pigs, Univ. of Minnesota, Mimeograph H. 120, 1953.
135. RICHARDSON D. Antibiotics for growing fattening swine Circular 320, Kansas State College, 34, 1953.
136. STARR, M. P., AND RYDMOLD, D. M.: Streptomycin resistance of coliform bacteria from turkeys fed streptomycin, *Am. J. Pub. Health* 41 1875 1951.
137. SLINGER, S. J., AND PEPPER, W. F. The effect of antibiotics on the turkey poult need for supplementary biotin and pantothenic acid, *Poultry Sc.* 3:653, 1954.
138. FRENCH, C. E., URAAI, J. A., INGRAM, R. H., AND SWIFT R. W. The effect of high levels of tetracycline and streptomycin on longevity in the rat, *J. Nutrition* 54 73, 1954.
139. EMERSON, G. A., AND SMITH D. G. Induction of nutritional deficiency by oral administration of streptomycin, *J. Pharmacol. & Exper. Therap.* 85:536, 1945.
140. COMBES R. H., KEEFE, T. J., BURGER, R., AND MAW W. A.: Modification of the biochemical responses of the immature pullet to estrogen by means of dietary streptomycin, *Nature* 166:692, 1950.
141. BIRLY J., AND MARCH, B.: The effect of aureomycin and vitamins on the growth rate of chicks, *Science* 114:830, 1951.
142. OLSON J. J., HUTCHINGS, B. L., AND WHITEHILL, A. R.: The effect of feeding aureomycin on the vitamin B₁₂ requirement of the chick, *Arch. Biochem.* 29 334, 1950.
143. STOKSTAD, E. L. R., AND JONES, T. H.: Effect of various levels of vitamin B₁₂ upon growth response produced by aureomycin in chicks, *Proc. Soc. Exper. Biol. & Med.* 76:73, 1951.
144. WAIBEL, P. E.; CRAVEN, W. W., AND BAUMANN, C. A. Effect of dietary antibiotics on the growth of chicks receiving suboptimum amounts of thiamine, by mouth or by injection, *J. Nutrition* 50 441, 1953.
145. WAIBEL, P. E.; SUNDG, M. L., AND CRAVEN, W. W.: Effect of addition of penicillin to the hen's ration on biotin and folio acid content of eggs, *Poultry Sc.* 31:631, 1952.
146. LINDEWILER, H. BAUMANN C. A., AND SNELL, E. E.: Effect of aureomycin on the response of rats to various forms of vitamin B₆, *J. Nutrition* 43:565, 1951.
147. LEE, H., AND BAUMANN C. A. Effects of certain antibiotics on the growth of rats fed diets limited in thiamine, riboflavin, or pantothenic acid, *J. Nutrition* 43:565, 1951.
148. McDANIEL, E. G., AND DAY, F. S. Effects of penicillin and aureomycin on rats fed pyridoxine-deficient diets, *Federation Proc.* 14 443, 1955.
149. COATES, M. E. DICKINSON C. D. HARRISON, G. F. AND KOE, S. K.: The effect of antibiotics on the growth of chicks deprived of vitamins of the B complex, *Biochem. J.* 49 lxviii, 1951.
150. NELSON, T. S., AND SCOTT H. M. Niacin deficiency in the chick as influenced by antibiotics, *Poultry Sc.* 32 601 1953.
151. SARETT H. P.: Effect of oral administration of streptomycin on urinary excretion of B vitamins in man, *J. Nutrition* 47 275, 1952.
152. HALVY S.; DIAMANT E. J. AND GUCCIONE, K.: The effect of antibiotics on the metabolism of nicotinic acid, biotin and folio acid in rats, *Brit. J. Nutrition* 6:57 1953.
153. SAURELICH, H. E.: Effect of streptomycin and penicillin upon the vitamin requirements of the rat, *J. Nutrition* 48 99, 1952.
154. SCHERER, H. E., AND JOHNSON, B. C.: Studies of antibiotics in weanling rats ad-

ministered suboptimum levels of certain B vitamins orally and parenterally J Nutrition 54:461 1954

153. CHOW B. F.; DAVIS, R. L., AND DAVIS S.: The effect of antibiotics and the composition of diets on fecal vitamin B₁₂ and biotin, J Nutrition 49:657 1953.
156. PETERSON G. E.; DECK, E. C., AND JOHANSSON K. R.: Influence of dietary aureomycin and carbohydrate on growth, intestinal microflora and vitamin B₁₂ synthesis of the rat, J Nutrition 51 171 1953.
157. CHAVEZ-MUNOZ, J. PONCHER, H. G. AND WARMAN H. A.: Vitamin B₁₂ sparing action of aureomycin in the rat, Proc. Soc. Exper Biol. & Med. 77:18 1951
158. CATTON D. V., BENEDSON R. W.; MADDOCK, H. M.; ARNOLD G. C., AND HOSEYER, P. C.: Effects of certain antibiotics and vitamin B₁₂ on pantothenic acid requirements of growing fattening swine, J Animal Sc. 12 51 1953.
159. SHERFF B. E.; GRUMMER, R. H.; PHILLIPS, P. H., AND BOHNESTEDT G.: Comparison of growth responses of 2-day-old pigs to streptomycin, aureomycin, and crude APF alone and in combination with B₁₂, J Animal Sc. 11:97 1952.
160. WAHLSTROM, R. C., AND JOHNSON, B. C.: Effect of cortisone and of aureomycin on baby pigs fed a vitamin B₁₂ deficient diet, Proc. Soc. Exper Biol. & Med. 78 112, 1951.
161. HARTBOCK, E. W.; BATCHELOR, E., AND JOHNSON B. C.: Effect of aureomycin on utilization of vitamin A in the growing male albino rat, Proc. Soc. Exper Biol. & Med. 83 43, 1953.
162. MURRAY T. K., AND CAMPBELL, J. A.: Effect of aureomycin on apparent utilization of vitamin A by rat, Federation Proc. 14:446, 1955
163. BARNER, R. S.; BRAUER, R. KON S. K., AND MITCHELL, K. G.: Effect of supplementing an all-vegetable pig-fattening meal with antibiotics Chem. & Indust. 29 713, 1952.
164. BURGESS, R. C., GLUCK, M. AND LAUGHLAND D. H.: Biochemical changes in the penicillin-fed chick, Poultry Sc. 32 444, 1953.
165. HIGH, E. G.: Influence of aureomycin, penicillin and vitamin B₁₂ on metabolism of carotene and vitamin A in rat, Federation Proc. 14 437 1955
166. ROSS, E., AND YACOWITZ, H.: Effect of penicillin on growth and bone ash of chicks fed different levels of vitamin D and phosphorus, Poultry Sc. 33:262, 1954.
167. MICROSKY B. B. NIELSON, A. M. GLUCK, M. AND BURGESS R.: Penicillin and calcium absorption, Arch. Biochem. 34 479 1951
168. GARUTIN A. R., AND SHAFNER, C. S.: A study of the physiological mechanisms affecting specific gravity of chicken eggs, Poultry Sc. 31 917 1952.
169. BOGDANOFF P. D. JR., AND SHAFNER, C. S.: Antibiotics and calcium metabolism Poultry Sc. 33 7 1954.
170. LINDBLAD, G. S., SLINGER, S. J. AND MOTZOK, I.: Effect of aureomycin on the calcium and phosphorus requirements of chicks and poults Poultry Sc. 33:452, 1954.
171. PEPPER, W. F. SLINGER, S. J., AND MOTZOK, I.: Effect of aureomycin on the manganese requirement of chicks fed varying levels of salt and phosphorus, Poultry Sc. 31:1054 1952.
172. MACLEIN L. J., DENTON C. A.; KELLOGG, W. L., AND BIRD, H. R.: Effect of dietary antibiotic upon feed efficiency and protein requirement of growing chickens Poultry Sc. 31 106 1952.
173. WEAKLEY C. E., JR. HARK, J. G.; ANDERSON G. C. BLETHNER, J. K., AND MASON J. A.: Protein utilization studies with simplified rations I. Ad libitum feeding of low nitrogen rations, Poultry Sc. 32 927 1953.

174. SLEDOGER, S. J., BERKEY, J. E., PEPPER, W. F., SNYDER, E. S., AND ARTHUR, D.: Effect of antibiotics on the protein requirement of broilers, *Poultry Sc.* 31:757 1952.
175. SCOTT, H. M., GOFFY, E. A., AND GLITA, W. A.: The protein requirement of the chick as influenced by aureomycin, *Poultry Sc.* 31:751 1952.
176. SAKENA, H. C., STARR, M. E., BLAYLOCK, L. G., CARVER, J. S., AND MCGOWAN, J.: Effect of dietary penicillin on the efficiency of protein utilization by chicks, *Arch. Biochem.* 44:346, 1953.
177. SAKENA, H. C., BLAYLOCK, L. G., CARVER, J. S., AND MCGOWAN, J.: Factors affecting the growth response of turkey poult to antibiotics, *Poultry Sc.* 32:788, 1953.
178. SLEDOGER, S. J., GARTLEY, K. M., PEPPER, W. F., AND HILL, D. C.: The influence of animal protein factor supplements and antibiotics on the incidence and severity of white feathers in turkeys, *J. Nutrition* 43:345 1951.
179. BERKEY, J., MARSH, B. E., AND SMITH, D. F.: The effect of feeding antibiotics on the carbohydrate and protein requirements of the chick, *Poultry Sc.* 31:863, 1952.
180. MATTERSON, L. D., SINGH, E. P., DECKER, L., AND KOFFER, A.: The influence of protein level, aureomycin and vitamin B₁₂ on the growth and feed efficiency of chicks, *Poultry Sc.* 31:940 1952.
181. WHITE-STEVENS, R., AND ZIEGLER, H. G.: Chlorotetracycline in relation to protein level of diet on growth efficiency of white rock broilers, *Poultry Sc.* 33:50, 1954.
182. CATRON, D. V., JENSEN, A. H., HOMER, P. G., MADDOCK, H. M., AND ASHTON, G. C.: Re-evaluation of protein requirements of growing fattening swine as influenced by feeding an antibiotic, *J. Animal Sc.* 11:221, 1952.
183. JENSEN, A. H., ACKER, D. C., MADDOCK, H. M., ASHTON, G. C., HOMER, P. G., HEADY, E. O., AND CATRON, D. V.: Different protein levels with and without antibiotics for growing-finishing swine: effect on growth rate and feed efficiency, *J. Animal Sc.* 14:69 1955.
184. BLACK, A., AND BRATZLER, J. W.: The effects of a vitamin B₁₂ supplement, vitamin B₁₂ and streptomycin on the metabolism of the rat, *J. Nutrition* 47:159, 1952.
185. KNOX, L. K., AND BLACK, A.: The influence of vitamin B₁₂ and antibiotics on protein and energy utilization in a low protein diet, *J. Nutrition* 48:477 1952.
186. CARROLL, R. W., HEMLEY, G. W., SUTLER, C. L., WILCOX, E. L., AND GRAHAM, W. R.: Absorption of nitrogen and amino acids from soybean meal as affected by heat treatment or supplementation with aureomycin and methionine, *Arch. Biochem.* 45:280, 1953.
187. HEMLEY, G. W., CARROLL, R. W., WILCOX, E. L., AND GRAHAM, W. R.: The effects of aureomycin and methionine supplements fed to rats receiving soybean meals, *Arch. Biochem.* 45:270, 1953.
188. BERRY, M. E., AND SCHUCK, C.: The effect of aureomycin on growth and protein utilization in the rat, *J. Nutrition* 54:271, 1954.
189. FORBES, R. M.: Studies on the influence of antibiotics and methionine on nitrogen utilization and basal metabolism of the growing male albino rat, *J. Nutrition* 53:374, 1954.
190. HARTBROOK, E. W., AND JOHNSON, B. C.: Effects of dietary terramycin and methionine supplements on fat and protein gains in weanling rats, *J. Nutrition* 51:205, 1953.
191. MEADE, R. J., AND FORBES, R. M.: The influence of aureomycin and vitamin B₁₂ alone and in combination on nitrogen utilization by growing swine, *J. Animal Sc.* In press, 1955.

192. MRADE, R. J. The influence of protein content of the diet and of aureomycin and/or vitamin B₁₂ supplementation upon performance of growing fattening pigs, *J. Animal Sc. In press*, 1955.
193. BROWN, P. B.; BECKER, D. E.; TERRY, S. W., AND CARD, L. E.: The effect of aureomycin on the growth and metabolism of the pig, *Arch. Biochem.* 41:378 1952.
194. RUSSO, J. M.; HANSON, L. E., AND JENSEN, J. J. Effects of aspartic acid and aureomycin on nitrogen balance in pigs, *Scientific Journal Series of the Minnesota Agricultural Experiment Station, J. Animal Sc. In press*, 1955.
195. GALK, E. F., AND FOLKER, J. P. The assimilation of amino acids by bacteria: 15. Action of antibiotics on nucleic acids and protein synthesis in *S. aureus* *Biochem. J.* 53:493, 1953.
196. LAMSTER, C. W.: Antibiotics as growth stimulants for dairy cattle *J. Dairy Sc. In press*, 1953.
197. CATRON, D. V. Antibiotics in swine nutrition, paper presented at the University of Wisconsin, Nutrition Conference 1953.
198. STOKSTAD, E. L. R., AND JUICK, T. H.: Growth-promoting effect of aureomycin on turkey poult, *Poultry Sc.* 29:611 1950.
199. MCGOWAN, J., BIRD, L. R., STERN, J. R., WILCOX, R. A., AND BEARIE, G. E.: The effect of aureomycin and streptomycin on growth of chicks and turkey poult, *Poultry Sc.* 29:771 1950.
200. MCGOWAN, J.; STERN, J. R.; WILCOX, R. A., AND CARVER, J. S. The effect of different antibiotics on growth of turkey poult, *Poultry Sc.* 30:492, 1951.
201. KRAMER, E. H., AND FRITZ, J. C. The response of chicks and poult to low levels of antibiotics, *Poultry Sc.* 30:921 1951.
202. WILLIAMS, O. M. AND HILL, J. E. The effect of antibiotics on the growth of two varieties of turkey poult, *Poultry Sc.* 31:769 1952.
203. HEUSER, G. F. AND NORRIS, L. C. Some results of feeding antibiotics to chickens, *Poultry Sc.* 31:857 1952.
204. LEWIS, R. W., AND SANFORD, P. E. Use of equivalent levels of antibiotic and vitamin B₁₂ supplements with cottonseed and soybean oil meals in chick diets, *Poultry Sc.* 32:268, 1953.
205. LILLIE, R. J. AND BIRD, H. R. Evaluation of antibiotic feed supplements and crystalline antibiotics in chick rations, *Poultry Sc.* 32:531 1953.
206. BRANSON, H. D., ANDERSON, G. W., AND HILL, D. C. Antibiotics and the growth of ducks, *Poultry Sc.* 32:835 1953.
207. RUNNELS, T. D., REYNOLDS, J. W., AND WALLER, E. F. The influence of various levels of penicillin and streptomycin on rate of growth and feed efficiency in broiler diets, *Poultry Sc.* 30:929 1951.
208. PEPPER, H. J.; ORENG, E. B.; BENEDICT, R. G., AND LEONOWITZ, L. A. The effect of feeding crude polymyxin D on the intestinal bacteria of chickens, *Poultry Sc.* 29:520, 1950.
209. BRANSON, H. D., AND HILL, D. C.: The comparative effect of antibiotics on the growth of poult, *Poultry Sc.* 30:793, 1951.
210. STERN, J. R.; SHERBURN, J. McN., AND MCGOWAN, J. Lack of growth response of turkey poult to certain antibiotics and bacteriostatic agents, *Poultry Sc.* 31:179 1952.
211. PEPPER, H. T., AND RODGER, N. E.: Effect of streptothricin on chick growth, *Poultry Sc.* 31:1104, 1952.

174. SLINGER, S. J. BERKEY J. E.; PEPPER, W. F. SNYDER, E. S., AND ARTHUR, D.: Effect of antibiotics on the protein requirement of broilers, *Poultry Sc.* 31:757 1952.
175. SCOTT H. M., GOTTI, E. A., AND GLETA, W. A. The protein requirement of the chick as influenced by aureomycin, *Poultry Sc.* 31 751 1952.
176. SAXENA, H. C. STARR, M. E. BLATLOCK, L. G., CARVER, J. S., AND MCGROVER, J.: Effect of dietary penicillin on the efficiency of protein utilization by chicks, *Arch. Biochem.* 44 846, 1953.
177. SAXENA, H. C.; BLATLOCK, L. G. CARVER, J. S., AND MCGROVER, J. Factors affecting the growth response of turkey poult to antibiotics, *Poultry Sc.* 32 788, 1953.
178. SLINGER, S. J.; GARTLEY K. M. PEPPER, W. F., AND HILL, D. C.: The influence of animal protein factor supplements and antibiotics on the incidence and severity of white feathers in turkeys, *J. Nutrition* 43 345 1951.
179. BIELY J. MARCH, B. E., AND SMITH, D. F. The effect of feeding antibiotics on the carbohydrate and protein requirements of the chick, *Poultry Sc.* 31:803 1952.
180. MATTHEWSON L. D.; SIMON E. P., DECKER, L., AND KOSKOFF A.: The influence of protein level, aureomycin and vitamin B₁₂ on the growth and feed efficiency of chicks, *Poultry Sc.* 31:940 1952.
181. WHITE-STEVENS, R., AND ZIEDEL, H. G. Chlortetracycline in relation to protein level of diet on growth efficiency of white rock broilers, *Poultry Sc.* 33 50, 1954.
182. CATROW D. V. JENSEN A. H., HOMETER, P. G. MADDOCK, H. M., AND ASHTON G. C.: Re-evaluation of protein requirements of growing fattening swine as influenced by feeding an antibiotic, *J. Animal Sc.* 11:221 1952.
183. JENSEN, A. H. ACKER, D. C. MADDOCK, H. M.; ASHTON G. C. HOMETER, P. G.; HEADY E. O. AND CATROW D. V. Different protein levels with and without antibiotics for growing finishing swine effect on growth rate and feed efficiency *J. Animal Sc.* 14:69 1953.
184. BLACK, A., AND BRATZLER, J. W. The effects of a vitamin B₁₂ supplement, vitamin B₁₂ and streptomycin on the metabolism of the rat, *J. Nutrition* 47 159, 1952.
185. KNOEREL, L. K., AND BLACK, A. The influence of vitamin B₁₂ and antibiotics on protein and energy utilization in a low protein diet, *J. Nutrition* 48:477 1952.
186. CARROLL, R. W.; HENKLEY G. W. SITTIER, C. L., WILCOX, E. L., AND GRAHAM, W. R.: Absorption of nitrogen and amino acids from soybean meal as affected by heat treatment or supplementation with aureomycin and methionine, *Arch. Biochem.* 45 260 1953.
187. HENKLEY G. W. CARROLL, R. W. WILCOX, E. L., AND GRAHAM, W. R.: The effects of aureomycin and methionine supplements fed to rats receiving soybean meal, *Arch. Biochem.* 45:270 1953.
188. BERRY M. E. AND SCHUCK, C. The effect of aureomycin on growth and protein utilization in the rat, *J. Nutrition* 54 271 1954.
189. FORMER, R. M. Studies on the influence of antibiotics and methionine on nitrogen utilization and basal metabolism of the growing male albino rat, *J. Nutrition* 53 275, 1954.
190. HARTROCK, E. W., AND JOHNSON B. C. Effects of dietary terramycin and methionine supplements on fat and protein gains in weanling rats, *J. Nutrition* 51:205, 1953.
191. MEADE, R. J., AND FORMER, R. M.: The influence of aureomycin and vitamin B₁₂ alone and in combination on nitrogen utilization by growing swine, *J. Animal Sc.* In press, 1955.

192. MEADE, R. J. The influence of protein content of the diet and of aureomycin and/or vitamin B₁₂ supplementation upon performance of growing fattening pigs. *J. Animal Sc.* In press, 1955.
193. BROWN, P. B.; BECKER, D. E., TERRILL, S. W., AND CARD, L. E.: The effect of aureomycin on the growth and metabolism of the pig. *Arch. Biochem.* 41: 878, 1952.
194. RUSSO, J. M., HANBOM, L. E., AND JEZESEKI, J. J.: Effects of aspartic acid and aureomycin on nitrogen balance in pigs. *Scientific Journal Series of the Minnesota Agricultural Experiment Station, J. Animal Sc.* In press, 1955.
195. GALE, E. F. AND FOLKES, J. P.: The assimilation of amino acids by bacteria: 15. Action of antibiotics on nucleic acids and protein synthesis in *S. aureus*. *Biochem. J.* 53: 493, 1953.
196. LAMETER, C. W. Antibiotics as growth stimulants for dairy cattle. *J. Dairy Sc.* In press, 1955.
197. CATRON, D. V. Antibiotics in swine nutrition, paper presented at the University of Wisconsin, Nutrition Conference, 1953.
198. STOKES, E. L. R., AND JUKES, T. H. Growth-promoting effect of aureomycin on turkey poult, *Poultry Sc.* 29: 811, 1950.
199. MCGOWAN, J., BING, L. R., STERN, J. R., WILCOX, R. A., AND BEARSE, G. E. The effect of aureomycin and streptomycin on growth of chicks and turkey poult, *Poultry Sc.* 29: 771, 1950.
200. MCGOWAN, J., STERN, J. R., WILCOX, R. A., AND CARVER, J. S. The effect of different antibiotics on growth of turkey poult, *Poultry Sc.* 30: 422, 1951.
201. KRAMER, E. H., AND FRITZ, J. C. The response of chicks and poult to low levels of antibiotics, *Poultry Sc.* 30: 921, 1951.
202. WILLIAMS, O. M. AND HILL, J. E. The effect of antibiotics on the growth of two varieties of turkey poult, *Poultry Sc.* 31: 788, 1952.
203. HEUSER, G. F., AND NORRIS, L. C. Some results of feeding antibiotics to chickens, *Poultry Sc.* 31: 857, 1952.
204. LEWIS, R. W. AND SANFORD, P. E. Use of equivalent levels of antibiotic and vitamin B₁₂ supplements with cottonseed and soybean oil meals in chick diets, *Poultry Sc.* 32: 268, 1953.
205. LITTLE, R. J. AND BIRD, H. R. Evaluation of antibiotic feed supplements and crystalline antibiotics in chick rations, *Poultry Sc.* 32: 531, 1953.
206. BRANTON, H. D., ANDERSON, G. W. AND HILL, D. C. Antibiotics and the growth of ducks, *Poultry Sc.* 32: 335, 1953.
207. RUNNELS, T. D., REYNOLDS, J. W. AND WALLER, E. F. The influence of various levels of penicillin and streptomycin on rate of growth and feed efficiency in broiler diets, *Poultry Sc.* 30: 929, 1951.
208. PEPPER, H. J., OBERG, E. B., BIEDERT, R. G. AND LINDENFELDER, L. A.: The effect of feeding crude polymyxin D on the intestinal bacteria of chickens, *Poultry Sc.* 29: 520, 1950.
209. BRANTON, H. D. AND HILL, D. C. The comparative effect of antibiotics on the growth of poult, *Poultry Sc.* 30: 793, 1951.
210. STERN, J. R., SHERBURN, J. MCN., AND MCGOWAN, J.: Lack of growth response of turkey poult to certain antibiotics and bacteriostatic agents, *Poultry Sc.* 31: 179, 1952.
211. PEZZER, H. T. AND RODGERS, N. E.: Effect of streptothricin on chick growth, *Poultry Sc.* 31: 1104, 1952.

174. SLINGER, S J; BERGLEY J E., PEPPER, W F.; SWYDER, E. S AND ARTHUR, D.: Effect of antibiotics on the protein requirement of broilers, *Poultry Sc.* 31 757 1952.
175. SCOTT H M., GOFFI, E A., AND GLETA, W A.: The protein requirement of the chick as influenced by aureomycin, *Poultry Sc.* 31 751, 1952.
176. SAKURA, H C.; STARR, M. E.; BLATLOCK, L. G., CARVER, J S., AND MCGINNIS, J: Effect of dietary penicillin on the efficiency of protein utilization by chicks, *Arch. Biochem.* 44 346, 1953.
177. SAKURA, H. C.; BLATLOCK, L. G. CARVER, J S., AND MCGINNIS, J Factors affecting the growth response of turkey poult to antibiotics, *Poultry Sc.* 32 788, 1953.
178. SLINGER, S J GARTLEY K. M. PEPPER, W F., AND HILL, D C.: The influence of animal protein factor supplements and antibiotics on the incidence and severity of white feathers in turkeys, *J Nutrition* 43 345 1951.
179. BELY J; MARSH, B. E., AND SMITH, D F The effect of feeding antibiotics on the carbohydrate and protein requirements of the chick, *Poultry Sc.* 31 803, 1952.
180. MATTERSON L. D., SIMON E. P., DECKER, L., AND KOSKOFF A.: The influence of protein level, aureomycin and vitamin B₁₂ on the growth and feed efficiency of chicks, *Poultry Sc.* 31 940 1952.
181. WHITE-STEVENS, R., AND ZETHEL, H. G Chlortetracycline in relation to protein level of diet on growth efficiency of white rock broilers, *Poultry Sc.* 33 50, 1954.
182. CATROW, D V., JENKIN A. H. HOMETER, P G MADDOCK, H. M., AND ARTHUR G. C.: Re-evaluation of protein requirements of growing-fattening swine as influenced by feeding an antibiotic, *J Animal Sc.* 11 281 1952.
183. JENKIN, A. H., ACKER, D C.; MADDOCK, H. M.; ARTHUR G. C.; HOMETER, P G., HEARD E. O AND CATROW D V: Different protein levels with and without antibiotics for growing finishing swine: effect on growth rate and feed efficiency *J Animal Sc.* 14 69 1953.
184. BLACK, A., AND BRATZLER, J W The effects of a vitamin B₁₂ supplement, vitamin B₁₂ and streptomycin on the metabolism of the rat, *J Nutrition* 47:159 1952.
185. KNOEREL, L. K., AND BLACK, A.: The influence of vitamin B₁₂ and antibiotics on protein and energy utilization in a low protein diet, *J Nutrition* 48:477 1952.
186. CARROLL, R. W HENLEY G W; STYLER, C. L., WILCOX, E. L., AND GRAHAM, W R.: Absorption of nitrogen and amino acids from soybean meal as affected by heat treatment or supplementation with aureomycin and methionine, *Arch. Biochem.* 45 280, 1953.
187. HENLEY G. W CARROLL, R W; WILCOX, E. L., AND GRAHAM, W R. The effects of aureomycin and methionine supplements fed to rats receiving soybean meals, *Arch. Biochem.* 45 270, 1953.
188. BERRY M. E. AND SCHUCK, C.: The effect of aureomycin on growth and protein utilization in the rat, *J Nutrition* 54 271 1954.
189. FORMER, R. M. Studies on the influence of antibiotics and methionine on nitrogen utilization and basal metabolism of the growing male albino rat, *J Nutrition* 53 275, 1954.
190. HARTBROOK, E. W., AND JOHNSON B. C. Effects of dietary terramycin and methionine supplements on fat and protein gains in weanling rats, *J Nutrition* 51 205, 1953.
191. MEADE, R. J AND FORMER, R. M The influence of aureomycin and vitamin B₁₂ alone and in combination on nitrogen utilization by growing swine, *J Animal Sc.* In press, 1955.

192. MEADE, R. J.: The influence of protein content of the diet and of aureomycin and/or vitamin B₁₂ supplementation upon performance of growing fattening pigs. *J. Animal Sc.* In press, 1955.
193. BROWN, P. B.; BECKER, D. E. TETULL, S. W., AND CARD, L. E.: The effect of aureomycin on the growth and metabolism of the pig. *Arch. Biochem.* 41:378 1952.
194. RUSSO, J. M., HANSON, L. E., AND JENSEN, J. J.: Effects of arsenic acid and aureomycin on nitrogen balance in pigs, Scientific Journal Series of the Minnesota Agricultural Experiment Station, *J. Animal Sc.* In press, 1955.
195. GALK, E. F., AND FOLKES, J. P.: The assimilation of amino acids by bacteria: 15 Action of antibiotics on nucleic acids and protein synthetase in *S. aureus*. *Biochem. J.* 53 493, 1953.
196. LASSITER, C. W.: Antibiotics as growth stimulants for dairy cattle, *J. Dairy Sc.* In press, 1953.
197. CATRON, D. V.: Antibiotics in swine nutrition, paper presented at the University of Wisconsin, Nutrition Conference, 1953.
198. STOKES, E. L. R., AND JONES, T. H.: Growth-promoting effect of aureomycin on turkey poult, *Poultry Sc.* 29 611, 1950.
199. MCGOWAN, J.; BERG, L. R. STERN, J. R. WILCOX, R. A., AND BEARSE, G. E.: The effect of aureomycin and streptomycin on growth of chicks and turkey poult, *Poultry Sc.* 29 771 1950.
200. MCGOWAN, J., STERN, J. R. WILCOX, R. A., AND CARVER, J. S.: The effect of different antibiotics on growth of turkey poult, *Poultry Sc.* 30 492, 1951.
201. KRAMER, E. H., AND FRITZ, J. C.: The response of chicks and poult to low levels of antibiotics, *Poultry Sc.* 30 921 1951.
202. WILLIAMS, O. M., AND HILL, J. E.: The effect of antibiotics on the growth of two varieties of turkey poult, *Poultry Sc.* 31 760 1952.
203. HEUER, C. F. AND NORMAN, L. C.: Some results of feeding antibiotics to chickens, *Poultry Sc.* 31 857 1952.
204. LEWIS, R. W., AND SANFORD, P. E.: Use of equivalent levels of antibiotic and vitamin B₁₂ supplements with cottonseed and soybean oil meals in chick diets, *Poultry Sc.* 32:268, 1953.
205. LILLIE, R. J. AND BIRD, H. R.: Evaluation of antibiotic feed supplements and crystalline antibiotics in chick rations, *Poultry Sc.* 32 531, 1953.
206. BRADY, H. D.; ANDERSON, G. W. AND HILL, D. C.: Antibiotics and the growth of ducks, *Poultry Sc.* 32:335 1953.
207. RUNNELS, T. D. REYNOLDS, J. W., AND WALLER, E. F.: The influence of various levels of penicillin and streptomycin on rate of growth and feed efficiency in broiler diets. *Poultry Sc.* 30:929 1951.
208. PEPPLER, H. J.; OJENG, E. B.; BENEDICT, R. G. AND LINDENFELDER, L. A.: The effect of feeding crude polymyxin D on the intestinal bacteria of chickens, *Poultry Sc.* 29 520, 1950.
209. BRADY, H. D. AND HILL, D. C.: The comparative effect of antibiotics on the growth of poult, *Poultry Sc.* 30 783, 1951.
210. STERN, J. R.; SIZEMUTH, J. MCN., AND MCGOWAN, J.: Lack of growth response of turkey poult to certain antibiotics and bacteriostatic agents, *Poultry Sc.* 31:179 1952.
211. PEPPLER, H. T., AND RODGERS, N. E.: Effect of streptothricin on chick growth, *Poultry Sc.* 31:1104, 1952.

174. SLINGER, S. J.; BENNETT J. E. PEPPER, W. F.; SNYDER, E. S., AND ARTHUR, D.: Effect of antibiotics on the protein requirement of broilers, *Poultry Sc.* 31:757 1952.
175. SCOTT H. M. GOFFI, E. A., AND GLINTA, W. A.: The protein requirement of the chick as influenced by aureomycin, *Poultry Sc.* 31 751, 1952.
176. SAKENA, H. C.; STANE, M. E. BLATLOCK, L. G., CARVER, J. S., AND MCGONNIE, J.: Effect of dietary penicillin on the efficiency of protein utilization by chicks, *Arch. Biochem.* 44 346, 1953.
177. SAKENA, H. C.; BLATLOCK, L. G. CARVER, J. S., AND MCGONNIE, J. Factors affecting the growth response of turkey poult to antibiotics, *Poultry Sc.* 32 768, 1953.
178. SLINGER, S. J. GARTLEY K. M., PEPPER, W. F. AND HILL, D. C.: The influence of animal protein factor supplements and antibiotics on the incidence and severity of white feathers in turkeys, *J Nutrition* 43 845 1951.
179. BIRLY J. MANCH, B. E., AND SMITH, D. F.: The effect of feeding antibiotics on the carbohydrate and protein requirements of the chick, *Poultry Sc.* 31 863, 1952.
180. MATTERSON L. D. SINGER, E. P. DECKER, L., AND KOSOFF A.: The influence of protein level, aureomycin and vitamin B₁₂ on the growth and feed efficiency of chicks, *Poultry Sc.* 31:940, 1952.
181. WHITE-STEVENS, R., AND ZIEBEL, H. G. Chlortetracycline in relation to protein level of diet on growth efficiency of white rock broilers, *Poultry Sc.* 33 50, 1954.
182. CATTON D. V. JENSEN A. H., HOMER, P. G. MADDOCK, H. M., AND ARNTON, G. C.: Re-evaluation of protein requirements of growing fattening swine as influenced by feeding an antibiotic, *J Animal Sc.* 11 221 1952.
183. JENSEN, A. H.; ACKER, D. C.; MADDOCK, H. M.; ARNTON G. C. HOMER, P. G., HEADY E. O. AND CATTON D. V.: Different protein levels with and without antibiotics for growing finishing swine: effect on growth rate and feed efficiency *J Animal Sc.* 14 69 1953
184. BLACK, A., AND BRATZLER, J. W.: The effects of a vitamin B₁₂ supplement, vitamin B₁₂ and streptomycin on the metabolism of the rat, *J Nutrition* 47:159 1952.
185. KNOEREL, L. K., AND BLACK, A.: The influence of vitamin B₁₂ and antibiotics on protein and energy utilization in a low protein diet, *J Nutrition* 48:477 1952.
186. CANNOLL, R. W. HEMLEY G. W.; SEITLER, C. L. WILCOX, E. L., AND GRAHAM, W. R.: Absorption of nitrogen and amino acids from soybean meal as affected by heat treatment or supplementation with aureomycin and methionine, *Arch. Biochem.* 45 260, 1953.
187. HEMLEY G. W., CANNOLL, R. W.; WILCOX, E. L., AND GRAHAM, W. R. The effects of aureomycin and methionine supplements fed to rats receiving soybean meal, *Arch. Biochem.* 45 270, 1953.
188. BERRY M. E. AND SCHUCK, C.: The effect of aureomycin on growth and protein utilization in the rat, *J Nutrition* 54 271, 1954.
189. FORBES, R. M. Studies on the influence of antibiotics and methionine on nitrogen utilization and basal metabolism of the growing male albino rat, *J Nutrition* 53:275, 1954.
190. HARTBROOK, E. W., AND JOHNSON B. C. Effects of dietary terramycin and methionine supplements on fat and protein gains in weanling rats, *J Nutrition* 51:205, 1953.
191. MEADE, R. J., AND FORBES, R. M. The influence of aureomycin and vitamin B₁₂ alone and in combination on nitrogen utilization by growing swine, *J Animal Sc.* In press, 1955.

192. MEADE, R. J.: The influence of protein content of the diet and of aureomycin and/or vitamin B₁₂ supplementation upon performance of growing fattening pigs, *J Animal Sc.* In press, 1953
193. BROWN P. B.; BECKER, D. E.; TERRILL, S. W., AND CARD, L. E.: The effect of aureomycin on the growth and metabolism of the pig, *Arch. Biochem.* 41:378, 1952.
194. RUSSO J. M. HANSON, L. E., AND JERESKI, J. J. Effects of arsenic acid and aureomycin on nitrogen balance in pigs, *Scientific Journal Series of the Minnesota Agricultural Experiment Station*, *J Animal Sc.* In press, 1953
195. GALE, E. F. AND FOLKER, J. P. The assimilation of amino acids by bacteria: 15. Action of antibiotics on nucleic acids and protein synthesis in *S aureus* *Biochem. J.* 53 493, 1953.
196. LASSITER, C. W. Antibiotics as growth stimulants for dairy cattle *J Dairy Sc.* In press, 1953
197. CATRON D. V. Antibiotics in swine nutrition, paper presented at the University of Wisconsin, Nutrition Conference, 1953.
198. STOKSTAD, E. L. R., AND JUKES, T. H. Growth-promoting effect of aureomycin on turkey poult, *Poultry Sc.* 29 811 1950
199. MCGOWAN, J.; BENO, L. R. STERN J. R. WILCOX, R. A., AND BEARER, G. E.: The effect of aureomycin and streptomycin on growth of chicks and turkey poult, *Poultry Sc.* 29 771, 1950.
200. MCGOWAN, J. STERN J. R., WILCOX, R. A., AND CARVER, J. S. The effect of different antibiotics on growth of turkey poult, *Poultry Sc.* 30:492, 1951
201. KRAMER, E. H., AND FRITZ, J. C. The response of chicks and poult to low levels of antibiotics, *Poultry Sc.* 30:921, 1951.
202. WILLIAMS, O. M., AND HILL, J. E. The effect of antibiotics on the growth of two varieties of turkey poult, *Poultry Sc.* 31 760 1952.
203. HEUSER, C. F., AND NORRIS, L. C.: Some results of feeding antibiotics to chickens, *Poultry Sc.* 31 857 1952.
204. LEWIS, R. W., AND SANFORD P. E. Use of equivalent levels of antibiotic and vitamin B₁₂ supplements with cottonseed and soybean oil meals in chick diets, *Poultry Sc.* 32:268, 1953.
205. LILLIE, R. J. AND BIRD, H. R. Evaluation of antibiotic feed supplements and crystal line antibiotics in chick rations, *Poultry Sc.* 32 531 1953.
206. BRADON H. D. ANDERSON G. W. AND HILL, D. C.: Antibiotics and the growth of ducks, *Poultry Sc.* 32:335 1953.
207. RUMMEL, T. D.; RETNOLDS, J. W. AND WALLER, E. F.: The influence of various levels of penicillin and streptomycin on rate of growth and feed efficiency in broiler diets, *Poultry Sc.* 30 929 1951.
208. PEPPER, H. J.; OHSANG, E. B.; BENEDICT R. G., AND LEDENFELDER, L. A.: The effect of feeding crude polymyxin D on the intestinal bacteria of chickens, *Poultry Sc.* 29 530, 1950.
209. BRADON, H. D. AND HILL, D. C. The comparative effect of antibiotics on the growth of poult, *Poultry Sc.* 30:793 1951
210. STERN J. R.; SIEBERTIL, J. McN., AND MCGOWAN, J.: Lack of growth response of turkey poult to certain antibiotics and bacteriostatic agents, *Poultry Sc.* 31:179 1952.
211. PEPPER, H. T., AND RODGERS, N. E. Effect of streptothricin on chick growth, *Poultry Sc.* 31:1104, 1952.

174. SLINGER, S. J. BENNEY, J. E. PEPPER, W. F.; SUTHER, E. S., AND ARTHUR, D.: Effect of antibiotics on the protein requirement of broilers, *Poultry Sc.* 31:757 1952.
175. SCOTT, H. M. GOFFI, E. A. AND GLIEBA, W. A.: The protein requirement of the chick as influenced by aureomycin, *Poultry Sc.* 31:751 1952.
176. SAKENA, H. C.; STARK, M. E.; BLAYLOCK, L. G., CARVER, J. S., AND MCGINNIS, J.: Effect of dietary penicillin on the efficiency of protein utilization by chicks, *Arch. Biochem.* 44:348, 1953.
177. SAKENA, H. C.; BLAYLOCK, L. G. CARVER, J. S., AND MCGINNIS, J. Factors affecting the growth response of turkey poults to antibiotics, *Poultry Sc.* 32:758, 1953.
178. SLINGER, S. J.; GARTLEY, K. M., PEPPER, W. F. AND HILL, D. C.: The influence of animal protein factor supplements and antibiotics on the incidence and severity of white feathers in turkeys, *J. Nutrition* 43:845 1951.
179. BULL, J. MARCH, B. E., AND SMITH, D. F. The effect of feeding antibiotics on the carbohydrate and protein requirements of the chick, *Poultry Sc.* 31:863, 1952.
180. MATTHEWSON, L. D. SINGH, E. P. DECKER, L., AND KOSOFF, A.: The influence of protein level, aureomycin and vitamin B₁₂ on the growth and feed efficiency of chicks, *Poultry Sc.* 31:840, 1952.
181. WHITE-STEVENS, R., AND ZEIBEL, H. G.: Chlorotetracycline in relation to protein level of diet on growth efficiency of white rock broilers. *Poultry Sc.* 33:50, 1954.
182. CATRON, D. V. JENSEN, A. H. HOMER, P. G. MADDOCK, H. M. AND ARNTON, G. C.: Re-evaluation of protein requirements of growing fattening swine as influenced by feeding an antibiotic, *J. Animal Sc.* 11:221, 1952.
183. JENSEN, A. H. ACKER, D. C.; MADDOCK, H. M.; ARNTON, G. C. HOMER, P. G., HEADY, E. O. AND CATRON, D. V.: Different protein levels with and without antibiotics for growing-finishing swine: effect on growth rate and feed efficiency. *J. Animal Sc.* 14:69 1953.
184. BLACK, A., AND BRATLER, J. W.: The effects of a vitamin B₁₂ supplement, vitamin B₁₂ and streptomycin on the metabolism of the rat, *J. Nutrition* 47:159 1952.
185. KROEMER, L. K., AND BLACK, A.: The influence of vitamin B₁₂ and antibiotics on protein and energy utilization in a low protein diet, *J. Nutrition* 48:477 1952.
186. CARROLL, R. W. HEDGLEY, G. W.; SITTLER, C. L., WILCOX, E. L., AND GRAHAM, W. R.: Absorption of nitrogen and amino acids from soybean meal as affected by heat treatment or supplementation with aureomycin and methionine, *Arch. Biochem.* 45:260 1953.
187. HEDGLEY, G. W., CARROLL, R. W., WILCOX, E. L., AND GRAHAM, W. R. The effects of aureomycin and methionine supplements fed to rats receiving soybean meal, *Arch. Biochem.* 45:270 1953.
188. BERRY, M. E. AND SCHUCK, C.: The effect of aureomycin on growth and protein utilization in the rat, *J. Nutrition* 54:271, 1954.
189. FORBES, R. M. Studies on the influence of antibiotics and methionine on nitrogen utilization and basal metabolism of the growing male albino rat, *J. Nutrition* 53:575, 1954.
190. HARTBROOK, E. W., AND JOHNSON, B. C. Effects of dietary terramycin and methionine supplements on fat and protein gains in weanling rats, *J. Nutrition* 51:205, 1953.
191. MEADE, R. J. AND FORBES, R. M.: The influence of aureomycin and vitamin B₁₂ alone and in combination on nitrogen utilization by growing swine, *J. Animal Sc.* In press, 1953.

19. MEADE, R. J.: The influence of protein content of the diet and of aureomycin and/or vitamin B₁ supplementation upon performance of growing fattening pigs. *J. Animal Sc.* In press, 1955
193. BROWN, P. B., BECKER, D. E., TERRILL, S. W., AND CARD, L. E.: The effect of aureomycin on the growth and metabolism of the pig. *Arch. Biochem.* 41: 378 1952.
194. RUSSO, J. M., HANSON, L. E., AND JACESEK, J. J.: Effects of arsenic acid and aureomycin on nitrogen balance in pigs. *Scientific Journal Series of the Minnesota Agricultural Experiment Station, J. Animal Sc.* In press, 1955
195. GALE, E. F., AND FOLKES, J. P.: The assimilation of amino acids by bacteria. 15. Action of antibiotics on nucleic acids and protein synthesis in *S. aureus*. *Biochem. J.* 53:493, 1953.
196. LAMETER, C. W.: Antibiotics as growth stimulants for dairy cattle. *J. Dairy Sc.* In press, 1955
197. CATRON, D. V.: Antibiotics in swine nutrition, paper presented at the University of Wisconsin, Nutrition Conference, 1953.
198. STOKSTAD, E. L. R. AND JONES, T. H.: Growth-promoting effect of aureomycin on turkey poult, *Poultry Sc.* 29: 811 1950
199. MCGINNIS, J.; BING, L. R.; STERN, J. R.; WILCOX, R. A., AND BEARER, C. E.: The effect of aureomycin and streptomycin on growth of chicks and turkey poult, *Poultry Sc.* 29: 771 1950.
200. MCGINNIS, J., STERN, J. R., WILCOX, R. A., AND CARVER, J. S.: The effect of different antibiotics on growth of turkey poult, *Poultry Sc.* 30: 492, 1951.
201. KRAMER, E. H., AND FRITZ, J. C.: The response of chicks and poult to low levels of antibiotics, *Poultry Sc.* 30: 921, 1951
202. WILLIAMS, O. M., AND HILL, J. E.: The effect of antibiotics on the growth of two varieties of turkey poult, *Poultry Sc.* 31: 769 1952.
203. HEDDER, C. F. AND NURSE, L. C.: Some results of feeding antibiotics to chickens, *Poultry Sc.* 31: 857 1952.
204. LEWIS, R. W. AND SANFORD, P. E.: Use of equivalent levels of antibiotic and vitamin B₁₂ supplements with cottonseed and soybean oil meals in chick diets, *Poultry Sc.* 32:268, 1953.
205. LITTLE, R. J. AND BIRD, H. R.: Evaluation of antibiotic feed supplements and crystal-line antibiotics in chick rations, *Poultry Sc.* 32: 531, 1953.
206. BRANSON, H. D.; ANDERSON, G. W. AND HILL, D. C.: Antibiotics and the growth of ducks, *Poultry Sc.* 32: 335, 1953.
207. RUNNELS, T. D.; REYNOLDS, J. W. AND WALLER, E. F.: The influence of various levels of penicillin and streptomycin on rate of growth and feed efficiency in broiler diets, *Poultry Sc.* 30: 929 1951
208. PEPPER, H. J.; OBERG, E. B.; BENEDICT, R. C., AND LUDWIGFELDER, L. A.: The effect of feeding crude polymyxin D on the intestinal bacteria of chickens, *Poultry Sc.* 29:520, 1950
209. BRANSON, H. D. AND HILL, D. C.: The comparative effect of antibiotics on the growth of poult, *Poultry Sc.* 30:793, 1951
210. STERN, J. R., SIEBENTHAL, J. MCN., AND MCGINNIS, J.: Lack of growth response of turkey poult to certain antibiotics and bacteriostatic agents, *Poultry Sc.* 31: 179 1952.
211. PEPPER, H. T. AND RODGERS, N. E.: Effect of streptothricin on chick growth, *Poultry Sc.* 31:1104 1952.

174. SLINGER, S J., BERKEY J E., PEPPER, W F., SUTHER, E. S., AND ARTHUR, D.: Effect of antibiotics on the protein requirement of broilers, *Poultry Sc.* 31:757 1952.
175. SCOTT H. M.; GOFFI, E. A., AND GLITA, W. A. The protein requirement of the chick as influenced by aureomycin, *Poultry Sc.* 31:751, 1952.
176. SAXENA, H. C. STARR, M. E. BLAYLOCK, L. G. CARVER, J. S., AND MCGOUGH, J.: Effect of dietary penicillin on the efficiency of protein utilization by chicks, *Arch. Biochem.* 44:346 1953.
177. SAXENA, H. C. BLAYLOCK, L. G.; CARVER, J. S., AND MCGOUGH, J.: Factors affecting the growth response of turkey poult to antibiotics, *Poultry Sc.* 32:788, 1953.
178. SLINGER, S J. GARTLEY E. M. PEPPER, W F., AND HILL, D. C.: The influence of animal protein factor supplements and antibiotics on the incidence and severity of white feathers in turkeys, *J. Nutrition* 43:345 1951.
179. BELY J. MARCH, B. E., AND SMITH, D. F. The effect of feeding antibiotics on the carbohydrate and protein requirements of the chick, *Poultry Sc.* 31:803, 1952.
180. MATTHESSON L. D.; SIMONSEN E. P., DECKER, L. AND KOROFF A.: The influence of protein level, aureomycin and vitamin B₁₂ on the growth and feed efficiency of chicks, *Poultry Sc.* 31:940 1952.
181. WHITE-STEVENS R., AND ZIEBEL, H. G.: Chlorotetracycline in relation to protein level of diet on growth efficiency of white rock broilers, *Poultry Sc.* 33:50, 1954.
182. CATRON D. V., JOHNSON A. H. HOMETER, P. G. MADDOCK, H. M., AND ARHTON, G. C. Re-evaluation of protein requirements of growing fattening swine as influenced by feeding an antibiotic, *J. Animal Sc.* 11:221, 1952.
183. JOHNSON A. H.; ACKER, D. C.; MADDOCK, H. M., ARHTON G. C.; HOMETER, P. G., HADY E. O., AND CATRON D. V. Different protein levels with and without antibiotics for growing-finishing swine: effect on growth rate and feed efficiency *J. Animal Sc.* 14:69 1953.
184. BLACK, A., AND BRATZLER, J. W. The effects of a vitamin B₁₂ supplement, vitamin B₁₂ and streptomycin on the metabolism of the rat, *J. Nutrition* 47:159 1952.
185. KROENIG, L. K., AND BLACK, A. The influence of vitamin B₁₂ and antibiotics on protein and energy utilization in a low protein diet, *J. Nutrition* 48:477 1952.
186. CARROLL, R. W.; HENSELEY G. W., SITTIER, C. L.; WILCOX, E. L., AND GRAHAM, W. R. Absorption of nitrogen and amino acids from soybean meal as affected by heat treatment or supplementation with aureomycin and methionine, *Arch. Biochem.* 45:260, 1953.
187. HENSELEY G. W.; CARROLL, R. W. WILCOX, E. L., AND GRAHAM, W. R.: The effects of aureomycin and methionine supplements fed to rats receiving soybean meals, *Arch. Biochem.* 45:270, 1953.
188. BERRY M. E. AND SCHUCK, C. The effect of aureomycin on growth and protein utilization in the rat, *J. Nutrition* 54:271, 1954.
189. FORNER, R. M. Studies on the influence of antibiotics and methionine on nitrogen utilization and basal metabolism of the growing male albino rat, *J. Nutrition* 53:275, 1954.
190. HARTBROOK, E. W., AND JOHNSON B. C. Effects of dietary terramycin and methionine supplements on fat and protein gains in weanling rats, *J. Nutrition* 51:205, 1953.
191. MEADE, R. J., AND FORNER, R. M. The influence of aureomycin and vitamin B₁₂ alone and in combination on nitrogen utilization by growing swine, *J. Animal Sc.* In press, 1955.

182. MEADE, R. J.: The influence of protein content of the diet and of aureomycin and/or vitamin B₁ supplementation upon performance of growing fattening pigs, J. Animal Sc. In press 1953
183. BROWN P. B.; BECKER, D. E., TERRILL, S. W., AND CARD, L. E.: The effect of aureomycin on the growth and metabolism of the pig, Arch. Biochem. 41:378, 1952.
184. RUSSO, J. M.; HANSON, L. E., AND JONES, J. J.: Effects of aspartic acid and aureomycin on nitrogen balance in pigs, Scientific Journal Series of the Minnesota Agricultural Experiment Station, J. Animal Sc. In press, 1953
185. GALE, E. F., AND FOLKES, J. P.: The assimilation of amino acids by bacteria: 15. Action of antibiotics on nucleic acids and protein synthesis in *S. aureus* Biochem. J. 53:493, 1953.
186. LASSITER, C. W. Antibiotics as growth stimulants for dairy cattle, J. Dairy Sc. In press, 1953
187. CATRON D. V. Antibiotics in swine nutrition, paper presented at the University of Wisconsin, Nutrition Conference, 1953.
188. STOKESAD, E. L. R., AND JONES, T. H. Growth-promoting effect of aureomycin on turkey poult, Poultry Sc. 29:611 1950.
189. MCGOWEN, J. BEAG, L. R., STERN J. R. WILCOX, R. A., AND BEARER, G. E. The effect of aureomycin and streptomycin on growth of chicks and turkey poult, Poultry Sc. 29:71 1950.
200. MCGOWEN, J. STERN J. R., WILCOX, R. A. AND CARVER, J. S. The effect of different antibiotics on growth of turkey poult, Poultry Sc. 30:492, 1951.
201. KRAMER, E. H., AND FRITZ, J. C. The response of chicks and poult to low levels of antibiotics, Poultry Sc. 30:921, 1951.
202. WILLIAMS, O. M., AND HILL, J. E.: The effect of antibiotics on the growth of two varieties of turkey poult, Poultry Sc. 31:769 1952.
203. HECHER, G. F., AND NORMAN, L. C. Some results of feeding antibiotics to chickens, Poultry Sc. 31:857 1952.
204. LEWIS, R. W. AND SANFORD, P. E. Use of equivalent levels of antibiotic and vitamin B₁₂ supplements with cottonseed and soybean oil meals in chick diets, Poultry Sc. 32:268, 1953
205. LILLIE, R. J. AND BIRD H. R. Evaluation of antibiotic feed supplements and crystalline antibiotics in chick rations, Poultry Sc. 32:531, 1953.
206. BRANTON H. D. ANDERSON G. W. AND HILL, D. C. Antibiotics and the growth of ducks, Poultry Sc. 32:335, 1953.
207. RUNNELS, T. D. RETHOLDS, J. W., AND WALLER, E. F. The influence of various levels of penicillin and streptomycin on rate of growth and feed efficiency in broiler diets, Poultry Sc. 30:929 1951.
208. PEPPLER, H. J.; OBERG, E. B. BENEDICT R. G. AND LINDENFELDER, L. A.: The effect of feeding crude polymyxin D on the intestinal bacteria of chickens, Poultry Sc. 29:520, 1950.
200. BRANTON H. D. AND HILL, D. C.: The comparative effect of antibiotics on the growth of poult, Poultry Sc. 30:793 1951
210. STERN J. R.; SIEBERTH, J. MCN., AND MCGOWEN, J.: Lack of growth response of turkey poult to certain antibiotics and bacteriostatic agents, Poultry Sc. 31:179 1952.
211. PEPPER, H. T., AND RODGERS, N. E. Effect of streptothricin on chick growth, Poultry Sc. 31:1104, 1952.

212. GERARD, W. E., READ D. C., AND PIERCE, J. M.: Growth stimulation. A comparative evaluation of several antibiotics on chick and swine growth, *J. Agric. Food Chem.* 1:784, 1953.
213. McCLEMMONT G. L., AND DUNCAN D. C.: Aureomycin supplement increases growth of chickens on high and low protein diets, *Agric. Gaz. N. S. W.* 63:461 1952.
214. BING, L. R. BEARSE, G. E.; MCGONNIE, J. AND MILLER, V. L.: The effect of removing supplemental aureomycin from the ration on the subsequent growth of chicks, *Arch. Biochem.* 29 404, 1950.
215. HILL, C. H., AND KELLY J. W.: The effect of fish meal on the response of chicks to high levels of penicillin, *Poultry Sc.* 33:657 1954.
216. BRANTON H. D., AND HILL, D. C.: Fish meal and the response of chicks to antibiotics, *Poultry Sc.* 32 151, 1953.
217. STOKSTAD, E. L. R. JUKES, T. H., AND WILLIAMS, W. L.: The growth-promoting effect of aureomycin on various types of diet, *Poultry Sc.* 32:1054, 1953.
218. HILLERMAN, J. P. KRATZER, F. H., AND WILSON W. O.: Food passage through chickens and turkeys and some regulating factors, *Poultry Sc.* 32:332, 1953.
219. BRANTON H. D., AND HILL, D. C.: Antibiotics and the growth of goslings, *Poultry Sc.* 31:1100 1952.
220. SLINGER, S. J.; SNYDER, E. S. AND PEPPER, W. F.: Effect of penicillin on the growth of goslings, *Poultry Sc.* 32 806, 1953.
221. BALDINI, J. T.; ROBERTS, R. E., AND KIRKPATRICK, C.: Antibiotic and vitamin B₁₂ supplements as related to the crude protein level of bobwhite quail diets, *Poultry Sc.* 32:563 1953.
222. HEUSER, G. F., AND SCOTT M. L.: Bowed legs in ducks, a nutritional disorder *Poultry Sc.* 33:187 1953.
223. MARKHAM, F. S., AND WONG, S. C.: *Plouropneumonia-like* organisms in the etiology of turkey sinusitis and chronic respiratory disease of chickens, *Poultry Sc.* 31:902, 1952.
224. WHITE-STEVENS, R., AND ZIEBEL, H. G.: The effect of chlortetracycline (Aureomycin) on the growth efficiency of broilers in the presence of chronic respiratory disease, *Poultry Sc.* 33 1164, 1954.
225. BRAUDE, R., WALLACE, H. D. AND CUMBA, T. J.: The value of antibiotics in the nutrition of swine - a review *Antib. & Chemot.* 3 271, 1953.
226. GILLESPIE, L., AND WALLACE, H. D.: Ictyocin as a feed ingredient for growing-fattening swine, *J. Animal Sc.* 13 30, 1954.
227. JUKES, T. H., STOKSTAD, E. L. R., TAYLOR, R. R.; CUMBA, T. J., EDWARDS, H. M., AND MEADOWS, G. B.: Growth-promoting effect of aureomycin on pigs, *Arch. Biochem.* 29 452, 1950.
228. CAUFORTER, L. E.: Effect of APF concentrate containing aureomycin on gestating, lactating and growing swine, *J. Animal Sc.* 10 657 1951.
229. WAHLSTROM, R. C.; TERRY, S. W. AND JOHNSON, B. C.: Effect of antibacterial agents on growth of baby pigs fed a "synthetic" diet, *Proc. Soc. Exper. Biol. & Med.* 75 710, 1950.
230. WILLIAMS, W. L.; OWEN, B. D.; ELLIOTT R. F. AND JUKES, T. H.: Aureomycin in baby pig milks and starters, *J. Animal Sc.* 11:781, 1952.
231. BLIGHT J. C.; KING, J. X., AND ELLER, N. R.: Effect of vitamin B₁₂ and aureomycin concentrates on the growth rate of unthrifty weanling pigs, *J. Animal Sc.* 11:92, 1952.

232. CARPENTER, L. E. More big news about aureomycin, *Horned Farmer* Aug., 1950
233. CATRON D. V., AND CUFF P.: Runt pigs can be saved, *Iowa Farm Sc.* 5:119 1951
234. BECKER, R. E.; TIERULL, S. W.; NESHET, R. O. AND MEADE, R. J.: Comparison of antibiotics for growing fattening pigs, Mimeograph AS 215, Univ of Ill. 1951.
235. BURNING, J. E., CUMHA, T. J.; PEARSON A. M., GLASCOCK, R. S., AND SIKELY A. L.: Effect of APF supplement on pigs fed different protein supplements, *Arch. Biochem.* 23:328, 1949
236. ROBINSON, K. L.; COSTY W. E., AND BURNETT G. S.: Some effects of procaine penicillin and of certain vitamin supplements in food consumption and liveweight gain of pigs, *J. Sc. Food Agric.* 3:445 1952.
237. BECKER, D. E.; TIERULL, S. W., AND NOTZOLD, R. A.: Supplementary protein and the response of the pig to antibiotics, *J. Animal Sc.* 14:492, 1955
238. LEFLEY K. C.; CATRON D. V., AND CULBERTSON C. C.: Dried whole aureomycin mash and meat and bone scraps for growing-fattening swine, *J. Animal Sc.* 9:608, 1950.
239. LUECKE, R. W., NEWLAND, H. W.; McMILLEN W. N., AND THORP F., JR.: The effect of antibiotics fed at low levels on the growth of weanling pigs, *J. Animal Sc.* 9:662, 1950.
240. BOWLAND, J. P.; BEACON, S. E., AND McELROY L. W. Animal protein factor and antibiotic supplementation of small grain rations for swine *J. Animal Sc.* 10:629 1951.
241. BRUGG, J. E.; ELBOD, R., AND BEESON W. M.: The effect of type of ration on the penicillin requirement for growing fattening pigs, *J. Animal Sc.* 10:1041, 1951.
242. BRIDGEMAN, M. F. AND WARDER, D. R.: Antibiotic studies with swine in dry lot and on alfalfa pasture, *J. Animal Sc.* 10:1041 1951.
243. HOGAN, A. G. LARLEY J. F. AND TRIBBLE, L. F. The value of procaine penicillin in rations for brood sows, *Prog. Report 16*, Univ of Missouri, 1951.
244. NOLAND, P. R. TUCKER, D. L., AND STEPHENSON E. L. Subcutaneous implantation of bacitracin in pellet form to stimulate growth of suckling pigs, *Arkansas Agric. Exper. Sta. Report Series 34* 1952.
245. PERRY T. W.; TERRASER, G. W., AND BEESON W. M.: The effects of bacitracin implants on weaning weights and subsequent feed lot performance of two-day old pigs, *J. Animal Sc.* 12:824, 1953.
246. SWENSON, M. J.; BUCHNER, R. G., GORTSCH, D. D.; ADRIEL, C. E., AND UNDERBERG, G. K. L. Growth of newborn pigs implanted subcutaneously with bacitracin pellets, *J. Animal Sc.* 13:78, 1954.
247. WALLACE, H. D., McKINNEY J. AND GILBERT, L. The influence of subcutaneous implantation of bacitracin and chlortetracycline (aureomycin) pellets on the growth and survival of suckling pigs, *Antib. & Chemo.* 4:663, 1954.
248. CLAWSON A. J.; SHEFFY B. E., AND WILLMAN J. P.: The value of implanted antibiotic pellets for suckling pigs, *J. Animal Sc.* 12:911, 1953.
249. TIERULL, S. W.; NESHET, R. O. AND KRAIER, J. L.: Comparison of solvent soybean oil meals and the effect of dried whey products and antibiotics in dry lot rations for pigs, *J. Animal Sc.* 10:828, 1951
250. HENDRICH A. A.; ROSS O. B.; WHITEHAIR, C. K., AND McVICAR, R. W.: The effect of various levels of penicillin upon growth of pigs, *Okla. Agric. Exper. Sta. Misc. Pub. No. MP 22*, 72, 1951.
251. WOODEN A. N. AND BUNYAN, J.: The administration of a procaine penicillin feeding supplement to runt pigs, *Vet. Rec* 64:647 1952.

252. ROBINSON W L.; KUNKLE, L. E., AND CAHILL, V. R.: The influence of various factors on the yields of pork cuts, *J. Animal Sc.* 11:752, 1952.
253. JENSEN A. H.; ARNTON, G. C.; MADDOCK, H. M. HOMERY, P. G., AND CATRON, D. V.: Effects of antibiotic and lysine supplementation on different protein levels for growing-fattening swine, *J. Animal Sc.* 11:767 1952.
254. KLINE, E. A. KASTELIC, J. QUINN L. Y., HOMERY, P. G., AND CATRON, D. V. The influence of feeding vitamin B₁₂, cobalt and antibiotics on the composition of pork tissue, *J. Animal Sc.* 11 750 1953.
255. WILSON G. D. BURKINS, J. E. BRAY R. W.; PHILLIPS, P. H., AND GRUMMER, R. H.: Pork carcass value as affected by protein level and supplementation with aureomycin and vitamin B₁₂ *J. Animal Sc.* 12 291 1953.
256. WALLACE, H. D. MELICEVIC, M., FRANKSON A. M. CUMHA, T. J. AND KOEHL, M.: The influence of aureomycin on the protein requirement and carcass characteristics of swine, *J. Animal Sc.* 13:177 1954.
257. BOHMAN V. R. HUNTER, J. E., AND MCCORMICK, J. The effect of graded levels of alfalfa and aureomycin upon growing fattening swine, *J. Animal Sc.* 14:499 1953.
258. GARDNER, K. E. NEVENS, W. B. FOLKERTS, T. AND JOHNSON, B. C. Growth stimulants and *ad libitum* feeding in the production of milk fed veal, *J. Animal Sc.* 11 :61, 1952.
259. SAKENA, H. C.; BLAYLOCK, L. G.; STADELMAN W. J. CARVER, J. S., AND MCGROCK, J.: Effect of penicillin on growth, feed efficiency and fattening of turkeys, *Poultry Sc.* 32:721, 1953.
260. SWICKARD, M. T.; HARKIN A. M. AND MARSDEN S. J. Palatability of turkeys fed experimental diets containing aureomycin and fish products, *Poultry Sc.* 33:726, 1953.
261. TAYLOR, F. R., JR., PATRICK, H., AND GOFF O. E.: Influence of arsenic and penicillin on dressing grades and quality of poultry *Poultry Sc.* 33:1085 1954.
262. VITAL, C. M. Effect of an APF-aureomycin supplement in swine rations, *Purdue Agric. Exper. Sta. Mimeograph* AH 65, 1951.
263. PERRY T. W. BEESON W. M., AND VOSTEREN B. W.: The effect of an antibiotic or a surfactant on the growth and carcass composition of swine, *J. Animal Sc.* 12 310, 1953.
264. BLOOM, S. AND KNOUT C. B.: The value of vitamin B₁₂, DL-methionine, penicillin and aureomycin in milk replacement formulae for dairy calves, *J. Animal Sc.* 10:1039 1951
265. RUBOFF L. L., DAVID, A. V. AND ALFORD, J. A.: Growth-promoting effect of aureomycin on young calves weaned from milk at an early age, *J. Nutrition* 45:289, 1951.
266. CLARK, H. E., HARRISON, D. L.; SOULE, R. P., JR., AND RICHARDSON D.: The nutritive value of proteins of muscle from hogs fed diets supplemented with aureomycin or terramycin hydrochloride, *J. Nutrition* 56 61 1953
267. COLBY R. W. RAU F. A. AND COUCH, J. R.: Effect of feeding an animal protein factor concentrate to young lambs, *Am. J. Physiol.* 163 418, 1950.
268. COLBY R. W. RAU F. A., AND DUNN, R. C.: Effect of feeding aureomycin to fattening lambs, *Proc. Soc. Exper. Biol. & Med.* 75 234, 1950.
- 69 COLBY R. W., RAU, F. A., AND MILLER, J. C. The effect of various antibiotics on fattening lambs, *J. Animal Sc.* 9 652, 1950
270. BELL, M. C.; WHITEHAIN, C. K., AND GALLUP W. D. The effect of aureomycin on digestion in steers, *Proc. Soc. Exper. Biol. & Med.* 76:284, 1951.

271. ANDRE, F: Modification of a standard feed mixture for fattening lambs. II. Iowa State College, Animal Husbandry Leaflet No. 175 February 1950
272. LOOSLI, J K., AND WALLACE, H D: Influence of APF and aureomycin on the growth of dairy calves, *Proc. Soc. Exper Biol. & Med.* 75:531 1950.
273. RUBOFF L. L.: A.P.F. supplements for calves, *J Animal Sc.* 9:906 1950
274. BARTLEY E. E. FOUNTAINE, F C., AND ATKESON F W: The effects of an APF concentrate containing aureomycin on the growth and well-being of young dairy calves, *J Animal Sc.* 9:646, 1950
275. JORDAN R. M. AND BELL, T D: Effect of aureomycin on growing and fattening lambs, *J Animal Sc.* 10 1051 1951
276. KNOTT C. B: Antibiotics in the growth of ruminant animals, *Antib. & Chemo.* 3 442, 1953.
277. BARTLEY E. E.: Antibiotics in calf nutrition, paper presented at Gordon Research Conference, Colby Jr College, N H 49 1951
278. BARTLEY E. E. ATKESON F W.; FRYER, H. C., AND FOUNTAINE, F C.: Antibiotics in dairy cattle nutrition. III. Effects of different levels of aureomycin intake upon the growth and well-being of dairy calves and the association of differences with changes in environment, *J Dairy Sc.* 37:259 1954.
279. BARTLEY E. E. FOUNTAINE, F C. ATKESON F W. AND FRYER, H. C.: Antibiotics in dairy cattle nutrition. IV. Comparative effects of different levels of crystalline aureomycin and a crude aureomycin fermentation product (Aurofao 2A) on the growth and well-being of young dairy calves. *J Dairy Sc.* 37 962, 1954.
280. RUBOFF L. L.: Antibiotic feed supplement (aureomycin) for dairy calves, *J Dairy Sc.* 34 652, 1951
281. RUBOFF L. L.; FUSSELL, J M. HYDE, C. E., AND CROWN R. M. Oral supplement versus intramuscular injection of aureomycin to young calves, *Annual Report 1952-53 of Louisiana Agric. Experiment Station, Research in Agriculture*, p. 59
282. RUBOFF L. L., HAQ, M O. BRANTON C. PATRICK, T E. AND DARENBOROUGH, G: Report on feeding aureomycin supplement to mature dairy bulls and lactating dairy cows, *J Animal Sc.* 11 774, 1952.
283. RUBOFF L. L. FUSSELL, J M. HYDE, C. E., AND CROWN, R. M.: Oral supplementation versus intramuscular injection of aureomycin to young calves, *J Dairy Sc.* 36:593, 1953.
284. RUBOFF L. L., AND HAQ, M. O: Studies on aureomycin and vitamin B₁₂ supplementations for dairy cows. II. Effect on production, composition and vitamin B₁₂ content of the milk, *J Dairy Sc.* 37 677 1954.
285. RUBOFF L. L. LANDAGORA, R. T. AND HESTER, H. II.: Effect of aureomycin on certain blood constituents (Hb packed RBC, Ca, P RBC and WBC) body temperatures weights of organs and tissues, and thickness of small intestine *J Dairy Sc.* 37 654 1954
286. RICHARDSON, C. W.; RONNING, M. BERGQUIST, E. R., AND NORTON C. L.: The effect of aureomycin upon the growth of dairy calves when administered orally subcutaneously and intramuscularly *J Dairy Sc.* 36 593, 1953.
287. ROY J H B; SHILLAM K. W G., PALMER, J., AND INGRAM, P L. The nutritive value of colostrum for the calf *Brit J Nutrition* 9:94 1955.
288. HAQ, M. O.; RUBOFF L. L., AND CELPI A. J JR.: Antibiotic feed and vitamin B₁₂ supplements for lactating dairy cows, *Science* 115 15 1952.

289. BARTLEY E. E., FOUNTAINE, F. E., AND AYERSON, F. W.: Antibiotics in dairy cattle nutrition. II. Effects of feeding an aureomycin product (Aurofac 2A) to lactating dairy cows, *J Dairy Sci.* 36:402, 1953.
290. WARNER, R. G.: Recent findings on antibiotics and surfactants for ruminants, *Proc. 1952, Cornell Nutrition Conf. for Feed Manufacturers*, Nov. 6-7 1952.
291. BELL, M. C.; WHITEHAIR, C. K., AND GALLUP W. D.: The effect of aureomycin on digestion in steers, *J Animal Sc.* 9:647 1950.
292. HORNE, L. H., JR.; SNAFF R. R., AND GALL, L. S.: The effects of antibiotics upon the digestion of feed nutrients by yearling steers, with bacteriological data, *J Animal Sc.* 14:243, 1953.
293. TOVERA, E.: Personal communication, 1954.
294. MATSUHIMA, J.; DOWE, T. W., AND ADAMS, C. H.: Effect of aureomycin in preventing liver abscess in cattle, *Proc. Soc. Exper. Biol. & Med.* 85:18, 1954.
295. PERRY T. W., BEESON W. M.; HORNBACK, E. C., AND MOHLER, M. T. Aureomycin for growing and fattening beef animals, *J Animal Sc.* 13:8, 1954.
296. DUTTMAN W. W., AND KEMLER, F. B.: Beef cattle feeding and breeding investigations for 1952-53, *KANSAS Agric. Exper. Sta. Cir. No. 298*, April 24, 1953.
297. DUTTMAN W. W., AND KEMLER, F. B. Beef cattle feeding investigations: 1953-54, *KANSAS Agric. Exper. Sta. Cir. No. 307* April 28, 1954.
298. TURNER, A. W., AND HODGKINS, V. E. Depression of ruminal digestion in adult sheep by aureomycin, *Austral. J. Agric. Res.* 3:453, 1952.
299. JORDAN R. M.: Aureomycin supplements in lamb fattening rations, *J Animal Sc.* 11:566, 1952.
300. JORDAN R. M., AND BELL, T. D.: Effect of aureomycin supplements on suckling lambs, *J Animal Sc.* 13:451, 1954.
301. JORDAN R. M. Lamb responses to antibiotics and pelleted rations, *J Animal Sc.* 13:53 1954.
302. HATFIELD E. E., AND GARROUS, U. S.: Aureomycin in a creep ration, *Illinois Agric. Exper. Sta. Bull. No. A.S. 296b*, 1953.
303. HATFIELD, E. E., AND GARROUS, U. S. Antibiotics in lamb rations, *Illinois Agric. Exper. Sta. Bull. No. A.S. 352b*, 1953.
304. HATFIELD, E. E., AND GARROUS, U. S.: Antibiotics in rations for lambs, *J Animal Sc.* 13:919, 1953.
305. HATFIELD, E. E., GARROUS, U. S., AND NORTON, H. W.: Antibiotic supplements in rations for growing and fattening lambs, *J Animal Sc.* 13:715 1954.
306. BRIDGES, J. H., MILLER, J. C., KAMMLADE, W. G., JR., AND KUNKEL, H. O.: Effect of various levels of aureomycin in fattening lambs, *J Animal Sc.* 12:600, 1953.
307. ELLIOTT R. F., AND ELLSWORTH, S. A.: The effect of aureomycin and type of ration on fattening lambs, *J Animal Sc.* 12:914, 1953.
308. BOTKINS, M. P., AND PAULEY, L.: Effect of aureomycin in various ratios of roughage to concentrate for feeder lambs, *Univ. of Wyo. Agric. Exper. Sta. Mimeo. Cir. No. 44*, 1954.
309. BOTKINS, M. P., AND PAULEY, L.: Control of death loss and digestive disturbances in feeder lambs, *Univ. of Wyo. Agric. Exper. Sta. Mimeo. Cir. No. 33*, 1953.
310. KUNKEL, H. O.: Unpublished data, *A. H. Dept. Texas Agric. Exper. Sta.*, 1954.

811. KUNKEL, H. O., AND PACKETT L. O., JR.: Unpublished data, A. H. Dept. Texas Agric. Exper. Sta., 1953.
812. KIDMAN D. M., AND RIDGELL, W. H.: Antibiotics in creep ration of suckling lambs J. Animal Sc. 11:769 1952.
813. BORDMAN V. R.; HUNTER, J. E., AND WALKER, L. Antibiotics and B-vitamins on lambs, J. Animal Sc. 14:1111 1953.
814. CARPENTER, L. E., AND LARSON N. Antibiotics and the reproduction of swine, J. Animal Sc. 12 821 1953.
815. VESTAL, C. M.; BEESON, W. M. ANDREWS, F. N.; HUTCHINGS, L. M. AND DOYLE, L. P. The effect of aureomycin on the development and livability of newborn pigs Purdue University Mimeograph, AH-87 1952.
816. DYER, I. A.: The gestation-lactation response of gilts fed a 17% crude protein soybean oil meal ration to alfalfa meal, dried corn distillers' solubles and supplement containing vitamin B₁₂ and antibiotics, University of Illinois, Dept. Animal Sc. Mimeograph, 9th Annual Report to Collaborators, 41 1952.
817. BEESON, W. M.; VESTAL, C. M.; ANDREWS, F. N.; HUTCHINGS, L. M. AND DOYLE, L. P. The effect of different levels of aureomycin on the birth weight and livability of newborn pigs, Purdue University Mimeo., AH 114, 1953.
818. DAVY R. J. GREEN W. W., AND STEVENSON J. W.: The effect of aureomycin on growth and reproduction of swine, J. Animal Sc. 14 507 1953.
819. MADDOCK, H. M., CATTON D. V., SPEER, V. C.; CUFF P. W. W., AND ASHTON G. C.: Concentration of chlortetracycline in sows' blood and milk after oral administration, Antib. & Chemo. 3:1075 1953.
820. WONG, S. C. KERSEY R. C. McMAHAN J. R. REYNOLDS, W. M., WARDEN W. K., AND MCGONNIS, J.: Effects of feeding terramycin and penicillin to laying hens on antibiotic deposition in eggs and growth of murine typhus rickettsiae in embryonated eggs, Federation Proc. 12:434 1953.
821. MANIAKULANDIA, A.; MYINT T. AND MCGONNIS J. Effect of terramycin and vitamin B₁₂ on hatchability Proc. Soc. Exper. Biol. & Med. 79 242, 1952.
822. SIZEMORE, J. R., LILLIE, R. J., DENTON C. A., AND BIRD, H. R.: Influence of aureomycin in the chick diet upon subsequent reproductive performance of laying hens, Poultry Sc. 32 618, 1953.
823. SLINGER, S. J.; FERGUSON A. E., AND McCOMACHIE, J. D.: A carry-over effect of penicillin from dam to progeny Poultry Sc. 31 172, 1952.
824. BENTLEY O. G. AND HERSHBERGER, T. V.: The effect of antibiotics on hatchability of hens' eggs and progeny growth performance, Poultry Sc. 33 641, 1954.
825. BENTLEY O. G.: Rat growth response of egg yolks from hens fed antibiotics, Federation Proc. 13 181 1954.
826. McFARLANE, W. D.; FULMER, H. L., AND JUKES, T. H.: Studies in embryonic mortality in the chick. I. The effect of diet upon the nitrogen, amino-nitrogen, tyrosine, tryptophan, cystine and iron content of the proteins and on the total copper of the hen's egg, Biochem. J. 24:1611, 1930.
827. SUNDH, M. L.; HALPERN, J. G., AND CRAVENS, W. W. The effect of vitamin B₁₂ supplements and antibiotic feed supplements on egg production and hatchability Poultry Sc. 31:617 1952.
828. BEAC, L. R. CARVER, J. S., BEARSE, C. E., AND MCGONNIS, J.: Antibiotics in the nutrition of laying hens, Washington Agric. Exper. Sta. Bull. No. 534, 10, 1952.

- 329 BROWN J. A., ROSELEY, A. R., AND CLANDININ D. R.: The use of penicillin in breeding rations, *Poultry Sc.* 32:576, 1953.
- 330 PETERSEN, C. F., AND LAMPMAN C. E.: Value of antibiotics in rations for egg production, *Poultry Sc.* 31:1067 1952.
331. HALSBROOK, E. R., BRECKLER, A. F. AND SMITH, E. P.: The value of fish scum and a vitamin B₁₂-antibiotic supplement in chicken and turkey mash mixtures containing ten percent meat scraps, *Poultry Sc.* 32:1058, 1953.
332. SHERWOOD, D. H. AND MILBY T. T.: Further tests with antibiotics for laying and breeding hens, *Poultry Sc.* 33:1031, 1954.
333. CARPENTER, K. J., AND DUCKWORTH, J.: Egg production of pullets fed on a ration containing aureomycin, *Nature* 169:669 1952.
334. JOHNSON E. L.: Vitamin B₁₂ requirements of hens as affected by choline and penicillin, *Poultry Sc.* 33:100, 1954.
- 335 ELAM, J. F., JACOBS, R. L., AND COUGH, J. R.: The effect of prolonged feeding of antibiotics upon the performance of laying hens, *Poultry Sc.* 32:782, 1953.
336. HETWANG, B. W.: High levels of antibiotics in the diets of laying chickens, *Poultry Sc.* 33:22, 1954.
- 337 BALLOU S. L.: Effect of high-level aureomycin feeding on rate of egg production, *Research Notes, Poultry Sc.* 33:867 1954.
338. HETWANG B. W.: High levels of antibiotics in the diets of chickens during hot weather Informal Poultry Nutr. Conf. Federation Meetings, San Francisco, 1955.
- 339 WHITE-STEVEN, R., AND ZIEGLER, H. G.: Unpublished experiments.
340. CARLSON C. W., AND KOHLMAYER, W.: Factors affecting the value of antibiotics in breeder diets, *Poultry Sc.* 33:1047 1954.
341. CARLSON C. W.; JONES, D. G.; KOHLMAYER, W. AND MOXON A. L.: The effects of vitamin B₁₂ and aureomycin on reproductive performance in turkeys, *Poultry Sc.* 32:884, 1953.
342. WHITE-STEVEN, R., ZIEGLER, H. G. AND SMITH, F.: In preparation.
- 343 CARSON J. R.; EATON, R. D., AND LUGENBUHL, R. E.: The effect of injections of an oil suspension of tetracycline on egg production and egg quality in hens affected with chronic respiratory disease, *Poultry Sc.* 33:589 1954.
344. ERSBOW B. H.: Prolonged survival of hyperthyroid rats fed penicillin and aureomycin residues, *Arch. Biochem.* 28:359, 1950.
- 345 ERSBOW B. H.: Distribution of an unidentified antithyrototoxic factor in materials of plant and animal origin, *Proc. Soc. Exper. Biol. & Med.* 74:391 1950.
- 346 MEYER, J., AND OGLE, R. C.: Antithyrototoxic effects of antibiotics in rats, *Proc. Soc. Exper. Biol. & Med.* 77:758, 1951.
- 347 CALEROCK, B.; HARRIS, W. D., AND JONES, R. S.: Antithyroid action of antibiotics, *Science* 119:128, 1954.
- 348 LIBBY D. A., AND MEYER, J.: Negative effects of antibiotics on thyroid gland, *Science* 120:354, 1954.
- 349 GRANT W. C.: Aureomycin and the thyroid gland, *Science* 120:724, 1954.
350. TAUNO, A., AND CHAIKOFF I. L.: Personal communication.
351. TELKKA, A., AND KUUSISTO, A. N.: Is the growth-promoting effect of antibiotics connected with thyroid function? *Acta Endocrinol.* 18:863 1954.

352. MAGNABRI, M. G. AND TURNER, C. W.: Nutritional requirements in hyperthyroidism of growing chicks, Univ. of Missouri Agric. Exper. Sta. Bull. 523 1953.
353. MCELLEN W. J. AND WALLER, E. F.: Antibiotics and thyroid size in growing chickens Poultry Sc. 33:1036 1954
354. GRANT W. C.: Unpublished data, 1955
355. MEYER, J.: Counteraction of cortisone inhibition of body hair and thymus growth by vitamin B₁₂ and aureomycin, Proc. Soc. Exper. Biol. & Med. 78:692, 1951
356. MEYER, J.: Beneficial effects of vitamin B₁₂ and aureomycin in rats given large doses of cortisone, Proc. Soc. Exper. Biol. & Med. 81:307 1952.
357. COHEN A. M., AND RACHIMILEWITZ, M.: Effect of aureomycin in rats with chronic alloxan diabetes, Proc. Soc. Exper. Biol. & Med. 83:50 1953.
358. BARKER, R. S.; BRAUER, R., AND MITCHELL, K. G.: Antibiotics and endocrine stimulants as promoters of growth in fattening pigs, Chem. & Indust. 17 410, 1953.
359. BAEZ, A. SRIKANTIA, S. G. FOWDER, I., AND SIKOR, E.: Metabolic action of aureomycin on liver: inhibition of anaerobic release of ferritin (VDM) Federation Proc. 14 6, 1955.
360. FINCH, J.: Effect of antibiotics on irreversibility to transfusion in hemorrhagic shock, Harvard M. Alumni Bull. 29 105 1952.
361. GUGGENHEIM, K., AND HALEVY S.: The effects of vitamin B₁₂ organ extracts, yeast and antibiotics on emetine toxicity in rats, J. Nutrition 53 129 1954
362. HOWLAND J. W. FURTH, F.; BENNETT L. R. COULTER, M., AND McDONNELL, G. M.: Studies on factors affecting the radiation syndrome. I. The effect of aureomycin and antibiotics on whole body irradiations, Report UR94, Univ. of Rochester Atomic Energy Project, 1949
363. MILLER, C. P. HALMOND, C. W., TOMPKINS, M. AND SHORTER, G.: The treatment of postirradiation infection with antibiotics: an experimental study on mice, J. Lab. & Clin. Med. 39 462, 1952.
364. GUSTAFSON G. E., AND KOLETEKY S.: Effect of oral tetracycline prior to whole body X-radiation, Proc. Soc. Exper. Biol. & Med. 78 489 1951.
365. FURTH, F. W., AND COULTER, M. P.: Studies on factors affecting the radiation syndrome, Report UR 116, Univ. of Rochester Atomic Energy Project, 1950.
366. FURTH, F. W.; COULTER, M. P. AND HOWLAND, J. W.: The effect of aureomycin and tetracycline on the X radiated rat, Am. J. Path. 28 185 1952.
367. FURTH, F. W. COULTER, M. P.; MILLER, R. W. HOWLAND, J. W. AND SWENNER, S. N.: The treatment of acute radiation syndrome in dogs with aureomycin and whole blood, J. Lab. & Clin. Med. 41:918, 1953.
368. KIELICK, R. L., AND SAKAMI, W.: A study of the mechanism of serine biosynthesis, J. Biol. Chem. 214:47 1955
369. ALEXANDER, N. AND GREENBERG, D. M.: Studies in the biosynthesis of serine, J. Biol. Chem. 214 821 1955
370. BEST, C. H.; LUCAS, C. C., AND RIMOUT, J. H.: The lipotropic factors, Ann. New York Acad. Sc. 57 646 1954
371. ABELL, M. R., AND BEVERIDGE, J. M. R.: Studies on hepatic necrosis induced by dietary means: IV. Conditions affecting the production and prevention of massive liver necrosis. A.M.A. Arch. Path. 52 428 1951
372. GORMY P.; STOKES, J. JR.; GOLDBLATT H., AND POPPER, H.: Antimicrobial agents

in the prevention of dietary hepatic injury (necrosis, cirrhosis) in rats, *J. Exper. Med.* 93:513, 1951.

373. GYORGY P.; STOKES, J., JR., AND GOLDBLATT H.: Antimicrobial agents in prevention of experimental dietary injury of liver. *Trans. A. Am. Physicians* 64:289, 1951.
374. FORBES, M.; SROCKMAN, G. D.; CHU E., AND GYORGY P.: Effect of chlortetracycline and penicillin on the intestinal flora of rats on necrogenic diet, *Antib. & Chemo.* 3: 1104, 1953.
375. LUCKEY T. D. REYNOLDS, J. A.; GYORGY P. AND FORBES, M.: Germ-free animals and liver necrosis, *Ann. New York Acad. Sc.* 57:932, 1954.
376. FORBES, M.; BARNES, L. A., MORLEY, H., AND GYORGY P.: Excretion of ether-soluble acids by rats on necrogenic diet with and without supplements of antibiotics, *Proc. Soc. Exper. Biol. & Med.* 84:162, 1953.
377. ROGERS, W. F., BURDICK, M. P., AND BURDITT G. R.: The effect of antibiotics on the excretion of phenolic compounds, *J. Lab. & Clin. Med.* 45:87, 1955.
378. SHAFER, J. M., FARQUHAR, J. D.; STOKES, J. JR., AND SROGOV V. M.: Studies on the use of aureomycin in hepatic disease. I. Aureomycin therapy in acute viral hepatitis, *Am. J. M. Sc.* 220:1, 1950.
379. MCNEIL, H. J., AND SOLOMON C.: Aureomycin in acute fulminant hepatitis: Report on successful treatment in three cases, *New York State J. Med.* 50:1393, 1950.
380. FARQUHAR, J. D., STOKES, J. JR.; WHITLOCK, C. M., BLUEMILK, L. W., JR., AND CAMERON, J. M.: Studies on the use of aureomycin in hepatic disease. III. A note on aureomycin therapy in hepatic coma, *Am. J. M. Sc.* 220:166, 1950.
381. RUMBLE, J. M.; MARION D. F. AND SANDERS, M.: Post hepatitis cirrhosis treated with aureomycin. A case report, *Gastroenterology* 14:432, 1950.
382. SHAFER, J. M., BLUEMILK, L. W. JR., SROGOV V. M., AND NEEFE, J. R.: Studies on the use of aureomycin in hepatic disease. IV. Aureomycin therapy in chronic liver disease, *Am. J. M. Sc.* 220:178, 1950.
383. DE LA HUEGA, J., GYORGY P., WALDSTEIN, S., KATZ, R., AND POPPER, H.: The effects of antimicrobial agents upon choline degradation in the intestinal tract, *J. Clin. Investigation* 32:1117, 1953.
384. KAPLAN, A.; COHEN, C.; NARAHARA, A., AND JONES, K.: Experimental differentiation between liver necrosis and liver cirrhosis and some dietary factors affecting their development, *Ann. New York Acad. Sc.* 57:688, 1954.
385. BARTER, J. H., AND CAMPBELL, H.: Effects of aureomycin on renal lesions, liver lipid and tissue choline in choline deficiency. *Proc. Soc. Exper. Biol. & Med.* 80:415, 1952.
386. CORMATZER, W. E., AND GALLO, D. G.: Effects of dietary aureomycin and sulfamizidine on phosphorylation in the liver of rats. *Proc. Soc. Exper. Biol. & Med.* 86:15, 1954.
387. GOLDBLOOM, R. S., AND STEIGMANN, F.: Aureomycin therapy in hepatic insufficiency. *Gastroenterology* 18:93, 1951.
388. SETO, J. T., AND LEPPER, M. H.: The effect of chlortetracycline, oxytetracycline, and tetracycline administered intravenously on hepatic fat content; a quantitative method of study including failure of some vitamins and other drugs to mitigate the effect, *Antib. & Chemo.* 4:666, 1954.
389. ZIMMERMAN, H. H., AND HUMMELER, F. L.: Effect of aureomycin on dehydrogenase systems of the rat liver. *Federation Proc.* 12:162, 1953.
390. SROGOV V. M., AND SUTHERLAND, D. A.: Fatty liver following aureomycin and terramycin therapy in chronic hepatic disease, *Gastroenterology* 18:598, 1951.

391. FALOOK, W. W., NOLL, J. W., AND PRION, J. T.: Nitrogen metabolism and liver histology during aureomycin administration in patients with hepatic disease, *J. Lab. & Clin. Med.* 41:590, 1953.
392. SUTHERLAND D. A., MANN J. D., GRIN, B. AND SELIGSON D.: Effect of aureomycin on liver fat and liver function, *Proc. Soc. Exper. Biol. & Med.* 77 458, 1951
393. JUCKER, T. H.: Antimetabolites and antibiotics as tools for research on blood formation, *Am. J. Clin. Nutrition* 3:56 1955.
394. LACITMAN H., GINSBERG, V., AND WATSON J.: Therapeutic effect of aureomycin in pernicious anemia, *Proc. Soc. Exper. Biol. & Med.* 74:884, 1950.
395. FOT H., KORDA, A., AND HARGREAVES, A.: Response of megaloblastic anemia of pregnancy to crystalline penicillin G, *Brit. M. J.* 1:1108 1951
396. FOT H., AND KORDA, A.: Treatment of megaloblastic anemias. Relation of penicillin to vitamin B₁₂ *Lancet* 265:1280, 1953.
397. FOT H., KORDA, A., AND HARGREAVES, A.: The response of megaloblastic macrocytic anemia to crystalline penicillin G. Preliminary report, *Brit. M. J.* 1 391, 1951.
398. DEXTRUMER, P. L.: Crystalline penicillin in macrocytic anaemia, *Antiseptic* 50:825 1953.
399. TRIBE, D. E., BOED, J. M., AND OSBONE, A. D.: Response of cobalt-deficient sheep to oral administration of penicillin, *Nature* 173:728, 1954.
400. WATSON G. M., AND WITTE, L. J.: Intestinal macrocytic anemia, *Brit. M. J.* 1:13, 1952.
401. URGLEY C. C. IV: Administration into buccal cavity into washed segment of intestine, or after partial sterilization of bowel, *Brit. M. J.* 2 915, 1950.
402. OLSSON, N. AND OLSSON, N. E.: Experiments with APF preparation or aureomycin in the feeding of growing and fattening pigs, *Kgl. Lantbrukshogsk. Stat. Hushjarnforsk. Särtryck*, No. 84, 20, 1951
403. BURENINE, J. E., GRUMMER, R. H., PHILLIPS, P. H. AND BOHNETT G.: The influence of crystalline aureomycin and vitamin B₁₂ on the protein utilization of growing-fattening swine, *J. Animal Sc.* 13:184, 1954.
404. MOCHIE, R. D., DOLZ, K. L., EATON H. D., AND ELLIOTT F. I.: Effect of adding various levels of an aureomycin supplement to starter on growth and blood constituents of Guernsey calves, *Storrs Agric. Exper. Sta. INF* 48 3, 1953.
405. VOELCKER, H., AND JACOBSON N. L.: Effects of various antibiotics and a detergent and of frequency of milk replacement feeding on young dairy calves, *J. Dairy Sc.* 36 592, 1953.
406. OWEN, F. G., AND ALLEN, R. S.: Growth and hematological response of calves to antibiotic and arsenical supplementation, *J. Dairy Sc.* 37:654 1954.
407. ACURIE, F. AND SCRIMSHAW N. S.: The effect of supplements of animal and vegetable protein, vitamin B₁₂ and aureomycin on hematological values in Central American school children, *Am. J. Clin. Nutrition* 3:225 1955.
408. MOLLER, H. AND GRAEBLE, O. E.: Pharmacology and toxicology of antibiotics, *J. Pharmacol. & Exper. Therap.* 93:1 1950.
409. CAMERON H. S.: The bactericidal action *in vivo* of aureomycin in an aureomycin fermentation residue against *Brucella suis*, *Cornell Vet.* 41 110 1951
410. FREUDT, F.: Aureomycin as a growth factor in premature infants, *Boll. Soc. Ital. biol. sper.* 27:1151 1951
411. ROBINSON P.: Controlled trial of aureomycin in premature twins and triplets, *Lancet* 1:53, 1952.

412. SNELLING, C. E., AND JOHNSON R.: The value of aureomycin in prevention of cross infection in the hospital for sick children, *Canad. M. A. J.* 66:6, 1952.
413. COOMEN F. J.: Studies of tetracycline in premature infants, *Pediatrics* 12 652, 1953.
414. KAYE, R. CAUGHEY R. H., AND MCCOY W. W.: Effects of vitamin B₁₂ and aureomycin on nitrogen retention in infants, *Pediatrics* 13:462, 1954.
415. DÉCHÈNE, E., Influence des antibiotiques sur la croissance, *Laval Med.* 13:924, 1953.
416. WITZEL, N. D., AND HORWOOD, H. H. Antibiotic supplements overcome growth failure in school children, *J. Agric. & Food Chem.* 2 1148, 1954.
417. SCRIMSHAW N. S.; GUDMAN M. A., AND TANDON O. B. Effect of aureomycin and penicillin on growth of Guatemalan school children, *Federation Proc.* 13:477 1954.
418. CARTER, C. H. Personal communication, 1953.
419. MACDOUGALL, L.: Personal communication, 1953.
420. HAIGHT T. H., AND PIERCE, W. E. Influence of small doses of antibiotics on the weight behavior of young males, *J. Lab. & Clin. Med.* 44 807 1954.
421. GARUZZA, G. J., JACKSON G. G., AND GREGG M. E.: Effect of aureomycin upon nitrogen balance and urinary riboflavin excretion in undernourished men, *J. Clin. Investigation* 31:631 1952.
422. TAYLOR, J. H.; GORDON W. S., AND BURRELL, P.: The effect of supplementing the diet of thoroughbred foals with aureomycin hydrochloride, *Vet. Rec.* 66:744, 1954.
423. MURPHY M. R. V. AND SREENIVARAYA, M.: Effect of antibiotics on the growth of the silkworm, *Bombyx mori*, *Nature* 172:684, 1953.
424. LAWRENCE, J. M., AND MCGOODER, J.: The effect of tetracycline on the growth of rabbits, *Arch. Biochem.* 37:164, 1952.
425. HUANG, T. C. ULRICH, H. E., AND MCKAY C. M.: Antibiotics, growth, food utilization and use of chromic oxide in studies with rabbits, *Federation Proc.* 13 462, 1954.
426. ANON. Antibiotics in rabbit rations materially reduce losses due to enteritis. (For animals-animal husbandry research) Report of the Chief of the Bureau of Animal Industry Agricultural Research Administration, 45, 1953.
427. LEROUXWONG, P., AND WHITMORE, R. Results of the use of aureomycin-B₁₂ feed supplement, *Small Stock Mag.* 35:29 1951.
428. VIJAYARAGHAVAN, P. K. MURPHY E. A. AND DUNN M. S.: The effect of aureomycin on the growth of mice, *Arch. Biochem.* 36:127 1952.
429. HAMER, D. M.; RAKE, G. MCKEE, C. M., AND MACPHILLANY H. B.: Toxicity of penicillin as prepared for clinical use, *Am. J. M. Sc.* 206:642, 1943.
430. AMERUS, C. M. SIDDI, C. N.; JOHNSON G. C., AND HARRISON J. W. E. Toxicity of penicillin and aureomycin in guinea pigs, *Antib. & Chemot.* 2 521, 1952.
431. HEITMAN, F. R.: Aureomycin in the treatment of experimental relapsing fever and leptospirosis icterohaemorrhagica. (Weil's disease) *Proc. Staff Meet., Mayo Clin.* 23 569 1948.
432. ROINE, P., AND ETTALA, T. Toxicity of aureomycin to guinea pigs, *Nature* 169 1014, 1952.
433. ROINE, P. RAITIO, A., AND VAKKIOVAARA, U.: Listeria infection in the guinea pig caused by feeding with aureomycin, *Nature* 172:767 1953.
434. ROINE, P. Effect of antibiotics on intestinal flora in guinea pigs, *Svensk Kem. Tidskr.* 68 143, 1954.
435. DREAU F. L.: Unpublished observations.

436. HOGAN, A. G.: Personal communication, 1955
437. KOWLER, A. R. MILLER, W. H., AND BROQUIST H. P.: Aureomycin chlortetracycline and the control of poultry spoilage. *Food Tech.* 9:151, 1955.
438. McVAY L. V. JR., AND STUNT D. H.: A long range study of the use of broad spectrum antibiotics in elderly people. In. *Antibiotics Annual 1953-1954*, New York, Medical Encyclopedia, Inc., 1953 p. 273.
439. MARHAM, F. S. PRICE, R. J., SEEGER, K., AND WHITE STEVENS R.: Dietary aureomycin and immune response to Newcastle disease and infectious bronchitis vaccination, *Poultry Sc.* 33:32, 1954



INDEX

Name Index

Including authors cited by name in text

- Abell, M. R., 83, ref. 371
 Ahnquist, H. J., 49 ref. 21
 Ambroz, C., 63, ref. 430
 Anderson, G. W., 35 refs. 91, 120-121
 Andre, F., 65, ref. 271
 Balkom, S. L., 78 ref. 337
 Baer, A., 83, ref. 359
 Barber R. S. 49 63, 83, refs. 163, 358
 Bartley E. E., 65 refs. 273, 274, 279
 Baumann, C. A., 45, ref. 147
 Baxter J. H., 87 ref. 383
 Becker D. E., 63 ref. 227
 Beeson, W. M., 73, ref. 317
 Bell, M. C., 65 ref. 270
 Bell, T. D., 66 69 refs. 275 299 300
 Bentley O. G., 78 ref. 324
 Berg, L. R., 59 77 refs. 214, 328
 Berry M. E., 54, ref. 168
 Beveridge, J. M. R., 85, ref. 371
 Bielby J. 44, 51, refs. 141, 179
 Bird, H. R., 22, 77 refs. 54, 205
 Black, A., 53, refs. 184, 185
 Blaylock, L. G., 35 ref. 124
 Bogdanoff, P. D., Jr., 50 ref. 169
 Bohman, V. R., 71, ref. 318
 Botkins, M. P. 70, refs. 306 309
 Bowland, J. P. 64 ref. 240
 Branton, H. D. 58, 59 64, refs. 35, 206, 216
 Bratzler J. W. 53, ref. 184
 Braude, R., 61, ref. 225
 Bridges, J. H., 70 ref. 306
 Broquist, H. P. 74, 94, refs. 41, 437
 Brown, F. B., 54 ref. 193
 Brown, J. A., 77 ref. 329
 Brown, P. B. 69 ref. 193
 Burgess, R. C., 47 48 49, ref. 23
 Burnside, J. E., 63, 89 refs. 235 403
 Calernick, B. 81, ref. 347
 Cameron, H. S., 80, ref. 409
 Campbell, H., 87 ref. 385
 Campbell, J. A., 49 ref. 102
 Carlson, C. W., 78, ref. 340
 Carpenter K. J. 77 ref. 333
 Carpenter L. E., 21 63, 71, refs. 27 228, 314
 Carroll, R. W. 53, refs. 186, 187
 Carson, J. R., 79 ref. 343
 Carter C. H., 91, ref. 418
 Castro, D. V. 21 47 51, 57 refs. 47 50, 158, 162, 197
 Chalkoff L. L., 81, ref. 350
 Chance, C. M. 83, refs. 109 110
 Chow B. F., 46, ref. 155
 Clark, H. E., 64, ref. 266
 Clawson, A. J. 64, ref. 243
 Coates, M. E., 21, 22, 26, 45 48, 49 refs. 30, 51 52, 60, 67 149
 Colby R. W. 65 69 refs. 287 299
 Common, R. H., 44, ref. 140
 Coodin, F. J. 90, ref. 413
 Cornatzer W. E., 57 ref. 368
 Couch, J. R., 35 ref. 126
 Cravioto-Munoz, J. 49, ref. 157
 Gmba, T. J., 21 62, ref. 49
 Daft, F. S., 45 ref. 148
 Davey R. J. 73, ref. 318
 Déchène, E., 90 ref. 415
 De la Huerga, J. 86 ref. 533
 Deshmukh, P. L., 88, ref. 593
 Dossan, P. L., 93, ref. 435
 Duckworth, J. 77 ref. 333
 Duttonman, W. W. 68, refs. 296, 297
 Dyer L. A., 73, ref. 316
 Elam, J. F., 29 32, 36, 76, 77 refs. 76, 88, 123, 335
 Elliott, R. F., 70 ref. 307
 Ellsworth, S. A., 70 ref. 307
 Emerson, G. A., 44, ref. 189
 Erithoff B. H., 81, refs. 344, 345

- Falcon, W. W., 83, ref. 391
 Fell, R. V., 37, ref. 129
 Ferris, E. F., 41, ref. 134
 Fine, J., 83, ref. 360
 Folkes, J. P., 55, ref. 105
 Forbes, B. M., 54, refs. 184, 191
 Foy H., 83, refs. 395-397
 French, C. E., 4., ref. 138
 Furth, F. W., 84, refs. 365-367
 Gabuten, A. R., 50, ref. 163
 Gabunda, G. J., 92, ref. 421
 Gale, E. F., 16, 55, refs. 9, 103
 Gallo, D. G., 87, ref. 398
 Garrigus U. S., 69, refs. 302-303
 Gordon, H. A., 45, ref. 68
 Gordon, W. S., 87, ref. 130
 Graves, O. E., 89, ref. 408
 Grant, W. C., 81, 82, refs. 49, 354
 Gustafson, G. E., 84, ref. 364
 Gyurgy P., 83, 88, refs. 45, 572
 Haight, T. H., 9., ref. 4.0
 Halbrook, E. R., 77, ref. 331
 Halevy S., 45, ref. 152
 Hamre, D. M., 93, ref. 429
 Hanson, L. E., 41, ref. 134
 Harned, B. K., 17, ref. 3
 Harbuck, E. W., 49, 54, refs. 161, 190
 Hatfield, E. E., 69, ref. 302
 Hellman, F. R., 93, ref. 431
 Hershberger T. V., 76, ref. 324
 Heywang, B. W., 77, 78, refs. 336, 338
 Hubbs, J. W., 33, ref. 108
 Hugh, E. G., 49, ref. 165
 Hull, C. H., 59, refs. 215, 216
 H. H. E. G., 23, ref. 61
 Hullerman, J. P., 59, ref. 218
 Hodgetz, A. W., 69, ref. 298
 Hogan, A. C., 93, ref. 436
 Hopwood, H. H., 90, ref. 416
 Horn, L. H. Jr., 83, ref. 111
 Howland, J. W., 84, ref. 362
 Huang, T. C., 93, ref. 425
 Huhtanen, C. N., 33, ref. 115
 Hummoller F. L., 87, ref. 399
 Jensen, A. H., 52, ref. 183
 Johnson, K. B., 46, refs. 87, 156
 Johnson, B. C., 46, 47, 54, 82, 83, refs. 154, 160, 190
 Johnson, E. L., 77, ref. 334
 Johnson, R., 90, ref. 412
 Jordan, R. M., 66, 69, refs. 275, 299-301
 Jukes, T. H., 44, ref. 143
 Kaplan, A., 87, ref. 384
 Kaya, R., 90, ref. 414
 Kelly J. W., 59, ref. 215
 Keiler E. M., 82, 33, ref. 107
 Kensler F. B., 68, refs. 296, 297
 King, J. O. L., 22, ref. 58
 Kinsman, D. M., 71, ref. 312
 Knodt, C. B., 32, 33, 60, ref. 107, 276
 Kohler A. R., 74, 91, refs. 41, 437
 Kohlmeier W., 78, ref. 340
 Koletsky S., 81, ref. 364
 Kondi, A., 83, refs. 395-397
 Kratzer F. H., 99, 34, refs. 73, 88
 Krider J. L., 21, ref. 49
 Kunkel, H. O., 74, refs. 310, 311
 Kunsisto, A. N., 82, ref. 331
 Lampman, C. E., 77, ref. 350
 Larson, N., 71, ref. 314
 Larson N. L., 23, ref. 61
 Lassiter C. W., 87, 66, ref. 196
 Lawrence J. M., 92, ref. 424
 Lepine, P., 25, ref. 67
 Lepper M. H., 87, ref. 388
 Leerwenburg, P., 93, ref. 427
 Lobby D. A., 59, 81, 95, refs. 133, 135, 348
 Lichtman, H., 88, ref. 394
 Lih, H., 45, ref. 147
 Lillie, R. J., 77, ref. 205
 Lindblad, G. S., 50, ref. 170
 Linkswiler H., 45, ref. 148
 Lodge, J. B., 83, ref. 114
 Loosli, J. K., 32, 65, refs. 105, 272
 Luckey T. D., 28, 29, refs. 44, 71
 McDaniel, E. C., 45, ref. 148
 Macdougall, L., 91, ref. 419
 McGinnis, J., 92, ref. 424
 Machlin, L. J., 50, ref. 172
 McVay L. V. Jr., 94, ref. 438
 Maddock, H. M., 74, ref. 319
 Maghrabi, M. G., 82, ref. 352
 March, B., 44, ref. 141
 Mariakulandia, A., 77, ref. 3.1
 Markham F. S., 95, ref. 439
 Marti-Ibáñez, Félix, 7
 Martin, G. J., 17, ref. 21
 Matsushima J., 68, ref. 294
 Matterson, L. D., 51, 59, ref. 34, 184
 Maurer S., 48, ref. 29
 Mayfield, O. J., 17, ref. 22
 Meade R. J., 54, refs. 191, 19.2
 Meltes, J., 81, 82, refs. 346, 348, 353, 359
 Mellen, W. J., 82, ref. 353
 Migicovsky B. B., 50, ref. 167
 Milby T. T., 77, ref. 33.2
 Miller C. P., 81, ref. 363
 Molitor H., 89, ref. 409
 Monson, W. F., 47, ref. 40
 Moore P. B., 17, ref. 2
 Morehouse, N. F., 17, ref. 22

- Morrison, A. B., 22, ref. 59
 Murray T. K., 49, ref. 163
 Murthy M. R. V., 92, ref. 423
 Nelson, T. S., 45, ref. 150
 Neumann, A. L., 82, ref. 106
 Noland, P. R., 63, 64, refs. 96, 244
 Ogle, R. C., 81, ref. 346
 Oleson, J. J., 44, ref. 143
 Olsson, N., 83, ref. 402
 Pankas, L., 70, ref. 303, 309
 Pepper W. F., 26, 42, 50, refs. 68, 187, 171
 Perrini, F., 90, ref. 410
 Perry T. W., 64, 68, refs. 245, 263, 295
 Peterson, C. F., 77, ref. 330
 Pierce, W. E., 92, ref. 420
 Radisson, J. J., 83, 34, ref. 112
 Richardson, C. W., 67, ref. 286
 Richardson, D., 41, ref. 135
 Riddell, W. H., 71, ref. 312
 Roberts, B. S., 85, ref. 123
 Robinson, K. L., 63, ref. 236
 Robinson, P., 90, ref. 411
 Rotne, P., 93, refs. 432-434
 Romero J. L., 35, ref. 119
 Ross, E., 49, ref. 166
 Roy J. H. B., 67, ref. 287
 Rusoff, L. L., 83, 65, 89, refs. 98, 113, 265, 278, 280-285
 Russo, J. M., 54, 55, ref. 194
 Sarett, H. P., 45, ref. 151
 Samberlich, H. E., 46, ref. 153
 Saxena, H. C., 51, 59, refs. 176, 177
 Sborov V. M., 87, ref. 890
 Schaffke, P. J., 89, 95, refs. 133, 185
 Schendel, H. E., 46, ref. 154
 Schmuck, C., 54, ref. 183
 Scott, H. M., 45, 51, refs. 150, 175
 Scott, M. L., 35, ref. 125
 Scribshaw N. S., 91, ref. 417
 Seto, J. T., 87, ref. 368
 Shaffner C. S., 50, refs. 168, 169
 Sheffy B. E., 47, 89, ref. 159
 Sherwood, D. H., 77, ref. 338
 Stoburth, J. McN., 29, refs. 72, 75
 Stinsen, E. P., 58, ref. 84
 Stinson, J. R., 74, refs. 42, 322
 Stinger S. J., 42, 51, 76, refs. 187, 178, 823
 Smith, D. G., 44, ref. 139
 Smyser C. F. Jr., 82, ref. 103
 Snelling, C. E., 90, ref. 412
 Speer V. C., 23, ref. 48
 Sprunt, D. H., 94, ref. 438
 Sreenivasaya, M., 92, ref. 423
 Stephenson, E. L., 87, ref. 129
 Stokstad, E. L. R., 44, ref. 143
 Sunde, M. L., 77, ref. 327
 Sutherland, D. A., 87, 88, refs. 360, 362
 Swenson, M. J., 64, ref. 246
 Tarver F. H., Jr., 64, ref. 261
 Tatrog, A., 81, ref. 350
 Taylor J. H., 37, 92, ref. 180, 422
 Telika, A., 82, ref. 351
 Terrill, S. W., 21, ref. 49
 Tovrea, E., 68, ref. 293
 Tribe, D. E., 89, ref. 399
 Turner, A. W., 69, ref. 298
 Turner C. W., 82, ref. 352
 Umbreit, W. W., 16, ref. 14
 Ungley C. C., 89, ref. 401
 Vostal, C. M., 64, 73, refs. 262, 315
 duVigneaud, V., 36, ref. 137
 Vijayaraghavan, P. K., 93, ref. 428
 Waibel, P. E., 83, 44, 45, 74, 77, refs. 132, 144, 145
 Wahlstrom, R. C., 47, 63, 82, 83, refs. 100, 229
 Wallace, H. D., 64, 65, refs. 247, 272
 Walker, E. F., 82, ref. 353
 Watson, C. M., 89, ref. 400
 Weakley C. E., Jr., 50, ref. 178
 Welch, Henry 7
 Wetzel, N. D., 90, ref. 416
 Whitt, E. G., 22, ref. 58
 White-Stevens, R., 51, 59, 78, refs. 181, 224, 839, 842
 Whitmore, R., 93, ref. 427
 Williams, W. L., 29, 87, 63, refs. 74, 131, 250
 Wilson, J. E., 86, ref. 127
 Witte, L. J., 89, ref. 400
 Wong, S. C., 74, ref. 320
 Yacowitz, H., 43, ref. 166
 Zefbel, H. G., 51, 59, 78, refs. 181, 224, 339
 Zimmerman, H. H., 87, ref. 359

Subject Index

- Anemia
- treatment with antibiotics, 88-89
- Animals - *See also under names of animals*
- antibiotic feeding in animal production, 57 79
 - conclusions, 85-90
 - germ-free:
 - effect of feeding antibiotics 26-29
 - reproduction:
 - effect of antibiotic feeding, 71 79
- Antibiotics
- effect on bacterial metabolism, 18-16
 - on blood formation, 88-89
 - on choline and methionine deficiency, 85
 - on egg production in hens, 74-79
 - on intestinal bacteria, 19-20
 - on intestinal flora, 29-32
 - on liver physiology 84-88
 - on manganese metabolism, 50
 - on nitrogen metabolism, 53-55
 - on nutrient requirements, 44-50
 - on protein requirement, 50-53
 - on reproduction in animals, 71 79
 - on rumen microorganisms, 82-84
 - on ruminants, 63-71
 - on stress conditions, 83
 - on vitamin requirements, 44-50
 - vitamin A, 47-49
 - vitamin B complex, 44-47
 - vitamin D 49-50
 - feeding in animal production, 57 79
 - conclusions, 85-90
 - growth effect:
 - antibiotic-tolerant intestinal microorganisms as a source of unidentified growth factors, 84-90
 - effect of clean and dirty environment on, 22-24
 - effect of prolonged use of antibiotic feeding 38-42
 - in germ-free animals, 26-29
 - in human subjects, 90-92
 - mechanism of 13 17 18 19-42
 - on chick embryos, 24-25
 - morphologic changes produced by in chicks, 25-28
 - pharmacology 89-90
 - physiologic effects of antibiotic feeding, 81-89
 - production of, 15
 - therapeutic use in anemia, 88-89
- Aureomycin - *See Chlortetracycline*
- Bacitracin
- growth effect
 - not active when injected, 20
 - structural formula of bacitracin A, 15
- Bacteria
- effect of antibiotics on metabolism, 18-16
 - effect of chloramphenicol, 16
 - effect of chlortetracycline, 16
 - effect of penicillin, 16
 - effect of streptomycin, 16
- intestinal
- effects of antibiotics on, 17 18, 19-20
- Bibliography 97 119
- Blood
- effect of antibiotics on blood formation, 88-89
- Calcium
- effect on antibiotic feeding on requirements, 49-50
- Calves
- effect of chlortetracycline feeding on growth, 66-67
- Carotene
- effect of antibiotic feeding on carotene requirements, 47-49

Cattle

- effect of antibiotic feeding in beef cattle, 67-69
- chlortetracycline, 67-69 70

Chick Embryos

- effect of antibiotics on growth, 24-25

Chickens - See also Poultry

- effect of antibiotic feeding on protein requirements, 50-51
- effect of hydrolyzed penicillin on growth, 36-38
- effect of prolonged use of antibiotic feeding, 88-91
- morphologic changes produced by feeding antibiotics, 25-26
- respiratory disease:
 - antibiotic feeding in, 59-61
 - chlortetracycline, 59-61

Children

- growth effect of antibiotics on, 90-92

Chloramphenicol

- effect on bacterial metabolism, 16

Chlortetracycline

- comparison with colostrum in feeding calves, 67
- effect on bacterial metabolism, 16
 - on chickens with chronic respiratory disease, 59-61
 - on egg production in hens 78-79
 - on guinea pigs, 93
 - on intestinal bacteria, 17 18
 - on nitrogen metabolism, 53-55
 - on protein requirements, 51-53
 - on rabbits, 62-63
 - on reproduction in pigs, 72-74
 - on rumen microorganisms, 32-34
 - on the thyroid gland, 81-82
 - on vitamin A requirements, 49
 - on vitamin B complex requirements, 44-47
 - vitamin B₁₂, 46-47

fermentation residues:

- in feeding pigs, 62-63

growth effect:

- active when injected, 20
- effect of environment on, 22-23, 24
- on beef cattle, 68-69 70
- on dairy calves, 66-67 68
- on horses, 62
- on lambs, 69-70
- on mice, 93
- on pigs, 63
- on premature infants, 90
- on silkworms, 92
- in prevention of liver necrosis, 85-86

- protective against alloxan diabetes in rats, 83

- structural formula, 14

therapeutic uses:

- in diseases of the liver in man, 86

toxicity:

- nontoxic on prolonged administration to human subjects, 94-95

Choline

- deficiency in diet, 84-85
- effect of antibiotics on, 85

Colostrum

- comparison with chlortetracycline in feeding calves, 67

Cows

- no effect of chlortetracycline on mature lactating cows, 67

Diabetes Mellitus

experimental:

- chlortetracycline protective against alloxan diabetes in rats, 83

Eggs

- effect of antibiotic feeding of the hen, 74-78
- increasing hatchability 74-76

Endocrine Glands

- effect of antibiotic feeding, 81-84

Environment

- effect of clean and dirty environment on antibiotic growth effect, 22-24

Foreword, 7 11

Guinea Pigs

- chlortetracycline not toxic for 93
- penicillin toxic for, 93

Hens

egg production-See also Eggs

- effect of antibiotic feeding on, 74-78

respiratory disease:

- effect of antibiotic feedings on egg production by hens with chronic respiratory disease, 78-79

Horses

- effect of chlortetracycline on, 62

Infants

premature:

- effect of chlortetracycline on, 90

Intestines

flora - See also Bacteria, intestinal.

- antibiotic-tolerant intestinal microorganisms as a source of unidentified growth factor 34-38
- effect of antibiotic feeding on, 29-32

Lambs

- effect of antibiotic feeding, 69-70
- chlortetracycline, 69-70

- Liver
 - effect of antibiotics on, 81-88
- diarrhea:
 - treatment with chlortetracycline in man, 86
- fat:
 - effect of antibiotics on, 87-88
- necrosis
 - beneficial effects of antibiotics, 21
 - prevention by chlortetracycline 85-86
- Manganese
 - effect of antibiotic feeding on manganese metabolism, 50
- Meat
 - effect of antibiotic feeding on quality of 64-65 91-95
- Methionine
 - deficiency in diet, 84-85
 - effect of antibiotics on, 85
- Milk
 - effect of chlortetracycline on, 83
- Nitrogen Metabolism
 - effect of antibiotic feeding on, 53-55
 - chlortetracycline 53-55
 - oxytetracycline, 54
- Nutrition
 - effect of feeding antibiotics on the requirement for nutrients, 43-55
- Oxytetracycline
 - effect on nitrogen metabolism, 54
 - on rabbits, 92-93
 - growth effect.
 - in germ-free poultry 26-29
 - structural formula, 14
- Penicillin
 - effect on bacterial metabolism, 16
 - on vitamin A requirements, 49
 - on vitamin B complex requirements, 44-45
 - on vitamin D requirements, 49-50
 - growth effect.
 - active when injected, 20
 - effect of environment on, 23-24
 - growth of animals receiving hydrolyzed penicillin, 36-38
 - structural formula, 14
 - toxicity:
 - toxic for guinea pigs, 93
- Phosphorus
 - effect of antibiotic feeding on requirements, 50
- Pigs
 - effect of antibiotic feeding on growth, 57-58 81-84
 - chlortetracycline, 83
 - effect of antibiotic feeding on protein requirements 51-53
 - effect of chlortetracycline on nitrogen metabolism, 54-55
 - effect of hydrolyzed penicillin on growth, 37
 - effect of prolonged use of antibiotic feeding, 40-42
 - reproduction.
 - effect of antibiotic feeding, 72-74
 - chlortetracycline, 72-74
- Poultry
 - effect of antibiotic feeding on growth, 58-59
- germ free
 - effect of feeding antibiotics, 26-29
- Protein
 - effect of antibiotic feeding on requirements, 50-53
 - chlortetracycline, 51-53
- Public Health
 - effect of antibiotic feeding of animals, 94-95
- Rabbits
 - effect of chlortetracycline and oxytetracycline on, 92-93
- Rumen
 - microorganisms-
 - effect of antibiotic feeding on, 32-34
- Ruminants
 - effect of antibiotic feeding, 65-71
- Silkworms
 - effect of chlortetracycline on, 92
- Streptomyces aureofaciens*
 - fermentation products:
 - effect on animal growth, 18, 21
- Streptomycin
 - effect on bacterial metabolism, 16
 - on intestinal bacteria, 17
 - on nitrogen metabolism, 53
 - growth effect-
 - not active when injected, 20
 - structural formula, 14
- Stress
 - effect of antibiotic feeding on stress conditions, 83
- Sulfonamides
 - in animal feeding, 16-17
- Terramycin - See *Oxytetracycline*
- Thyroid Gland
 - effect of chlortetracycline on, 81-82
- Vitamin A
 - effect of antibiotics on requirements, 47-49

— chlortetracycline, 49

— penicillin, 49

Vitamin B Complex

— effect of antibiotic feeding on requirements for 44-47

— effect of chlortetracycline, 44-47

— effect of penicillin, 44-45

Vitamin B₁₂

— effect of chlortetracycline on requirements, 46-47

— role in choline and methionine deficiency 85

Vitamin D

— effect of antibiotic feeding on requirements, 49-50

— penicillin, 49-50

Vitamins

— deficiency produced by sulfonamide 16-17

— effect of antibiotics on requirements 44-50

Antibiotics in nutrition

by Thomas H. Jukes, Ph.D.

Research Division AMERICAN CYANAMID COMPANY
NEW YORK

Foreword by HENRY WELCH, M.D. AND
FELIX MARTI IBAÑEZ, M.D.

A Publication of

CALIFORNIA ENCYCLOPEDIA, INC.
30 EAST 60th STREET NEW YORK 22, N. Y.